

Table 1.15C Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Numbers in Thousands, 2011 and 2012

Dependence or Abuse Status	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)
DEPENDENCE				
None	761	809	300	321
Both Illicit Drugs and Alcohol	87	105	63	70
Illicit Drugs or Alcohol	206	261	122	151
Illicit Drugs	127	165	86	98
Alcohol	182	223	107	135
DEPENDENCE OR ABUSE				
None	737	781	288	310
Both Illicit Drugs and Alcohol	99	132	66	80
Illicit Drugs or Alcohol	251	311	145	168
Illicit Drugs	136	180	91	104
Alcohol	229	277	130	154

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.15D Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Percentages, 2011 and 2012

Dependence or Abuse Status	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)
DEPENDENCE				
None	0.30	0.31	0.13	0.14
Both Illicit Drugs and Alcohol	3.69	3.99	4.61	4.58
Illicit Drugs or Alcohol	1.44	1.50	1.05	1.16
Illicit Drugs	2.14	2.21	1.93	1.88
Alcohol	1.75	1.85	1.27	1.44
DEPENDENCE OR ABUSE				
None	0.31	0.31	0.13	0.14
Both Illicit Drugs and Alcohol	2.75	2.78	2.58	2.66
Illicit Drugs or Alcohol	1.09	1.12	0.71	0.76
Illicit Drugs	1.79	1.86	1.53	1.47
Alcohol	1.21	1.25	0.77	0.86

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.15P Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Dependence or Abuse Status	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)
DEPENDENCE				
None	0.3199		0.7852	
Both Illicit Drugs and Alcohol	0.4466		0.7472	
Illicit Drugs or Alcohol	0.0185		0.3167	
Illicit Drugs	0.4031		0.8578	
Alcohol	0.0193		0.2720	
DEPENDENCE OR ABUSE				
None	0.5009		0.9058	
Both Illicit Drugs and Alcohol	0.1056		0.8018	
Illicit Drugs or Alcohol	0.0034		0.1839	
Illicit Drugs	0.2288		0.9470	
Alcohol	0.0034		0.1272	

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2011 and 2012

Dependence or Abuse Status	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
DEPENDENCE												
Both Illicit Drugs and Alcohol	96	120	87	105	63	70	37	53	44	46	39	54
Illicit Drugs or Alcohol	286	349	206	261	122	151	87	126	126	145	211	242
Illicit Drugs	170	211	127	165	86	98	60	80	75	93	113	131
Alcohol	257	297	182	223	107	135	79	110	111	116	187	210
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	120	158	99	132	66	80	47	64	55	74	71	83
Illicit Drugs or Alcohol	401	463	251	311	145	168	107	141	164	196	349	371
Illicit Drugs	195	252	136	180	91	104	65	84	83	108	138	171
Alcohol	374	412	229	277	130	154	96	129	152	177	331	335

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2011 and 2012

Dependence or Abuse Status	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.04	0.05	0.21	0.24	0.68	0.72	0.36	0.44	0.20	0.21	0.02	0.03
Illicit Drugs or Alcohol	0.12	0.15	0.48	0.56	1.24	1.43	0.83	1.02	0.56	0.63	0.11	0.12
Illicit Drugs	0.07	0.09	0.30	0.37	0.91	1.00	0.58	0.66	0.34	0.41	0.06	0.07
Alcohol	0.11	0.13	0.42	0.49	1.11	1.29	0.76	0.91	0.49	0.51	0.10	0.11
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	0.05	0.07	0.24	0.29	0.71	0.82	0.45	0.53	0.25	0.33	0.04	0.04
Illicit Drugs or Alcohol	0.17	0.20	0.57	0.65	1.40	1.54	1.01	1.13	0.72	0.82	0.17	0.18
Illicit Drugs	0.08	0.11	0.33	0.39	0.95	1.05	0.63	0.69	0.38	0.47	0.07	0.09
Alcohol	0.16	0.18	0.52	0.59	1.31	1.43	0.91	1.04	0.67	0.76	0.17	0.17

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Dependence or Abuse Status	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.4172		0.4635		0.9931		0.2919		0.7308		0.9567	
Illicit Drugs or Alcohol	0.0050		0.0036		0.0943		0.0089		0.2779		0.5481	
Illicit Drugs	0.0289		0.1244		0.6705		0.2253		0.2617		0.2963	
Alcohol	0.0409		0.0124		0.1257		0.0099		0.5747		0.9725	
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	0.1799		0.1393		0.5796		0.3646		0.2695		0.8595	
Illicit Drugs or Alcohol	0.0099		0.0013		0.0776		0.0247		0.0732		0.7145	
Illicit Drugs	0.0059		0.0304		0.4689		0.2292		0.0611		0.1699	
Alcohol	0.0824		0.0053		0.0910		0.0334		0.2031		0.7471	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2011 and 2012

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	195	252	136	180	91	104	65	84	83	108	138	171
18-25	83	94	51	62	27	30	23	32	36	40	72	77
26-49	145	197	103	147	74	87	53	73	55	84	98	126
50 or Older	93	123	71	79	43	57	30	27	48	48	62	94
MALE	161	210	100	138	54	81	50	64	67	88	124	157
18-25	64	76	35	47	17	21	18	23	27	32	61	68
26-49	121	166	76	114	48	70	40	56	47	68	92	118
50 or Older	76	114	51	65	*	*	*	22	39	38	58	93
FEMALE	107	119	92	100	72	64	39	56	48	53	54	60
18-25	47	52	37	42	23	21	16	24	23	25	32	34
26-49	80	94	71	79	57	52	32	47	32	36	37	45
50 or Older	54	48	49	46	38	32	*	16	27	29	22	13

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2011 and 2012

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	0.08	0.11	0.33	0.39	0.95	1.05	0.63	0.69	0.38	0.47	0.07	0.09
18-25	0.24	0.27	0.71	0.81	1.86	1.85	1.32	1.49	0.95	1.06	0.25	0.26
26-49	0.15	0.20	0.51	0.67	1.39	1.62	0.99	1.26	0.57	0.80	0.13	0.16
50 or Older	0.09	0.12	0.46	0.48	1.52	1.80	0.86	0.59	0.53	0.55	0.07	0.11
MALE	0.14	0.19	0.61	0.78	1.64	2.11	1.38	1.35	0.73	1.00	0.13	0.16
18-25	0.37	0.44	1.26	1.50	3.45	3.52	2.81	2.83	1.60	1.95	0.39	0.43
26-49	0.25	0.35	0.96	1.28	2.59	3.16	2.09	2.36	1.16	1.62	0.23	0.29
50 or Older	0.16	0.24	0.85	1.06	*	*	*	1.22	1.07	1.23	0.14	0.22
FEMALE	0.09	0.10	0.36	0.38	1.17	1.07	0.59	0.77	0.37	0.39	0.06	0.06
18-25	0.28	0.30	0.87	0.96	2.35	2.09	1.40	1.88	1.12	1.19	0.23	0.25
26-49	0.16	0.19	0.58	0.62	1.66	1.66	0.93	1.38	0.55	0.58	0.10	0.12
50 or Older	0.10	0.09	0.53	0.45	2.20	1.59	*	0.59	0.52	0.52	0.05	0.03

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	0.0059		0.0304		0.4689		0.2292		0.0611		0.1699	
18-25	0.3932		0.2941		0.1573		0.2510		0.1029		1.0000	
26-49	0.0049		0.0209		0.2513		0.1334		0.1031		0.1428	
50 or Older	0.4911		0.9425		0.6583		0.7283		0.8190		0.4380	
MALE	0.0206		0.0163		0.2665		0.9144		0.0183		0.4071	
18-25	0.9086		0.2163		0.4211		0.3334		0.1501		0.2919	
26-49	0.0178		0.0473		0.4853		0.5907		0.0530		0.2464	
50 or Older	0.2765		0.5450		*		*		0.7645		0.3411	
FEMALE	0.0943		0.5066		0.8784		0.1667		0.7706		0.0963	
18-25	0.0903		0.8071		0.2308		0.6334		0.3253		0.0484	
26-49	0.1084		0.2322		0.4791		0.1587		0.8116		0.2518	
50 or Older	0.5585		0.6108		0.6291		*		0.5050		0.4682	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2011 and 2012

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	374	412	229	277	130	154	96	129	152	177	331	335
18-25	116	114	72	81	31	35	35	44	50	51	120	110
26-49	259	288	175	215	100	113	75	106	122	140	229	219
50 or Older	238	250	129	143	78	98	56	61	78	86	203	222
MALE	300	349	165	211	82	128	80	107	117	125	271	303
18-25	83	88	48	51	20	23	22	30	37	34	85	87
26-49	214	238	122	165	59	92	54	86	94	102	199	196
50 or Older	194	230	97	117	*	*	*	55	62	63	168	210
FEMALE	219	216	153	157	101	87	60	76	96	108	166	147
18-25	79	72	52	59	23	26	28	31	33	38	73	57
26-49	160	159	124	122	80	64	52	63	78	84	106	106
50 or Older	127	118	76	83	58	52	14	27	48	59	102	85

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2011 and 2012

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	0.16	0.18	0.52	0.59	1.31	1.43	0.91	1.04	0.67	0.76	0.17	0.17
18-25	0.34	0.33	0.91	0.93	1.96	2.00	1.84	1.91	1.23	1.27	0.35	0.33
26-49	0.26	0.29	0.80	0.90	1.77	1.90	1.35	1.72	1.16	1.25	0.28	0.27
50 or Older	0.24	0.24	0.84	0.86	2.70	2.88	1.58	1.33	0.87	0.98	0.23	0.25
MALE	0.27	0.31	0.97	1.15	2.35	2.93	2.08	2.14	1.22	1.38	0.27	0.30
18-25	0.49	0.51	1.56	1.54	3.67	3.56	3.22	3.30	2.03	2.01	0.50	0.50
26-49	0.44	0.50	1.45	1.64	3.09	3.68	2.76	3.29	2.11	2.23	0.46	0.47
50 or Older	0.42	0.48	1.58	1.92	*	*	*	3.00	1.67	1.96	0.41	0.49
FEMALE	0.18	0.18	0.57	0.57	1.57	1.35	0.88	1.03	0.73	0.77	0.17	0.15
18-25	0.46	0.42	1.14	1.17	2.34	2.42	2.19	2.20	1.53	1.63	0.48	0.40
26-49	0.32	0.32	0.94	0.91	2.19	1.88	1.47	1.76	1.27	1.29	0.28	0.28
50 or Older	0.24	0.21	0.82	0.80	3.23	2.49	0.65	0.99	0.90	1.06	0.23	0.19

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	0.0824		0.0053		0.0910		0.0334		0.2031		0.7471	
18-25	0.8871		0.2641		0.3261		0.6820		0.5935		0.1912	
26-49	0.1199		0.0047		0.0907		0.0061		0.3883		0.4748	
50 or Older	0.1829		0.3977		0.6702		0.6839		0.5933		0.3468	
MALE	0.0246		0.0011		0.0306		0.0832		0.1131		0.6914	
18-25	0.5363		0.5605		0.9952		0.1835		0.7776		0.2126	
26-49	0.1240		0.0052		0.1203		0.0156		0.3076		0.6353	
50 or Older	0.0279		0.1112		*		*		0.2973		0.0932	
FEMALE	0.7928		0.5055		0.9737		0.4599		0.5954		0.1515	
18-25	0.6183		0.3332		0.2331		0.5068		0.2569		0.6022	
26-49	0.6491		0.3148		0.5707		0.2517		0.7605		0.6181	
50 or Older	0.2379		0.6917		0.3407		0.1788		0.8702		0.1620	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.19C Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2011 and 2012

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	401	463	251	311	145	168	107	141	164	196	349	371
18-25	128	133	85	93	38	40	39	49	59	59	135	134
26-49	281	321	194	240	111	126	84	116	130	151	239	240
50 or Older	251	269	139	160	85	107	59	63	86	96	214	236
MALE	321	383	182	230	88	135	86	113	130	140	287	335
18-25	92	105	56	63	23	27	24	34	44	43	98	108
26-49	232	261	138	181	66	101	61	92	104	112	208	216
50 or Older	204	243	102	125	*	*	*	56	67	70	180	225
FEMALE	230	241	166	181	114	101	67	88	101	117	172	156
18-25	83	78	59	64	29	30	31	35	36	41	79	64
26-49	171	178	133	138	89	74	56	72	80	87	111	113
50 or Older	135	126	87	94	66	61	21	29	54	66	105	86

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.19D Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2011 and 2012

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	0.17	0.20	0.57	0.65	1.40	1.54	1.01	1.13	0.72	0.82	0.17	0.18
18-25	0.37	0.39	1.00	1.00	2.20	2.10	1.92	1.93	1.35	1.41	0.38	0.39
26-49	0.29	0.33	0.88	0.98	1.89	2.08	1.50	1.82	1.23	1.32	0.29	0.30
50 or Older	0.25	0.26	0.90	0.95	2.91	3.09	1.68	1.37	0.94	1.07	0.25	0.27
MALE	0.29	0.34	1.05	1.24	2.52	2.98	2.24	2.27	1.35	1.53	0.29	0.33
18-25	0.54	0.61	1.65	1.73	3.99	3.81	3.36	3.26	2.23	2.32	0.56	0.60
26-49	0.48	0.54	1.59	1.73	3.28	3.82	3.03	3.42	2.28	2.39	0.48	0.51
50 or Older	0.44	0.51	1.65	2.03	*	*	*	3.05	1.79	2.16	0.44	0.52
FEMALE	0.19	0.20	0.62	0.65	1.72	1.57	1.00	1.17	0.78	0.84	0.18	0.16
18-25	0.48	0.45	1.20	1.24	2.59	2.60	2.34	2.37	1.62	1.75	0.50	0.43
26-49	0.34	0.36	1.01	1.02	2.39	2.19	1.58	1.96	1.31	1.34	0.29	0.30
50 or Older	0.25	0.23	0.93	0.90	3.63	2.86	0.95	1.06	1.00	1.17	0.23	0.19

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.19P Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
TOTAL	0.0099		0.0013		0.0776		0.0247		0.0732		0.7145	
18-25	0.5307		0.1272		0.7470		0.2577		0.1645		0.5361	
26-49	0.0123		0.0010		0.0311		0.0045		0.2471		0.9052	
50 or Older	0.1753		0.3912		0.6344		0.8010		0.5308		0.3493	
MALE	0.0030		0.0002		0.0182		0.1353		0.0166		0.3459	
18-25	0.9213		0.1723		0.7525		0.0683		0.4838		0.4208	
26-49	0.0200		0.0016		0.0711		0.0326		0.1205		0.7826	
50 or Older	0.0220		0.0883		*		*		0.1933		0.0896	
FEMALE	0.8332		0.4153		0.9537		0.2289		0.6657		0.2909	
18-25	0.3783		0.4164		0.8281		0.9128		0.1550		0.9891	
26-49	0.3315		0.1748		0.3065		0.1089		0.8802		0.8959	
50 or Older	0.2172		0.6418		0.4015		0.3097		0.7702		0.1301	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.