

Table 1.31C Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2011 and 2012

Location of Treatment/Counseling ¹	Total		Any	Any	Serious	Serious	Moderate	Moderate	Mild	Mild	No	No
	(2011)	(2012)	Mental Illness (2011)	Mental Illness (2012)								
Outpatient Mental Health Clinic or Center	221	211	189	166	125	108	117	100	91	80	108	122
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	330	343	242	248	148	151	120	146	164	133	201	226
Doctor's Office - Not Part of a Clinic	186	206	164	167	109	99	88	96	82	89	104	117
Outpatient Medical Clinic	129	107	92	94	60	67	52	45	47	48	91	64
Partial Day Hospital or Day Treatment Program	59	63	54	61	44	48	19	31	25	18	23	16
School or University Setting/Clinic/Center ²	14	19	13	17	7	6	7	14	8	6	8	9
Some Other Place ³	63	50	52	39	45	34	17	16	20	12	34	31

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.31D Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2011 and 2012

Location of Treatment/Counseling ¹	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
Outpatient Mental Health Clinic or Center	1.26	1.20	1.67	1.47	2.68	2.56	3.52	2.82	2.45	2.48	1.80	1.89
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	1.39	1.42	1.67	1.77	2.63	2.82	3.49	3.26	3.07	3.11	2.25	2.47
Doctor's Office - Not Part of a Clinic	1.08	1.17	1.44	1.52	2.33	2.34	2.93	2.80	2.23	2.66	1.71	1.85
Outpatient Medical Clinic	0.79	0.67	0.90	0.93	1.47	1.71	1.87	1.47	1.41	1.57	1.55	1.09
Partial Day Hospital or Day Treatment Program	0.37	0.40	0.54	0.62	1.09	1.27	0.70	1.00	0.77	0.62	0.41	0.28
School or University Setting/Clinic/Center ²	0.10	0.12	0.13	0.17	0.17	0.17	0.28	0.46	0.24	0.22	0.15	0.16
Some Other Place ³	0.41	0.32	0.53	0.40	1.16	0.91	0.63	0.51	0.63	0.42	0.62	0.54

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.31P Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Location of Treatment/Counseling ¹	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
Outpatient Mental Health Clinic or Center	0.9167		0.6414		0.5892		0.2302		0.8498		0.5447	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.7830		0.7088		0.8216		0.3117		0.1926		0.9817	
Doctor's Office - Not Part of a Clinic	0.7609		0.7103		0.4595		0.9088		0.1963		0.9692	
Outpatient Medical Clinic	0.3169		0.9045		0.6372		0.4403		0.9930		0.1965	
Partial Day Hospital or Day Treatment Program	0.4920		0.3800		0.3832		0.5407		0.9046		0.7462	
School or University Setting/Clinic/Center ²	0.3588		0.2015		0.7746		0.0799		0.5598		0.7149	
Some Other Place ³	0.1356		0.1219		0.2163		0.9384		0.2317		0.7031	

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.