

Table 1.51C Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Type of Illicit Drug Use in the Past Year: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
ILLCIT DRUGS¹	161	176	90	94	54	58
Marijuana and Hashish	120	149	65	81	39	52
Cocaine	61	76	40	46	18	30
Crack	27	43	20	17	12	25
Heroin	21	*	12	*	9	5
Hallucinogens	55	57	42	37	28	21
LSD	19	28	14	13	10	11
PCP	*	*	*	*	*	*
Ecstasy	50	47	39	29	25	17
Inhalants	32	47	25	24	6	14
Nonmedical Use of Psychotherapeutics ^{2,3}	124	134	63	83	50	46
Pain Relievers	107	119	51	73	36	35
OxyContin [®]	40	56	23	43	13	16
Tranquilizers	85	91	43	57	32	26
Stimulants ³	54	72	32	59	28	34
Methamphetamine ³	44	59	29	47	28	32
Sedatives	*	27	13	9	12	13
ILLCIT DRUGS OTHER THAN MARIJUANA¹	130	142	74	85	49	49

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.51D Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Type of Illicit Drug Use in the Past Year: Standard Errors of Percentages, 2011 and 2012

Drug	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
ILLCIT DRUGS¹	0.45	0.46	0.26	0.25	0.16	0.16
Marijuana and Hashish	0.43	0.50	0.25	0.29	0.15	0.19
Cocaine	1.55	1.62	1.10	1.01	0.49	0.67
Crack	4.18	4.80	3.36	1.96	1.97	2.79
Heroin	3.69	*	2.12	*	1.66	0.87
Hallucinogens	1.49	1.40	1.18	0.98	0.80	0.57
LSD	2.61	2.87	1.92	1.39	1.36	1.24
PCP	*	*	*	*	*	*
Ecstasy	2.28	1.88	1.87	1.22	1.23	0.74
Inhalants	2.83	4.08	2.35	2.27	0.55	1.36
Nonmedical Use of Psychotherapeutics ^{2,3}	0.90	0.84	0.48	0.55	0.39	0.31
Pain Relievers	1.04	1.00	0.52	0.65	0.38	0.32
OxyContin [®]	2.67	3.67	1.60	3.07	0.95	1.25
Tranquilizers	1.69	1.50	0.92	0.99	0.69	0.47
Stimulants ³	2.08	2.18	1.29	1.86	1.14	1.12
Methamphetamine ³	4.36	4.65	3.03	4.03	2.97	2.85
Sedatives	*	4.92	3.04	1.78	2.93	2.56
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.74	0.71	0.44	0.44	0.30	0.26

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.51P Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Type of Illicit Drug Use in the Past Year: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
ILLCIT DRUGS¹	0.2826		0.3594		0.3316	
Marijuana and Hashish	0.0130		0.0421		0.0344	
Cocaine	0.4403		0.8627		0.2271	
Crack	0.3727		0.4422		0.4034	
Heroin	*		*		0.4225	
Hallucinogens	0.8196		0.9012		0.4967	
LSD	0.8056		0.6704		0.5914	
PCP	*		*		*	
Ecstasy	0.9490		0.6247		0.7346	
Inhalants	0.3360		0.7043		0.0687	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.7206		0.3328		0.6692	
Pain Relievers	0.9037		0.4227		0.7045	
OxyContin [®]	0.5860		0.4616		0.8917	
Tranquilizers	0.5169		0.7099		0.3636	
Stimulants ³	0.2763		0.1199		0.6545	
Methamphetamine ³	0.1770		0.2277		0.8729	
Sedatives	*		0.6692		0.8812	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.6989		0.6241		0.7763	

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52C Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Tobacco Product and Alcohol Use in the Past Year and Past Month: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
PAST YEAR						
Tobacco Products ¹	198	203	116	109	78	76
Cigarettes	187	192	112	101	77	68
Smokeless Tobacco	61	66	34	44	16	16
Cigars	101	111	63	76	36	49
Alcohol	266	265	131	129	87	86
PAST MONTH						
Tobacco Products ¹	190	192	111	101	75	69
Cigarettes	172	183	108	94	74	61
Smokeless Tobacco	50	50	24	29	15	13
Cigars	72	83	48	55	31	33
Pipe Tobacco	41	45	20	31	18	28
Alcohol	235	231	108	105	76	69
Binge Alcohol Use ²	163	170	75	62	60	53
Heavy Alcohol Use ²	103	100	49	46	34	31

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52D Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Tobacco Product and Alcohol Use in the Past Year and Past Month: Standard Errors of Percentages, 2011 and 2012

Substance	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
PAST YEAR						
Tobacco Products ¹	0.25	0.25	0.15	0.14	0.10	0.10
Cigarettes	0.28	0.29	0.17	0.15	0.12	0.11
Smokeless Tobacco	0.57	0.59	0.32	0.40	0.16	0.14
Cigars	0.44	0.45	0.27	0.31	0.16	0.20
Alcohol	0.16	0.15	0.08	0.08	0.05	0.05
PAST MONTH						
Tobacco Products ¹	0.28	0.28	0.17	0.15	0.11	0.10
Cigarettes	0.30	0.32	0.19	0.17	0.13	0.11
Smokeless Tobacco	0.64	0.59	0.31	0.34	0.19	0.15
Cigars	0.60	0.64	0.40	0.43	0.26	0.26
Pipe Tobacco	2.09	1.88	1.04	1.31	0.95	1.16
Alcohol	0.18	0.17	0.08	0.08	0.06	0.05
Binge Alcohol Use ²	0.28	0.29	0.13	0.11	0.11	0.09
Heavy Alcohol Use ²	0.65	0.59	0.32	0.28	0.22	0.18

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52P Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Tobacco Product and Alcohol Use in the Past Year and Past Month: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
PAST YEAR						
Tobacco Products ¹	0.3183		0.8081		0.4960	
Cigarettes	0.2567		0.9497		0.8483	
Smokeless Tobacco	0.8378		0.4668		0.9820	
Cigars	0.5246		0.2841		0.2705	
Alcohol	0.3974		0.6155		0.2038	
PAST MONTH						
Tobacco Products ¹	0.4605		0.8142		0.3923	
Cigarettes	0.3327		0.9783		0.7372	
Smokeless Tobacco	0.7682		0.8391		0.6644	
Cigars	0.8434		0.4856		0.7610	
Pipe Tobacco	0.7123		0.7166		0.9162	
Alcohol	0.4365		0.9212		0.2669	
Binge Alcohol Use ²	0.2199		0.5298		0.0709	
Heavy Alcohol Use ²	0.7185		0.7524		0.7521	

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.