

**Table 1.63C Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2011 and 2012**

Substance	Total <sup>1</sup> (2011)	Total <sup>1</sup> (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
<b>PAST YEAR USE</b>						
Illicit Drugs <sup>2</sup>	589	639	209	206	592	634
Marijuana and Hashish	503	549	172	179	510	543
Cocaine	167	217	65	74	151	201
Crack	68	111	35	42	59	100
Heroin	71	78	47	45	46	57
Hallucinogens	138	172	66	53	122	157
LSD	57	83	20	31	53	73
PCP	19	32	9	14	16	28
Ecstasy	103	132	49	47	90	121
Inhalants	99	115	47	56	86	101
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	357	398	138	150	334	370
Pain Relievers	297	349	123	121	277	327
OxyContin <sup>®</sup>	102	103	44	58	89	83
Tranquilizers	207	249	88	113	189	217
Stimulants <sup>4</sup>	149	160	68	70	125	138
Methamphetamine <sup>4</sup>	118	113	54	46	98	97
Sedatives	63	65	39	34	46	56
Illicit Drugs Other Than Marijuana <sup>2</sup>	397	450	153	160	376	421
<b>PAST MONTH USE</b>						
Daily Cigarette Use <sup>5</sup>	674	688	204	195	653	703
Heavy Alcohol Use <sup>6</sup>	423	448	115	122	417	442

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

<sup>1</sup> Estimates in the Total column represent all persons aged 18 or older, including those with unknown past year MDE information.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>5</sup> Daily Cigarette Use is defined as smoking on each of the past 30 days.

<sup>6</sup> Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.63D Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2011 and 2012**

Substance	Total <sup>1</sup> (2011)	Total <sup>1</sup> (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
<b>PAST YEAR USE</b>						
Illicit Drugs <sup>2</sup>	0.25	0.27	1.20	1.20	0.25	0.27
Marijuana and Hashish	0.22	0.23	1.03	1.05	0.22	0.23
Cocaine	0.07	0.09	0.41	0.46	0.07	0.09
Crack	0.03	0.05	0.23	0.26	0.03	0.05
Heroin	0.03	0.03	0.31	0.28	0.02	0.03
Hallucinogens	0.06	0.07	0.43	0.33	0.06	0.07
LSD	0.02	0.04	0.14	0.19	0.02	0.03
PCP	0.01	0.01	0.06	0.09	0.01	0.01
Ecstasy	0.04	0.06	0.32	0.29	0.04	0.06
Inhalants	0.04	0.05	0.31	0.34	0.04	0.05
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.15	0.17	0.86	0.90	0.15	0.16
Pain Relievers	0.13	0.15	0.76	0.74	0.13	0.15
OxyContin <sup>®</sup>	0.04	0.04	0.29	0.36	0.04	0.04
Tranquilizers	0.09	0.11	0.57	0.69	0.09	0.10
Stimulants <sup>4</sup>	0.06	0.07	0.45	0.44	0.06	0.06
Methamphetamine <sup>4</sup>	0.05	0.05	0.35	0.28	0.05	0.04
Sedatives	0.03	0.03	0.26	0.21	0.02	0.03
Illicit Drugs Other Than Marijuana <sup>2</sup>	0.17	0.19	0.94	0.97	0.17	0.19
<b>PAST MONTH USE</b>						
Daily Cigarette Use <sup>5</sup>	0.29	0.29	1.19	1.14	0.29	0.30
Heavy Alcohol Use <sup>6</sup>	0.18	0.19	0.72	0.74	0.18	0.20

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

<sup>1</sup> Estimates in the Total column represent all persons aged 18 or older, including those with unknown past year MDE information.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>5</sup> Daily Cigarette Use is defined as smoking on each of the past 30 days.

<sup>6</sup> Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.63P Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE):  
P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Total <sup>1</sup> (2011)	Total <sup>1</sup> (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
<b>PAST YEAR USE</b>						
Illicit Drugs <sup>2</sup>	0.0002		0.9945		0.0002	
Marijuana and Hashish	0.0213		1.0000		0.0227	
Cocaine	0.0028		0.5048		0.0037	
Crack	0.0268		0.4862		0.0391	
Heroin	0.5344		0.5265		0.1688	
Hallucinogens	0.1886		0.1638		0.0354	
LSD	0.0739		0.6775		0.0707	
PCP	0.2411		0.6772		0.2562	
Ecstasy	0.0776		0.6125		0.0285	
Inhalants	0.9655		0.7805		0.7891	
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.0002		0.0840		0.0013	
Pain Relievers	0.0020		0.5661		0.0028	
OxyContin <sup>®</sup>	0.3556		0.4422		0.1236	
Tranquilizers	0.0039		0.0751		0.0267	
Stimulants <sup>4</sup>	0.0113		0.3146		0.0362	
Methamphetamine <sup>4</sup>	0.4219		0.7020		0.3747	
Sedatives	0.4685		0.6851		0.2434	
Illicit Drugs Other Than Marijuana <sup>2</sup>	0.0000		0.2892		0.0001	
<b>PAST MONTH USE</b>						
Daily Cigarette Use <sup>5</sup>	0.8593		0.2426		0.5573	
Heavy Alcohol Use <sup>6</sup>	0.0811		0.0859		0.1961	

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

<sup>1</sup> Estimates in the Total column represent all persons aged 18 or older, including those with unknown past year MDE information.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>5</sup> Daily Cigarette Use is defined as smoking on each of the past 30 days.

<sup>6</sup> Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.