

Table 1.6C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

Age Category	Total (2011)	Total (2012)	Male (2011)	Male (2012)	Female (2011)	Female (2012)
TOTAL	325	341	190	211	261	258
18	17	18	11	10	13	15
19	24	22	15	13	18	18
20	24	24	13	13	20	20
21	24	24	17	17	17	17
22	17	20	8	11	15	17
23	19	22	12	15	15	18
24	20	20	10	14	17	13
25	20	20	13	14	15	14
26-29	90	94	55	70	70	65
30-34	95	101	61	66	69	73
35-39	95	121	52	75	81	95
40-44	103	100	50	66	87	76
45-49	106	102	60	75	88	70
50-54	153	122	101	85	119	89
55-59	108	108	53	69	91	85
60-64	111	120	81	81	76	88
65 or Older	67	135	29	68	57	109

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.6D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

Age Category	Total (2011)	Total (2012)	Male (2011)	Male (2012)	Female (2011)	Female (2012)
TOTAL	0.14	0.14	0.17	0.19	0.22	0.21
18	0.36	0.39	0.44	0.42	0.59	0.65
19	0.53	0.48	0.68	0.58	0.81	0.82
20	0.52	0.52	0.59	0.58	0.85	0.89
21	0.52	0.51	0.74	0.73	0.73	0.76
22	0.38	0.44	0.37	0.52	0.70	0.70
23	0.47	0.50	0.64	0.68	0.70	0.79
24	0.48	0.48	0.45	0.68	0.84	0.63
25	0.48	0.49	0.65	0.69	0.73	0.68
26-29	0.51	0.53	0.64	0.77	0.80	0.76
30-34	0.48	0.50	0.63	0.69	0.67	0.68
35-39	0.48	0.60	0.53	0.75	0.79	0.92
40-44	0.49	0.48	0.52	0.66	0.78	0.71
45-49	0.47	0.46	0.55	0.70	0.78	0.62
50-54	0.65	0.51	0.89	0.74	0.97	0.72
55-59	0.56	0.52	0.57	0.68	0.92	0.81
60-64	0.60	0.69	0.91	1.01	0.80	0.97
65 or Older	0.17	0.32	0.17	0.37	0.25	0.46

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment. Serious mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.6P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Age Category	Total (2011)	Total (2012)	Male (2011)	Male (2012)	Female (2011)	Female (2012)
TOTAL	0.3361		0.2021		0.7894	
18	0.7497		0.5455		0.3954	
19	0.5279		0.8822		0.5155	
20	0.6671		0.5381		0.9918	
21	0.4289		0.6682		0.4899	
22	0.0464		0.0249		0.4371	
23	0.3085		0.9766		0.2010	
24	0.6333		0.0680		0.3700	
25	0.4854		0.4135		0.8193	
26-29	0.5419		0.9587		0.4609	
30-34	0.3198		0.9264		0.2000	
35-39	0.5895		0.2634		0.9147	
40-44	0.6583		0.2347		0.1795	
45-49	0.7773		0.3447		0.5922	
50-54	0.0538		0.1446		0.2122	
55-59	0.9631		0.6849		0.7347	
60-64	0.2051		0.6011		0.2183	
65 or Older	0.0024		0.0688		0.0088	

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.7C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Demographic Characteristic	Total (2011)	Total (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26-49 (2011)	Aged 26-49 (2012)	Aged 50+ (2011)	Aged 50+ (2012)
TOTAL	325	341	59	59	213	228	224	254
GENDER								
Male	190	211	35	37	122	153	140	149
Female	261	258	46	45	172	166	174	191
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	309	300	54	53	199	192	218	229
White	284	284	46	52	196	179	197	213
Black or African American	84	104	19	16	61	66	55	79
American Indian or Alaska Native	45	35	3	3	15	15	*	*
Native Hawaiian or Other Pacific Islander	*	7	*	1	*	*	*	*
Asian	60	59	16	10	32	51	*	28
Two or More Races	36	26	14	13	15	16	29	16
Hispanic or Latino	99	155	24	26	76	119	51	89
CURRENT EMPLOYMENT								
Full-Time	185	184	32	31	142	148	109	107
Part-Time	131	110	32	33	96	82	78	63
Unemployed	77	112	23	27	60	90	42	60
Other ¹	223	258	34	31	123	138	183	213

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.7D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2011 and 2012

Demographic Characteristic	Total (2011)	Total (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26-49 (2011)	Aged 26-49 (2012)	Aged 50+ (2011)	Aged 50+ (2012)
TOTAL	0.14	0.14	0.17	0.17	0.22	0.23	0.22	0.25
GENDER								
Male	0.17	0.19	0.20	0.21	0.25	0.32	0.30	0.31
Female	0.22	0.21	0.27	0.26	0.34	0.33	0.32	0.35
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.16	0.15	0.20	0.19	0.25	0.24	0.24	0.25
White	0.18	0.17	0.22	0.24	0.31	0.29	0.26	0.27
Black or African American	0.31	0.37	0.39	0.32	0.50	0.52	0.55	0.75
American Indian or Alaska Native	3.34	2.52	1.43	1.07	3.04	2.70	*	*
Native Hawaiian or Other Pacific Islander	*	0.76	*	0.73	*	*	*	*
Asian	0.52	0.50	0.85	0.56	0.54	0.81	*	0.76
Two or More Races	1.08	0.79	1.79	1.62	1.21	1.24	2.21	1.19
Hispanic or Latino	0.29	0.44	0.33	0.37	0.42	0.65	0.57	0.95
CURRENT EMPLOYMENT								
Full-Time	0.16	0.15	0.25	0.22	0.21	0.22	0.29	0.28
Part-Time	0.40	0.33	0.33	0.33	0.81	0.69	0.68	0.53
Unemployed	0.55	0.79	0.50	0.63	0.98	1.40	1.27	1.66
Other ¹	0.32	0.35	0.40	0.37	0.81	0.88	0.38	0.42

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.7P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Demographic Characteristic	Total (2011)	Total (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26-49 (2011)	Aged 26-49 (2012)	Aged 50+ (2011)	Aged 50+ (2012)
TOTAL	0.3361		0.1673		0.7013		0.4759	
GENDER								
Male	0.2021		0.1961		0.1423		0.9432	
Female	0.7894		0.4418		0.4852		0.3749	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.6576		0.2644		0.2977		0.9139	
White	0.4137		0.0547		0.1410		0.8863	
Black or African American	0.1974		0.5574		0.7456		0.0912	
American Indian or Alaska Native	0.7696		0.7505		0.8757		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.8945		0.2019		0.2855		*	
Two or More Races	0.7590		0.5694		0.6222		0.2874	
Hispanic or Latino	0.0004		0.3564		0.0035		0.0331	
CURRENT EMPLOYMENT								
Full-Time	0.7994		0.4660		0.7372		0.8672	
Part-Time	0.1853		0.5327		0.2512		0.6198	
Unemployed	0.0135		0.0307		0.0428		0.4778	
Other ¹	0.3373		0.2244		0.8937		0.3032	

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.8C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Geographic/Socioeconomic Characteristic	Total (2011)	Total (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26-49 (2011)	Aged 26-49 (2012)	Aged 50+ (2011)	Aged 50+ (2012)
TOTAL	325	341	59	59	213	228	224	254
GEOGRAPHIC REGION								
Northeast	126	136	19	23	83	94	81	93
Midwest	127	119	23	23	88	83	85	82
South	188	227	36	37	121	132	137	177
West	184	174	37	31	127	138	122	117
COUNTY TYPE								
Large Metro	239	266	44	48	161	185	162	188
Small Metro	191	208	42	38	127	133	129	147
250K – 1 Mil. Pop.	157	142	34	29	112	96	100	112
< 250K Pop.	103	151	24	24	60	94	80	94
Nonmetro	125	125	17	20	88	75	79	91
Urbanized	81	68	11	11	59	44	53	49
Less Urbanized	85	98	10	12	64	54	56	72
Completely Rural	29	46	7	12	21	35	18	28
POVERTY LEVEL¹								
Less Than 100%	182	185	39	35	110	131	130	124
100-199%	148	163	28	29	106	100	97	126
200% or More	231	243	39	40	157	165	159	171
EDUCATION								
< High School	132	149	24	23	91	87	84	118
High School Graduate	156	190	30	36	100	127	120	144
Some College	186	181	44	40	126	131	114	123
College Graduate	171	168	22	19	108	123	126	110
HEALTH INSURANCE²								
Private	217	209	50	48	154	153	134	150
Medicaid/CHIP ³	159	160	20	22	106	107	117	113
Other ⁴	160	231	16	14	69	92	147	203
No Coverage	143	162	29	32	107	131	89	87

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment. Serious mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁴ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.8D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: Standard Errors of Percentages, 2011 and 2012

Geographic/Socioeconomic Characteristic	Total (2011)	Total (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26-49 (2011)	Aged 26-49 (2012)	Aged 50+ (2011)	Aged 50+ (2012)
TOTAL	0.14	0.14	0.17	0.17	0.22	0.23	0.22	0.25
GEOGRAPHIC REGION								
Northeast	0.30	0.32	0.32	0.37	0.47	0.54	0.43	0.48
Midwest	0.25	0.23	0.32	0.31	0.43	0.40	0.38	0.36
South	0.22	0.26	0.28	0.29	0.33	0.36	0.37	0.47
West	0.34	0.32	0.45	0.37	0.54	0.59	0.55	0.51
COUNTY TYPE								
Large Metro	0.19	0.20	0.23	0.24	0.29	0.32	0.31	0.36
Small Metro	0.25	0.28	0.34	0.30	0.38	0.45	0.41	0.45
250K – 1 Mil. Pop.	0.30	0.29	0.46	0.37	0.49	0.47	0.46	0.52
< 250K Pop.	0.41	0.59	0.47	0.51	0.60	0.98	0.79	0.85
Nonmetro	0.33	0.31	0.34	0.39	0.61	0.52	0.45	0.46
Urbanized	0.52	0.39	0.51	0.47	0.95	0.68	0.77	0.60
Less Urbanized	0.46	0.54	0.44	0.52	0.88	0.80	0.63	0.80
Completely Rural	0.74	0.98	1.56	2.40	1.59	2.32	0.85	1.06
POVERTY LEVEL¹								
Less Than 100%	0.53	0.47	0.39	0.34	0.72	0.77	1.41	1.04
100-199%	0.31	0.35	0.32	0.34	0.51	0.51	0.52	0.68
200% or More	0.15	0.16	0.24	0.24	0.24	0.26	0.22	0.23
EDUCATION								
< High School	0.39	0.42	0.41	0.42	0.67	0.64	0.59	0.74
High School Graduate	0.22	0.26	0.25	0.30	0.38	0.48	0.37	0.43
Some College	0.29	0.28	0.34	0.30	0.47	0.50	0.48	0.49
College Graduate	0.24	0.24	0.43	0.36	0.31	0.36	0.41	0.37
HEALTH INSURANCE²								
Private	0.14	0.14	0.25	0.22	0.23	0.24	0.19	0.21
Medicaid/CHIP ³	0.65	0.66	0.37	0.40	0.98	1.05	1.47	1.34
Other ⁴	0.29	0.38	0.67	0.52	1.04	1.24	0.31	0.40
No Coverage	0.37	0.41	0.32	0.40	0.48	0.59	1.13	1.01

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁴ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.8P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Geographic/Socioeconomic Characteristic	Total (2011)	Total (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26-49 (2011)	Aged 26-49 (2012)	Aged 50+ (2011)	Aged 50+ (2012)
TOTAL	0.3361		0.1673		0.7013		0.4759	
GEOGRAPHIC REGION								
Northeast	0.5127		0.2178		0.4399		0.9674	
Midwest	0.4336		0.3991		0.4920		0.5062	
South	0.1444		0.5986		0.6531		0.0347	
West	0.8317		0.6352		0.3335		0.3980	
COUNTY TYPE								
Large Metro	0.0931		0.3281		0.0582		0.7667	
Small Metro	0.9911		0.8298		0.6215		0.6417	
250K – 1 Mil. Pop.	0.8731		0.5466		0.2139		0.4418	
< 250K Pop.	0.8793		0.6110		0.4834		0.8281	
Nonmetro	0.4794		0.0538		0.0267		0.4531	
Urbanized	0.1415		0.6132		0.0661		0.8247	
Less Urbanized	1.0000		0.1819		0.1266		0.2766	
Completely Rural	0.5094		0.1100		0.8454		0.7375	
POVERTY LEVEL¹								
Less Than 100%	0.6173		0.4597		0.8902		0.3378	
100-199%	0.2363		0.5056		0.8012		0.0921	
200% or More	0.5596		0.3498		0.5911		0.8972	
EDUCATION								
< High School	0.8727		0.8367		0.3438		0.2463	
High School Graduate	0.0044		0.0808		0.1216		0.0592	
Some College	0.4805		0.3900		0.7684		0.3741	
College Graduate	0.7021		0.8315		0.7699		0.4283	
HEALTH INSURANCE²								
Private	0.6938		0.6073		0.5826		0.3417	
Medicaid/CHIP ³	0.7660		0.1769		0.5405		0.9877	
Other ⁴	0.0646		0.3282		0.4451		0.0803	
No Coverage	0.1185		0.0885		0.1006		0.8283	

*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates in 2011 used in the comparisons may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁴ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.