

**Table 2.10C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE):
Standard Errors of Numbers in Thousands, 2011 and 2012**

Dependence or Abuse	Total¹ (2011)	Total¹ (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
DEPENDENCE						
Illicit Drugs ²	37	32	17	16	32	29
Marijuana	32	29	13	14	28	27
Illicit Drugs Other Than Marijuana ²	20	15	13	10	15	12
Alcohol	26	25	12	13	23	21
Both Illicit Drugs and Alcohol ²	14	12	9	7	11	10
Illicit Drugs or Alcohol ²	43	39	19	19	39	35
DEPENDENCE OR ABUSE						
Illicit Drugs ²	47	45	21	21	43	41
Marijuana	41	41	16	18	38	38
Illicit Drugs Other Than Marijuana ²	29	24	16	14	24	19
Alcohol	45	40	21	19	40	36
Both Illicit Drugs and Alcohol ²	30	25	14	13	26	21
Illicit Drugs or Alcohol ²	58	55	27	25	54	51

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

¹ Estimates in the Total column represent all persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.10D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE):
Standard Errors of Percentages, 2011 and 2012**

Dependence or Abuse	Total¹ (2011)	Total¹ (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
DEPENDENCE						
Illicit Drugs ²	0.15	0.13	0.81	0.69	0.14	0.13
Marijuana	0.13	0.12	0.63	0.60	0.12	0.12
Illicit Drugs Other Than Marijuana ²	0.08	0.06	0.65	0.44	0.07	0.05
Alcohol	0.10	0.10	0.57	0.58	0.10	0.09
Both Illicit Drugs and Alcohol ²	0.06	0.05	0.43	0.32	0.05	0.04
Illicit Drugs or Alcohol ²	0.17	0.15	0.91	0.82	0.17	0.15
DEPENDENCE OR ABUSE						
Illicit Drugs ²	0.19	0.18	0.97	0.87	0.19	0.18
Marijuana	0.16	0.16	0.76	0.79	0.17	0.16
Illicit Drugs Other Than Marijuana ²	0.12	0.09	0.78	0.60	0.11	0.09
Alcohol	0.18	0.16	1.00	0.80	0.17	0.16
Both Illicit Drugs and Alcohol ²	0.12	0.10	0.69	0.56	0.11	0.10
Illicit Drugs or Alcohol ²	0.23	0.22	1.21	1.01	0.23	0.22

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

¹ Estimates in the Total column represent all persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.10P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): P Values from Tests of Differences of Percentages, 2012 Versus 2011

Dependence or Abuse	Total¹ (2011)	Total¹ (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
DEPENDENCE						
Illicit Drugs ²	0.0067		0.3054		0.0128	
Marijuana	0.0832		0.9670		0.0722	
Illicit Drugs Other Than Marijuana ²	0.0023		0.0236		0.0409	
Alcohol	0.1669		0.9466		0.1029	
Both Illicit Drugs and Alcohol ²	0.1321		0.4878		0.1176	
Illicit Drugs or Alcohol ²	0.0084		0.6022		0.0097	
DEPENDENCE OR ABUSE						
Illicit Drugs ²	0.0158		0.3778		0.0151	
Marijuana	0.2122		0.4837		0.0900	
Illicit Drugs Other Than Marijuana ²	0.0005		0.0238		0.0039	
Alcohol	0.1403		0.1491		0.2612	
Both Illicit Drugs and Alcohol ²	0.1784		0.4100		0.2586	
Illicit Drugs or Alcohol ²	0.0134		0.1519		0.0194	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

¹ Estimates in the Total column represent all persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.