

**Table 2.12C Type of Treatment Received in the Past Year for Depression among Persons Aged 12 to 17 with a Past Year Major Depressive Episode (MDE), by Demographic and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

Demographic/Socioeconomic Characteristic	Saw or Talked to Health Professional OR Used Prescription Medication (2011)	Saw or Talked to Health Professional OR Used Prescription Medication (2012)	Saw or Talked to Health Professional Only (2011)	Saw or Talked to Health Professional Only (2012)	Used Prescription Medication and Did Not See or Talk to Health Professional (2011)	Used Prescription Medication and Did Not See or Talk to Health Professional (2012)	Saw or Talked to Health Professional AND Used Prescription Medication (2011)	Saw or Talked to Health Professional AND Used Prescription Medication (2012)
<b>TOTAL</b>	35	37	26	27	10	9	21	23
<b>GENDER</b>								
Male	20	16	14	12	8	5	11	9
Female	31	33	22	25	6	7	19	20
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	32	32	22	23	10	8	20	21
White	26	28	17	18	9	7	17	19
Black or African American	13	14	10	12	4	3	8	5
American Indian or Alaska Native	*	*	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*
Hispanic or Latino	15	18	14	15	2	4	8	9
<b>POVERTY LEVEL<sup>1</sup></b>								
Less Than 100%	17	21	12	15	4	5	11	13
100-199%	18	17	13	12	6	3	10	11
200% or More	25	26	19	20	7	7	15	16
<b>HEALTH INSURANCE<sup>2</sup></b>								
Private	26	27	18	20	9	6	18	17
Medicaid/CHIP <sup>3</sup>	22	23	17	18	5	6	13	13
Other <sup>4</sup>	*	*	5	5	*	4	*	6
No Coverage	*	8	*	8	*	1	3	2

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

NOTE: Respondents with unknown past year treatment data were excluded.

NOTE: Health Professionals include general practitioner or family doctor; other medical doctor (e.g., cardiologist, gynecologist, urologist); psychologist; psychiatrist or psychotherapist; social worker; counselor; other mental health professional (e.g., mental health nurse or other therapist where type is not specified); and nurse, occupational therapist, or other health professional.

<sup>1</sup> Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>4</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.12D Type of Treatment Received in the Past Year for Depression among Persons Aged 12 to 17 with a Past Year Major Depressive Episode (MDE), by Demographic and Socioeconomic Characteristics: Standard Errors of Percentages, 2011 and 2012**

Demographic/Socioeconomic Characteristic	Saw or Talked to Health Professional OR Used Prescription Medication (2011)	Saw or Talked to Health Professional OR Used Prescription Medication (2012)	Saw or Talked to Health Professional Only (2011)	Saw or Talked to Health Professional Only (2012)	Used Prescription Medication and Did Not See or Talk to Health Professional (2011)	Used Prescription Medication and Did Not See or Talk to Health Professional (2012)	Saw or Talked to Health Professional AND Used Prescription Medication (2011)	Saw or Talked to Health Professional AND Used Prescription Medication (2012)
<b>TOTAL</b>	1.45	1.35	1.20	1.10	0.50	0.40	0.97	0.97
<b>GENDER</b>								
Male	2.80	2.35	2.19	1.87	1.38	0.84	1.77	1.45
Female	1.71	1.61	1.40	1.34	0.43	0.44	1.18	1.18
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	1.62	1.55	1.25	1.23	0.62	0.48	1.13	1.16
White	1.79	1.76	1.33	1.33	0.74	0.57	1.35	1.42
Black or African American	4.25	4.06	3.82	3.76	1.65	1.10	3.03	1.84
American Indian or Alaska Native	*	*	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*
Hispanic or Latino	3.19	2.72	3.13	2.40	0.53	0.71	1.91	1.53
<b>POVERTY LEVEL<sup>1</sup></b>								
Less Than 100%	3.17	2.91	2.63	2.39	0.97	0.83	2.32	2.21
100-199%	2.91	2.95	2.41	2.40	1.17	0.69	1.95	2.26
200% or More	2.00	1.71	1.60	1.47	0.66	0.55	1.32	1.24
<b>HEALTH INSURANCE<sup>2</sup></b>								
Private	1.89	1.73	1.47	1.41	0.79	0.48	1.42	1.30
Medicaid/CHIP <sup>3</sup>	2.49	2.37	2.06	1.98	0.67	0.74	1.71	1.59
Other <sup>4</sup>	*	*	4.38	4.20	*	2.88	*	4.41
No Coverage	*	4.61	*	4.54	*	0.97	2.17	1.05

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

NOTE: Respondents with unknown past year treatment data were excluded.

NOTE: Health Professionals include general practitioner or family doctor; other medical doctor (e.g., cardiologist, gynecologist, urologist); psychologist; psychiatrist or psychotherapist; social worker; counselor; other mental health professional (e.g., mental health nurse or other therapist where type is not specified); and nurse, occupational therapist, or other health professional.

<sup>1</sup> Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>4</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.12P Type of Treatment Received in the Past Year for Depression among Persons Aged 12 to 17 with a Past Year Major Depressive Episode (MDE), by Demographic and Socioeconomic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Demographic/Socioeconomic Characteristic	Saw or Talked to Health Professional OR Used Prescription Medication (2011)	Saw or Talked to Health Professional OR Used Prescription Medication (2012)	Saw or Talked to Health Professional Only (2011)	Saw or Talked to Health Professional Only (2012)	Used Prescription Medication and Did Not See or Talk to Health Professional (2011)	Used Prescription Medication and Did Not See or Talk to Health Professional (2012)	Saw or Talked to Health Professional AND Used Prescription Medication (2011)	Saw or Talked to Health Professional AND Used Prescription Medication (2012)
<b>TOTAL</b>	0.9030		0.8706		0.7063		0.9082	
<b>GENDER</b>								
Male	0.0943		0.2439		0.4258		0.5239	
Female	0.4706		0.6862		0.7715		0.6968	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.8468		0.5543		0.5273		0.5274	
White	0.7182		0.9549		0.5425		0.4513	
Black or African American	0.2502		0.9482		0.6730		0.1243	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	*		*		*		*	
Two or More Races	*		*		*		*	
Hispanic or Latino	0.6536		0.6158		0.3191		0.6742	
<b>POVERTY LEVEL<sup>1</sup></b>								
Less Than 100%	0.6373		0.7156		0.7344		0.6854	
100-199%	0.4339		0.7083		0.3533		0.8122	
200% or More	0.5101		0.7757		0.9013		0.5191	
<b>HEALTH INSURANCE<sup>2</sup></b>								
Private	0.8610		0.9092		0.1592		0.7807	
Medicaid/CHIP <sup>3</sup>	0.4466		0.6065		0.4841		0.4041	
Other <sup>4</sup>	*		0.9825		*		*	
No Coverage	*		*		*		0.1516	

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

NOTE: Respondents with unknown past year treatment data were excluded.

NOTE: Health Professionals include general practitioner or family doctor; other medical doctor (e.g., cardiologist, gynecologist, urologist); psychologist; psychiatrist or psychotherapist; social worker; counselor; other mental health professional (e.g., mental health nurse or other therapist where type is not specified); and nurse, occupational therapist, or other health professional.

<sup>1</sup> Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>4</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.