

**Table 2.9C Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2011 and 2012**

Substance	Total <sup>1</sup> (2011)	Total <sup>1</sup> (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
<b>PAST YEAR USE</b>						
Illicit Drugs <sup>2</sup>	93	83	38	36	94	90
Marijuana and Hashish	83	75	32	31	81	79
Cocaine	20	19	9	10	16	17
Crack	7	8	5	3	5	7
Heroin	11	9	4	4	9	8
Hallucinogens	35	32	14	15	31	27
LSD	18	16	7	8	14	13
PCP	7	11	5	3	6	10
Ecstasy	29	23	12	12	25	19
Inhalants	40	34	17	16	35	31
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	53	53	24	23	50	50
Pain Relievers	48	47	22	21	45	45
OxyContin <sup>®</sup>	18	17	10	8	16	15
Tranquilizers	31	31	14	15	28	27
Stimulants <sup>4</sup>	23	23	11	11	19	20
Methamphetamine <sup>4</sup>	16	12	7	6	13	9
Sedatives	12	11	5	7	11	9
Illicit Drugs Other Than Marijuana <sup>2</sup>	65	64	30	28	64	63
<b>PAST MONTH USE</b>						
Daily Cigarette Use <sup>5</sup>	28	25	11	9	25	23
Heavy Alcohol Use <sup>6</sup>	25	25	8	10	23	22

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

<sup>1</sup> Estimates in the Total column represent all persons aged 12 to 17, including those with unknown past year MDE information.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>5</sup> Daily Cigarette Use is defined as smoking on each of the past 30 days.

<sup>6</sup> Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.9D Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2011 and 2012**

Substance	Total <sup>1</sup> (2011)	Total <sup>1</sup> (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
<b>PAST YEAR USE</b>						
Illicit Drugs <sup>2</sup>	0.37	0.33	1.51	1.28	0.38	0.35
Marijuana and Hashish	0.33	0.30	1.37	1.18	0.34	0.31
Cocaine	0.08	0.08	0.45	0.44	0.07	0.07
Crack	0.03	0.03	0.23	0.15	0.02	0.03
Heroin	0.05	0.04	0.19	0.19	0.04	0.04
Hallucinogens	0.14	0.13	0.68	0.68	0.13	0.12
LSD	0.07	0.07	0.33	0.35	0.06	0.06
PCP	0.03	0.04	0.24	0.12	0.02	0.05
Ecstasy	0.12	0.09	0.58	0.54	0.11	0.09
Inhalants	0.16	0.14	0.80	0.69	0.15	0.14
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.21	0.21	1.09	0.96	0.21	0.21
Pain Relievers	0.19	0.19	1.02	0.88	0.19	0.19
OxyContin <sup>®</sup>	0.07	0.07	0.48	0.36	0.07	0.07
Tranquilizers	0.12	0.13	0.70	0.64	0.12	0.12
Stimulants <sup>4</sup>	0.09	0.09	0.55	0.49	0.08	0.09
Methamphetamine <sup>4</sup>	0.06	0.05	0.37	0.25	0.06	0.04
Sedatives	0.05	0.05	0.26	0.31	0.05	0.04
Illicit Drugs Other Than Marijuana <sup>2</sup>	0.26	0.26	1.29	1.09	0.26	0.26
<b>PAST MONTH USE</b>						
Daily Cigarette Use <sup>5</sup>	0.11	0.10	0.53	0.41	0.11	0.10
Heavy Alcohol Use <sup>6</sup>	0.10	0.10	0.41	0.43	0.10	0.10

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

<sup>1</sup> Estimates in the Total column represent all persons aged 12 to 17, including those with unknown past year MDE information.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>5</sup> Daily Cigarette Use is defined as smoking on each of the past 30 days.

<sup>6</sup> Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.9P Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Total <sup>1</sup> (2011)	Total <sup>1</sup> (2012)	MDE (2011)	MDE (2012)	No MDE (2011)	No MDE (2012)
<b>PAST YEAR USE</b>						
Illicit Drugs <sup>2</sup>	0.0423		0.3254		0.0368	
Marijuana and Hashish	0.1093		0.4919		0.0683	
Cocaine	0.0658		0.6214		0.0677	
Crack	0.2752		0.4251		0.4104	
Heroin	0.1301		0.8424		0.1800	
Hallucinogens	0.0511		0.5917		0.0055	
LSD	0.7848		0.5110		0.7710	
PCP	0.4936		0.5430		0.3666	
Ecstasy	0.0005		0.4607		0.0001	
Inhalants	0.0011		0.0955		0.0034	
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.1606		0.5784		0.0993	
Pain Relievers	0.0251		0.2986		0.0203	
OxyContin <sup>®</sup>	0.1875		0.2381		0.2431	
Tranquilizers	0.6277		0.6510		0.3780	
Stimulants <sup>4</sup>	0.4204		0.7933		0.4640	
Methamphetamine <sup>4</sup>	0.2272		0.3495		0.1707	
Sedatives	0.8751		0.6129		0.3564	
Illicit Drugs Other Than Marijuana <sup>2</sup>	0.0050		0.1899		0.0063	
<b>PAST MONTH USE</b>						
Daily Cigarette Use <sup>5</sup>	0.0384		0.0542		0.0771	
Heavy Alcohol Use <sup>6</sup>	0.1412		0.7244		0.1187	

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

<sup>1</sup> Estimates in the Total column represent all persons aged 12 to 17, including those with unknown past year MDE information.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>5</sup> Daily Cigarette Use is defined as smoking on each of the past 30 days.

<sup>6</sup> Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.