

Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	121,078 ^b	124,808	38,287 ^b	41,479	22,454 ^a	23,863
12	343	327	197	204	100	100
13	582	518	361	355	163	184
14	851	802	615	557	287	285
15	1,247	1,134	971 ^b	811	496 ^a	399
16	1,584	1,464	1,236	1,154	692	620
17	1,766	1,800	1,355	1,387	784	789
18	2,237	2,322	1,740	1,805	1,063	1,061
19	2,242	2,221	1,678	1,625	1,021	1,034
20	2,613	2,513	1,822	1,829	1,129	1,079
21	2,473	2,598	1,527	1,665	939	946
22	2,596	2,710	1,567	1,674	912	1,004
23	2,410 ^a	2,665	1,340	1,409	829	793
24	2,512	2,505	1,255	1,339	739	782
25	2,450	2,444	1,139	1,205	720	683
26-29	10,047	10,587	3,997	4,529	2,530	2,506
30-34	11,635	11,853	3,802	4,248	2,150 ^a	2,605
35-39	10,744	10,781	2,874	3,048	1,611	1,718
40-44	11,273	10,998	2,373	2,775	1,297	1,471
45-49	12,366	12,626	2,409	2,862	1,438	1,640
50-54	14,278	14,212	2,505	2,829	1,538	1,681
55-59	10,607	11,686	1,794	2,216	1,127	1,355
60-64	7,573	7,972	1,075	1,010	495	604
65 or Older	6,647 ^b	8,073	655	943	394	525

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	47.0	48.0	14.9 ^b	16.0	8.7	9.2
12	9.2	8.1	5.3	5.0	2.7	2.5
13	14.0	12.9	8.7	8.8	3.9	4.6
14	20.1	19.1	14.5	13.3	6.8	6.8
15	29.4	27.1	22.9 ^b	19.3	11.7 ^a	9.5
16	36.3	34.8	28.3	27.5	15.9	14.7
17	41.8	42.1	32.1	32.4	18.6	18.4
18	47.2	49.3	36.7	38.4	22.4	22.5
19	52.1	52.2	39.0	38.2	23.7	24.3
20	58.3	58.3	40.6	42.4	25.2	25.0
21	56.6	58.8	35.0	37.7	21.5	21.4
22	60.3	60.9	36.4	37.6	21.2	22.6
23	60.5	61.0	33.6	32.2	20.8	18.1
24	60.2	61.3	30.1	32.8	17.7	19.2
25	62.0	61.2	28.8	30.2	18.2	17.1
26-29	59.0	61.7	23.5	26.4	14.9	14.6
30-34	60.0	60.0	19.6	21.5	11.1 ^a	13.2
35-39	54.5	55.5	14.6	15.7	8.2	8.8
40-44	55.4	54.5	11.7	13.8	6.4	7.3
45-49	57.3	59.0	11.2 ^a	13.4	6.7	7.7
50-54	61.9	60.7	10.9	12.1	6.7	7.2
55-59	56.0	56.8	9.5	10.8	6.0	6.6
60-64	41.9 ^a	47.6	5.9	6.0	2.7	3.6
65 or Older	16.5 ^a	19.3	1.6	2.3	1.0	1.3

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	107,842 ^a	111,239	29,739 ^a	31,513	18,071	18,855
12	48	57	38	44	15	9
13	213	204	168	164	88	84
14	445	437	372	352	192	183
15	890	788	757 ^b	615	378	332
16	1,279	1,205	1,033	968	594	508
17	1,502	1,540	1,186	1,219	700	684
18	2,018	2,040	1,581	1,609	965	940
19	2,042	1,963	1,490	1,465	937	939
20	2,392	2,301	1,638	1,609	1,028	968
21	2,264	2,337	1,349	1,468	861	831
22	2,388	2,477	1,357	1,429	795	856
23	2,198 ^a	2,438	1,141	1,209	701	687
24	2,261	2,282	1,050	1,129	626	684
25	2,233	2,214	948	969	613	579
26-29	9,063	9,635	3,280	3,447	2,096	2,041
30-34	10,442	10,732	2,760 ^a	3,259	1,625 ^b	2,130
35-39	9,503	9,651	1,994	2,099	1,234	1,291
40-44	10,065	9,642	1,606	1,836	914	1,027
45-49	11,364	11,507	1,641	2,008	1,038	1,119
50-54	13,204	13,271	1,815	1,866	1,129	1,198
55-59	9,797	10,913	1,317	1,517	855	986
60-64	7,000	7,427	802	739	392	404
65 or Older	5,230 ^a	6,177	416	494	295	375

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	41.9	42.8	11.5 ^a	12.1	7.0	7.3
12	1.3	1.4	1.0	1.1	0.4	0.2
13	5.1	5.1	4.0	4.1	2.1	2.1
14	10.5	10.4	8.8	8.4	4.5	4.4
15	21.0	18.8	17.8 ^b	14.7	8.9	7.9
16	29.3	28.7	23.7	23.0	13.6	12.1
17	35.6	36.0	28.1	28.5	16.6	16.0
18	42.6	43.3	33.4	34.2	20.4	20.0
19	47.5	46.1	34.7	34.4	21.8	22.1
20	53.3	53.4	36.5	37.3	22.9	22.5
21	51.9	52.9	30.9	33.2	19.7	18.8
22	55.5	55.7	31.5	32.1	18.5	19.3
23	55.1	55.8	28.6	27.7	17.6	15.7
24	54.2	55.9	25.2	27.6	15.0	16.7
25	56.5	55.4	24.0	24.3	15.5	14.5
26-29	53.3	56.2	19.3	20.1	12.3	11.9
30-34	53.8	54.3	14.2 ^a	16.5	8.4 ^a	10.8
35-39	48.2	49.6	10.1	10.8	6.3	6.6
40-44	49.5	47.8	7.9	9.1	4.5	5.1
45-49	52.7	53.7	7.6	9.4	4.8	5.2
50-54	57.3	56.7	7.9	8.0	4.9	5.1
55-59	51.8	53.0	7.0	7.4	4.5	4.8
60-64	38.7 ^a	44.4	4.4	4.4	2.2	2.4
65 or Older	13.0	14.8	1.0	1.2	0.7	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	36,921	37,688	3,857 ^b	4,671	1,369	1,650
12	4	3	1	2	*	1
13	12	8	9	8	1	*
14	22	20	17	13	1	1
15	76	39	46 ^a	21	21	8
16	89	64	61	47	17	5
17	135	132	95	90	28	15
18	259	279	147	168	55	54
19	360	346	200	191	62	41
20	501	457	261	235	91 ^a	45
21	527	535	227	201	68	46
22	570	639	189	260	48	78
23	645	656	169	203	45	53
24	634	700	183	164	58	38
25	756	655	188	179	37	42
26-29	2,933	3,330	359 ^a	608	148	174
30-34	3,429	3,730	482	551	134	250
35-39	3,202	3,220	300	386	154	167
40-44	3,983	3,774	251	330	79	120
45-49	5,239	5,356	314	412	136	211
50-54	6,652	6,281	196	231	95	109
55-59	3,781	4,258	77 ^a	256	53	117
60-64	2,073	2,093	77	116	34	73
65 or Older	1,040	1,115	8	*	5	*

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	14.3	14.5	1.5 ^b	1.8	0.5	0.6
12	0.1	0.1	0.0	0.1	*	0.0
13	0.3	0.2	0.2	0.2	0.0	*
14	0.5	0.5	0.4	0.3	0.0	0.0
15	1.8	0.9	1.1 ^a	0.5	0.5	0.2
16	2.0	1.5	1.4	1.1	0.4	0.1
17	3.2	3.1	2.3	2.1	0.7	0.4
18	5.5	5.9	3.1	3.6	1.2	1.2
19	8.4	8.1	4.7	4.5	1.4	1.0
20	11.2	10.6	5.8	5.5	2.0 ^a	1.0
21	12.1	12.1	5.2	4.6	1.5	1.1
22	13.2	14.4	4.4	5.8	1.1	1.8
23	16.2	15.0	4.2	4.6	1.1	1.2
24	15.2	17.1	4.4	4.0	1.4	0.9
25	19.1	16.4	4.7	4.5	0.9	1.1
26-29	17.2	19.4	2.1 ^a	3.5	0.9	1.0
30-34	17.7	18.9	2.5	2.8	0.7	1.3
35-39	16.2	16.6	1.5	2.0	0.8	0.9
40-44	19.6	18.7	1.2	1.6	0.4	0.6
45-49	24.3	25.0	1.5	1.9	0.6	1.0
50-54	28.8	26.8	0.9	1.0	0.4	0.5
55-59	20.0	20.7	0.4 ^a	1.2	0.3	0.6
60-64	11.5	12.5	0.4	0.7	0.2	0.4
65 or Older	2.6	2.7	0.0	*	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	8,214	9,015	625 ^a	921	228 ^a	443
12	1	*	1	*	*	*
13	*	1	*	0	*	*
14	5	2	3	1	*	*
15	27	8	4	3	2	*
16	10	12	9	7	0	1
17	24	12	12	7	1	*
18	20	39	7	17	3	7
19	47	48	16	20	3	2
20	86	67	20	27	6	5
21	62	59	11	12	2	5
22	99	91	16	17	2	5
23	117	113	11	11	2	3
24	139	116	12	19	1	7
25	143	121	14	16	2	3
26-29	635	717	36	79	7	15
30-34	867 ^a	1,192	44	129	12	78
35-39	917	918	71	73	19	53
40-44	1,052	1,126	107	88	26	27
45-49	1,214	1,467	123	172	84	97
50-54	1,551	1,455	51	117	37	57
55-59	702	836	23	71	14	60
60-64	397	484	29	32	*	17
65 or Older	98	132	5	*	5	*

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	3.2	3.5	0.2 ^a	0.4	0.1 ^a	0.2
12	0.0	*	0.0	*	*	*
13	*	0.0	*	0.0	*	*
14	0.1	0.0	0.1	0.0	*	*
15	0.6	0.2	0.1	0.1	0.0	*
16	0.2	0.3	0.2	0.2	0.0	0.0
17	0.6	0.3	0.3	0.2	0.0	*
18	0.4	0.8	0.1	0.4	0.1	0.2
19	1.1	1.1	0.4	0.5	0.1	0.0
20	1.9	1.6	0.4	0.6	0.1	0.1
21	1.4	1.3	0.3	0.3	0.0	0.1
22	2.3	2.0	0.4	0.4	0.0	0.1
23	2.9	2.6	0.3	0.3	0.0	0.1
24	3.3	2.8	0.3	0.5	0.0	0.2
25	3.6	3.0	0.3	0.4	0.1	0.1
26-29	3.7	4.2	0.2	0.5	0.0	0.1
30-34	4.5 ^a	6.0	0.2	0.7	0.1	0.4
35-39	4.6	4.7	0.4	0.4	0.1	0.3
40-44	5.2	5.6	0.5	0.4	0.1	0.1
45-49	5.6	6.8	0.6	0.8	0.4	0.5
50-54	6.7	6.2	0.2	0.5	0.2	0.2
55-59	3.7	4.1	0.1	0.3	0.1	0.3
60-64	2.2	2.9	0.2	0.2	*	0.1
65 or Older	0.2	0.3	0.0	*	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	36,362	37,908	4,069	4,306	972	1,134
12	14	14	5	8	1	6
13	46	35	18	25	6	13
14	75	89	55	61	22	17
15	170	127	129 ^a	87	47	24
16	297 ^a	220	227	171	84 ^a	45
17	327	339	214	200	55	54
18	525	551	347	344	82 ^a	137
19	596	586	353	375	99	105
20	750	721	378	371	74	78
21	809	776	303	277	93	71
22	809	909	303	273	67	75
23	877	858	239	224	43	69
24	812	857	195	188	50	34
25	852	845	204	184	49	31
26-29	3,437	3,833	353 ^a	545	64	110
30-34	4,447	4,653	288	351	82	85
35-39	3,411	3,895	216	249	24	56
40-44	3,409	3,173	106	82	7	21
45-49	3,841	3,885	82	112	19	49
50-54	4,898	4,332	38	35	*	35
55-59	3,240	3,822	9	16	2	4
60-64	1,911	2,229	*	69	*	*
65 or Older	807	1,159	5	57	*	13

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	14.1	14.6	1.6	1.7	0.4	0.4
12	0.4	0.4	0.1	0.2	0.0	0.2
13	1.1	0.9	0.4	0.6	0.1	0.3
14	1.8	2.1	1.3	1.5	0.5	0.4
15	4.0	3.0	3.0 ^a	2.1	1.1	0.6
16	6.8 ^a	5.2	5.2	4.1	1.9	1.1
17	7.7	7.9	5.1	4.7	1.3	1.3
18	11.1	11.7	7.3	7.3	1.7 ^a	2.9
19	13.9	13.8	8.2	8.8	2.3	2.5
20	16.7	16.7	8.4	8.6	1.7	1.8
21	18.5	17.6	6.9	6.3	2.1	1.6
22	18.8	20.4	7.0	6.1	1.6	1.7
23	22.0	19.6	6.0	5.1	1.1	1.6
24	19.5	21.0	4.7	4.6	1.2	0.8
25	21.6	21.2	5.2	4.6	1.2	0.8
26-29	20.2	22.3	2.1 ^a	3.2	0.4	0.6
30-34	22.9	23.6	1.5	1.8	0.4	0.4
35-39	17.3 ^a	20.0	1.1	1.3	0.1	0.3
40-44	16.8	15.7	0.5	0.4	0.0	0.1
45-49	17.8	18.1	0.4	0.5	0.1	0.2
50-54	21.2	18.5	0.2	0.1	*	0.1
55-59	17.1	18.6	0.0	0.1	0.0	0.0
60-64	10.6	13.3	*	0.4	*	*
65 or Older	2.0	2.8	0.0	0.1	*	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	20,523	21,095	1,861	1,693	624	493
12	197	172	89	73	35	31
13	312 ^b	222	152 ^a	104	47	43
14	351	316	181	142	48	38
15	348	324	145	108	28	18
16	346	296	141	125	28	27
17	323	283	112	95	36	31
18	335	358	103	107	11	28
19	316	312	90	68	38	16
20	389	325	86	64	27	16
21	357	350	65	88	26	35
22	443	399	69	56	18	12
23	445	386	42	30	17	8
24	399	374	33	15	5	1
25	422	413	13 ^b	45	3	7
26-29	1,829	1,891	105	121	29	39
30-34	2,176 ^b	2,735	89	66	59	18
35-39	2,232	2,180	50	94	14	14
40-44	2,227	2,275	67	90	42	5
45-49	2,235	2,097	45	117	45	43
50-54	2,049	1,951	116	39	15	26
55-59	1,702	1,878	30	37	30	37
60-64	566	762	24	9	10	*
65 or Older	525	795	14	*	14	*

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	8.0	8.1	0.7	0.7	0.2	0.2
12	5.3	4.2	2.4	1.8	0.9	0.8
13	7.5 ^b	5.5	3.7 ^a	2.6	1.1	1.1
14	8.3	7.5	4.3	3.4	1.1	0.9
15	8.2	7.7	3.4	2.6	0.7	0.4
16	7.9	7.0	3.2	3.0	0.6	0.6
17	7.7	6.6	2.7	2.2	0.8	0.7
18	7.1	7.6	2.2	2.3	0.2	0.6
19	7.3	7.3	2.1	1.6	0.9	0.4
20	8.7	7.5	1.9	1.5	0.6	0.4
21	8.2	7.9	1.5	2.0	0.6	0.8
22	10.3	9.0	1.6	1.3	0.4	0.3
23	11.2 ^a	8.8	1.0	0.7	0.4	0.2
24	9.6	9.2	0.8	0.4	0.1	0.0
25	10.7	10.3	0.3 ^b	1.1	0.1	0.2
26-29	10.7	11.0	0.6	0.7	0.2	0.2
30-34	11.2 ^a	13.8	0.5	0.3	0.3	0.1
35-39	11.3	11.2	0.3	0.5	0.1	0.1
40-44	10.9	11.3	0.3	0.4	0.2	0.0
45-49	10.4	9.8	0.2	0.5	0.2	0.2
50-54	8.9	8.3	0.5	0.2	0.1	0.1
55-59	9.0	9.1	0.2	0.2	0.2	0.2
60-64	3.1	4.6	0.1	0.1	0.1	*
65 or Older	1.3	1.9	0.0	*	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	51,243 ^b	54,389	14,657 ^b	16,666	6,119 ^a	6,831
12	162	164	105	119	52	66
13	213	219	129	149	52	70
14	301	305	214	198	90	100
15	480	469	356	302	134	114
16	631	580	471	394	177	159
17	735	745	475	479	183	185
18	922	1,037	538	634	211	224
19	941	990	574	556	230	217
20	1,225	1,213	667	701	253	252
21	1,210	1,211	563	563	193	204
22	1,252	1,405	582 ^a	743	230	297
23	1,269	1,328	501	546	196	235
24	1,290	1,266	513	525	219	205
25	1,264	1,259	427	460	178	198
26-29	5,030	5,650	1,569 ^a	1,951	693	730
30-34	5,364	5,814	1,687	1,807	745	773
35-39	4,259	4,791	1,147	1,304	484	559
40-44	4,489	4,623	1,176	1,296	459	516
45-49	5,092	5,329	940	1,074	402	445
50-54	6,067	5,912	919	1,289	506	564
55-59	4,581	4,774	599	832	250	380
60-64	2,485	2,654	270	300	103	200
65 or Older	1,981 ^a	2,652	235	442	80	137

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	19.9 ^a	20.9	5.7 ^b	6.4	2.4	2.6
12	4.3	4.0	2.8	3.0	1.4	1.6
13	5.1	5.5	3.1	3.7	1.3	1.7
14	7.1	7.2	5.1	4.7	2.1	2.4
15	11.3	11.2	8.4	7.2	3.2	2.7
16	14.4	13.8	10.8	9.4	4.1	3.8
17	17.4	17.4	11.3	11.2	4.3	4.3
18	19.5	22.0	11.4 ^a	13.5	4.4	4.8
19	21.9	23.3	13.4	13.1	5.4	5.1
20	27.3	28.1	14.9	16.3	5.6	5.8
21	27.7	27.4	12.9	12.7	4.4	4.6
22	29.1	31.6	13.5 ^a	16.7	5.3	6.7
23	31.8	30.4	12.6	12.5	4.9	5.4
24	30.9	31.0	12.3	12.8	5.3	5.0
25	32.0	31.5	10.8	11.5	4.5	5.0
26-29	29.6 ^a	32.9	9.2 ^a	11.4	4.1	4.3
30-34	27.7	29.4	8.7	9.1	3.8	3.9
35-39	21.6 ^a	24.6	5.8	6.7	2.5	2.9
40-44	22.1	22.9	5.8	6.4	2.3	2.6
45-49	23.6	24.9	4.4	5.0	1.9	2.1
50-54	26.3	25.3	4.0	5.5	2.2	2.4
55-59	24.2	23.2	3.2	4.0	1.3	1.8
60-64	13.7	15.9	1.5	1.8	0.6	1.2
65 or Older	4.9	6.3	0.6	1.1	0.2	0.3

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	34,247 ^b	37,045	11,143 ^b	12,489	4,471	4,862
12	134	147	92	107	48	59
13	177	185	111	125	42	59
14	258	246	169	162	75	88
15	398	383	297	247	116	93
16	561	497	413 ^a	329	150	134
17	621	611	401	362	132	124
18	771	871	438	499	149	171
19	785	772	444	410	177	148
20	987	979	492	528	164	190
21	974	957	416	361	147	137
22	1,026	1,134	442	550	174	208
23	995	1,031	384	410	133	167
24	1,025	1,020	403	394	150	138
25	1,036	996	334	354	126	152
26-29	4,018	4,498	1,180	1,459	538	559
30-34	3,893	4,337	1,355	1,288	546	551
35-39	2,959 ^a	3,586	859	994	338	397
40-44	2,947	3,273	842	997	330	382
45-49	3,064	3,255	655	810	310	346
50-54	3,210	3,269	579 ^a	919	303	348
55-59	2,254	2,694	426	613	165	163
60-64	1,052	1,259	211	233	92	135
65 or Older	1,099	1,047	199	336	65	111

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2011 and 2012

Age Category	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	13.3 ^b	14.2	4.3 ^b	4.8	1.7	1.9
12	3.6	3.6	2.5	2.7	1.3	1.5
13	4.2	4.6	2.7	3.1	1.0	1.5
14	6.1	5.9	4.0	3.9	1.8	2.1
15	9.4	9.1	7.0	5.9	2.7	2.2
16	12.9	11.8	9.5	7.8	3.4	3.2
17	14.7	14.3	9.5	8.5	3.1	2.9
18	16.3	18.5	9.2	10.6	3.1	3.6
19	18.3	18.1	10.3	9.6	4.1	3.5
20	22.0	22.7	11.0	12.3	3.7	4.4
21	22.3	21.7	9.5	8.2	3.4	3.1
22	23.8	25.5	10.3	12.4	4.0	4.7
23	25.0	23.6	9.6	9.4	3.3	3.8
24	24.6	25.0	9.7	9.6	3.6	3.4
25	26.2	24.9	8.4	8.9	3.2	3.8
26-29	23.6	26.2	6.9	8.5	3.2	3.3
30-34	20.1	22.0	7.0	6.5	2.8	2.8
35-39	15.0 ^a	18.4	4.4	5.1	1.7	2.0
40-44	14.5	16.2	4.1	4.9	1.6	1.9
45-49	14.2	15.2	3.0	3.8	1.4	1.6
50-54	13.9	14.0	2.5 ^a	3.9	1.3	1.5
55-59	11.9	13.1	2.2	3.0	0.9	0.8
60-64	5.8	7.5	1.2	1.4	0.5	0.8
65 or Older	2.7	2.5	0.5	0.8	0.2	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.