

**Table 1.78A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	51,243 <sup>b</sup>	54,389	14,657 <sup>b</sup>	16,666	6,119 <sup>a</sup>	6,831
<b>GEOGRAPHIC DIVISION</b>						
Northeast	8,532	9,116	2,466	2,738	1,048	1,055
New England	2,350	2,599	691	686	278	277
Middle Atlantic	6,182	6,517	1,775	2,052	770	778
Midwest	10,685	11,045	3,144	3,372	1,393	1,302
East North Central	7,529	8,059	2,272	2,484	1,039	1,039
West North Central	3,156	2,986	871	887	354	263
South	18,419	19,039	5,334 <sup>a</sup>	6,075	2,145 <sup>a</sup>	2,661
South Atlantic	9,515	9,991	2,652	2,911	975 <sup>a</sup>	1,249
East South Central	3,095	2,954	927	1,058	422	492
West South Central	5,810	6,094	1,754 <sup>a</sup>	2,106	749	919
West	13,606 <sup>b</sup>	15,189	3,713 <sup>b</sup>	4,481	1,534	1,812
Mountain	4,405	4,630	1,239	1,354	469	540
Pacific	9,202 <sup>b</sup>	10,559	2,473 <sup>a</sup>	3,127	1,065	1,273
<b>COUNTY TYPE</b>						
Large Metro	27,040 <sup>b</sup>	30,081	7,672 <sup>b</sup>	9,421	3,036 <sup>b</sup>	3,871
Small Metro	16,833	16,870	4,844	4,990	2,122	2,121
250K – 1 Mil. Pop.	11,692	11,547	3,207	3,377	1,398	1,493
< 250K Pop.	5,142	5,324	1,637	1,612	724	628
Nonmetro	7,369	7,438	2,141	2,256	961	839
Urbanized	3,398	3,645	1,101	1,062	485	394
Less Urbanized	3,196	3,084	873	1,010	405	378
Completely Rural	775	709	168	184	72	66

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.78B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	19.9 <sup>a</sup>	20.9	5.7 <sup>b</sup>	6.4	2.4	2.6
<b>GEOGRAPHIC DIVISION</b>						
Northeast	18.2	19.3	5.3	5.8	2.2	2.2
New England	19.1	20.9	5.6	5.5	2.3	2.2
Middle Atlantic	17.9	18.7	5.1	5.9	2.2	2.2
Midwest	19.2	19.8	5.6	6.0	2.5	2.3
East North Central	19.5	20.8	5.9	6.4	2.7	2.7
West North Central	18.6	17.5	5.1	5.2	2.1	1.5
South	19.4	19.8	5.6 <sup>a</sup>	6.3	2.3 <sup>a</sup>	2.8
South Atlantic	19.0	19.6	5.3	5.7	1.9	2.5
East South Central	20.2	19.2	6.1	6.9	2.8	3.2
West South Central	19.6	20.2	5.9	7.0	2.5	3.1
West	22.7 <sup>a</sup>	25.1	6.2 <sup>a</sup>	7.4	2.6	3.0
Mountain	24.3	25.1	6.8	7.3	2.6	2.9
Pacific	22.1 <sup>a</sup>	25.0	5.9 <sup>a</sup>	7.4	2.6	3.0
<b>COUNTY TYPE</b>						
Large Metro	19.8 <sup>b</sup>	21.6	5.6 <sup>b</sup>	6.8	2.2 <sup>b</sup>	2.8
Small Metro	20.9	21.5	6.0	6.4	2.6	2.7
250K – 1 Mil. Pop.	21.5	21.8	5.9	6.4	2.6	2.8
< 250K Pop.	19.7	20.8	6.3	6.3	2.8	2.5
Nonmetro	18.3	17.6	5.3	5.3	2.4	2.0
Urbanized	20.2	19.7	6.6	5.7	2.9	2.1
Less Urbanized	16.7	16.2	4.6	5.3	2.1	2.0
Completely Rural	18.5	14.9	4.0	3.9	1.7	1.4

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.79A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	2,522	2,482	1,749	1,642	688	693
<b>GEOGRAPHIC DIVISION</b>						
Northeast	375	324	264	215	99	84
New England	89	74	60	55	21	23
Middle Atlantic	286	250	204 <sup>a</sup>	160	78	61
Midwest	539	510	384	356	167	147
East North Central	368	374	268	256	119	107
West North Central	172	136	116	100	48	40
South	941	984	652	638	269	287
South Atlantic	401	455	278	296	112	136
East South Central	173	164	119	109	58	45
West South Central	367	366	255	233	99	105
West	667	664	449	433	153	175
Mountain	209	229	139	166	53	67
Pacific	458	435	310	267	100	108
<b>COUNTY TYPE</b>						
Large Metro	1,284	1,248	886	841	315	351
Small Metro	803	800	577	528	255	226
250K – 1 Mil. Pop.	546	537	396	364	171	157
< 250K Pop.	257	263	181	164	84	69
Nonmetro	435	433	286	272	118	116
Urbanized	184	217	123	123	53	54
Less Urbanized	214	175	137	114	54	48
Completely Rural	38	41	26	35	11	14

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.79B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	10.1	10.0	7.0	6.6	2.8	2.8
<b>GEOGRAPHIC DIVISION</b>						
Northeast	8.8	7.6	6.2	5.1	2.3	2.0
New England	8.0	6.7	5.4	4.9	1.9	2.1
Middle Atlantic	9.0	8.0	6.4	5.1	2.4	1.9
Midwest	9.9	9.4	7.0	6.6	3.1	2.7
East North Central	9.7	9.9	7.0	6.8	3.1	2.8
West North Central	10.5	8.3	7.1	6.1	3.0	2.4
South	10.2	10.6	7.0	6.9	2.9	3.1
South Atlantic	8.7	9.8	6.0	6.4	2.4	2.9
East South Central	11.7	11.1	8.0	7.4	3.9	3.1
West South Central	11.6	11.5	8.1	7.3	3.1	3.3
West	11.1	11.1	7.5	7.3	2.6	2.9
Mountain	11.3	12.3	7.5	8.9	2.9	3.6
Pacific	11.0	10.6	7.5	6.5	2.4	2.6
<b>COUNTY TYPE</b>						
Large Metro	9.6	9.2	6.7	6.2	2.4	2.6
Small Metro	10.2	10.8	7.4	7.1	3.3	3.1
250K – 1 Mil. Pop.	10.2	10.5	7.4	7.1	3.2	3.1
< 250K Pop.	10.4	11.4	7.3	7.1	3.4	3.0
Nonmetro	11.4	11.1	7.5	6.9	3.1	3.0
Urbanized	10.9	12.0	7.3	6.8	3.1	3.0
Less Urbanized	12.1	10.3	7.7	6.7	3.0	2.9
Completely Rural	10.6	10.0	7.4	8.4	3.2	3.3

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.80A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	9,373	9,709	4,366 <sup>a</sup>	4,729	1,711	1,832
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,452 <sup>a</sup>	1,637	752 <sup>a</sup>	876	285	362
New England	435	480	227	238	89	101
Middle Atlantic	1,017 <sup>a</sup>	1,157	525 <sup>a</sup>	638	196	261
Midwest	2,031	2,072	987	969	401	338
East North Central	1,411	1,448	695	687	292	265
West North Central	619	624	293	283	109 <sup>a</sup>	72
South	3,465	3,541	1,554	1,693	614	694
South Atlantic	1,731	1,782	761	876	290	351
East South Central	608	540	287	257	131	124
West South Central	1,126	1,220	506	561	192	219
West	2,425	2,459	1,072	1,190	411	438
Mountain	818	777	390	342	129	133
Pacific	1,607	1,682	682 <sup>a</sup>	848	283	305
<b>COUNTY TYPE</b>						
Large Metro	4,754 <sup>b</sup>	5,352	2,221 <sup>b</sup>	2,687	886 <sup>a</sup>	1,056
Small Metro	3,273	3,074	1,526	1,490	598	554
250K – 1 Mil. Pop.	2,112	1,940	975	932	352	357
< 250K Pop.	1,161	1,134	551	558	247	197
Nonmetro	1,346	1,283	619	552	227	223
Urbanized	698	618	324	242	111	91
Less Urbanized	546	554	258	259	104	115
Completely Rural	102	111	37	51	12	16

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.80B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	27.3	28.1	12.7 <sup>a</sup>	13.7	5.0	5.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	23.7 <sup>a</sup>	26.6	12.3 <sup>a</sup>	14.2	4.7	5.9
New England	27.1	29.6	14.2	14.7	5.5	6.2
Middle Atlantic	22.5	25.5	11.6 <sup>a</sup>	14.1	4.3	5.8
Midwest	27.7	28.1	13.5	13.2	5.5	4.6
East North Central	27.9	28.5	13.7	13.5	5.8	5.2
West North Central	27.2	27.3	12.9	12.4	4.8 <sup>a</sup>	3.2
South	27.5	27.8	12.3	13.3	4.9	5.4
South Atlantic	26.9	27.3	11.8	13.4	4.5	5.4
East South Central	30.3	26.7	14.3	12.7	6.5	6.1
West South Central	27.1	29.0	12.2	13.3	4.6	5.2
West	29.5	29.6	13.0	14.3	5.0	5.3
Mountain	33.1	31.1	15.8	13.7	5.2	5.4
Pacific	27.9	28.9	11.8 <sup>a</sup>	14.6	4.9	5.2
<b>COUNTY TYPE</b>						
Large Metro	25.8 <sup>b</sup>	28.3	12.1 <sup>b</sup>	14.2	4.8	5.6
Small Metro	29.4	28.4	13.7	13.8	5.4	5.1
250K – 1 Mil. Pop.	29.8	28.2	13.8	13.5	5.0	5.2
< 250K Pop.	28.6	28.9	13.6	14.2	6.1	5.0
Nonmetro	28.4	26.5	13.0	11.4	4.8	4.6
Urbanized	31.6	27.8	14.7 <sup>a</sup>	10.9	5.0	4.1
Less Urbanized	26.1	25.4	12.3	11.9	5.0	5.3
Completely Rural	23.0	24.9	8.3	11.4	2.6	3.6

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.81A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	39,348 <sup>b</sup>	42,198	8,542 <sup>b</sup>	10,296	3,720	4,306
<b>GEOGRAPHIC DIVISION</b>						
Northeast	6,704	7,155	1,450	1,647	664	609
New England	1,826	2,045	403	393	168	152
Middle Atlantic	4,878	5,110	1,047	1,254	496	457
Midwest	8,115	8,463	1,772	2,046	825	818
East North Central	5,750	6,236	1,309	1,542	629	667
West North Central	2,365	2,227	463	504	196	151
South	14,014	14,514	3,128 <sup>a</sup>	3,744	1,262 <sup>a</sup>	1,679
South Atlantic	7,384	7,755	1,613	1,739	573	762
East South Central	2,313	2,251	522	693	233	322
West South Central	4,317	4,509	993 <sup>a</sup>	1,313	457	595
West	10,514 <sup>b</sup>	12,067	2,192 <sup>a</sup>	2,858	969	1,199
Mountain	3,378	3,625	711	846	287	339
Pacific	7,136 <sup>a</sup>	8,442	1,481 <sup>a</sup>	2,012	682	860
<b>COUNTY TYPE</b>						
Large Metro	21,002 <sup>b</sup>	23,480	4,565 <sup>b</sup>	5,892	1,835 <sup>a</sup>	2,465
Small Metro	12,758	12,997	2,741	2,972	1,268	1,341
250K – 1 Mil. Pop.	9,034	9,070	1,836	2,082	875	979
< 250K Pop.	3,724	3,927	905	890	393	362
Nonmetro	5,588	5,722	1,236	1,433	616	500
Urbanized	2,516	2,809	654	697	321	249
Less Urbanized	2,436	2,356	478	637	247	215
Completely Rural	635	557	105	99	49	36

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.81B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	19.8 <sup>a</sup>	21.0	4.3 <sup>b</sup>	5.1	1.9	2.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	18.4	19.5	4.0	4.5	1.8	1.7
New England	19.0	21.1	4.2	4.1	1.7	1.6
Middle Atlantic	18.1	18.9	3.9	4.6	1.8	1.7
Midwest	18.9	19.6	4.1	4.7	1.9	1.9
East North Central	19.3	20.8	4.4	5.1	2.1	2.2
West North Central	18.1	16.9	3.5	3.8	1.5	1.1
South	19.1	19.5	4.3	5.0	1.7 <sup>a</sup>	2.3
South Atlantic	18.9	19.5	4.1	4.4	1.5	1.9
East South Central	19.6	18.9	4.4	5.8	2.0	2.7
West South Central	19.3	19.8	4.4	5.8	2.0	2.6
West	23.1 <sup>a</sup>	26.1	4.8 <sup>a</sup>	6.2	2.1	2.6
Mountain	24.4	25.8	5.1	6.0	2.1	2.4
Pacific	22.5 <sup>a</sup>	26.2	4.7 <sup>a</sup>	6.2	2.1	2.7
<b>COUNTY TYPE</b>						
Large Metro	20.0 <sup>a</sup>	22.0	4.3 <sup>b</sup>	5.5	1.7 <sup>a</sup>	2.3
Small Metro	20.7	21.6	4.4	4.9	2.1	2.2
250K – 1 Mil. Pop.	21.5	22.2	4.4	5.1	2.1	2.4
< 250K Pop.	19.0	20.3	4.6	4.6	2.0	1.9
Nonmetro	17.7	17.1	3.9	4.3	1.9	1.5
Urbanized	19.5	19.4	5.1	4.8	2.5	1.7
Less Urbanized	15.9	15.6	3.1	4.2	1.6	1.4
Completely Rural	18.7	14.3	3.1	2.5	1.4	0.9

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.82A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	48,721 <sup>b</sup>	51,908	12,908 <sup>b</sup>	15,025	5,431 <sup>a</sup>	6,138
<b>GEOGRAPHIC DIVISION</b>						
Northeast	8,157	8,792	2,202	2,523	949	971
New England	2,261	2,525	631	631	257	254
Middle Atlantic	5,896	6,267	1,571	1,892	692	718
Midwest	10,146	10,535	2,760	3,016	1,226	1,156
East North Central	7,161	7,685	2,004	2,229	921	932
West North Central	2,985	2,850	756	787	305	224
South	17,479	18,055	4,682 <sup>a</sup>	5,438	1,876 <sup>a</sup>	2,374
South Atlantic	9,114	9,536	2,374	2,615	863	1,113
East South Central	2,921	2,791	809	950	364	447
West South Central	5,443	5,728	1,499 <sup>a</sup>	1,873	649	814
West	12,939 <sup>b</sup>	14,525	3,264 <sup>b</sup>	4,048	1,380	1,637
Mountain	4,196	4,401	1,101	1,188	415	472
Pacific	8,743 <sup>b</sup>	10,124	2,163 <sup>b</sup>	2,860	965	1,165
<b>COUNTY TYPE</b>						
Large Metro	25,756 <sup>b</sup>	28,833	6,786 <sup>b</sup>	8,579	2,721 <sup>b</sup>	3,520
Small Metro	16,031	16,070	4,267	4,461	1,866	1,895
250K – 1 Mil. Pop.	11,146	11,010	2,811	3,013	1,227	1,336
< 250K Pop.	4,885	5,061	1,456	1,448	640	559
Nonmetro	6,934	7,005	1,855	1,984	843	723
Urbanized	3,214	3,427	978	939	432	340
Less Urbanized	2,982	2,909	736	896	351	330
Completely Rural	737	668	142	150	60	53

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.82B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	20.9 <sup>a</sup>	22.1	5.5 <sup>b</sup>	6.4	2.3	2.6
<b>GEOGRAPHIC DIVISION</b>						
Northeast	19.1	20.5	5.2	5.9	2.2	2.3
New England	20.2	22.3	5.6	5.6	2.3	2.2
Middle Atlantic	18.8	19.8	5.0	6.0	2.2	2.3
Midwest	20.2	20.9	5.5	6.0	2.4	2.3
East North Central	20.5	21.9	5.7	6.4	2.6	2.7
West North Central	19.4	18.4	4.9	5.1	2.0	1.4
South	20.3	20.7	5.4 <sup>a</sup>	6.2	2.2 <sup>a</sup>	2.7
South Atlantic	20.0	20.6	5.2	5.7	1.9	2.4
East South Central	21.1	20.0	5.8	6.8	2.6	3.2
West South Central	20.5	21.3	5.6 <sup>a</sup>	7.0	2.4	3.0
West	24.0 <sup>a</sup>	26.6	6.1 <sup>a</sup>	7.4	2.6	3.0
Mountain	25.7	26.6	6.7	7.2	2.5	2.9
Pacific	23.3 <sup>a</sup>	26.6	5.8 <sup>a</sup>	7.5	2.6	3.1
<b>COUNTY TYPE</b>						
Large Metro	20.9 <sup>b</sup>	22.9	5.5 <sup>b</sup>	6.8	2.2 <sup>b</sup>	2.8
Small Metro	22.0	22.6	5.9	6.3	2.6	2.7
250K – 1 Mil. Pop.	22.7	23.0	5.7	6.3	2.5	2.8
< 250K Pop.	20.7	21.7	6.2	6.2	2.7	2.4
Nonmetro	19.1	18.3	5.1	5.2	2.3	1.9
Urbanized	21.3	20.6	6.5	5.6	2.9	2.0
Less Urbanized	17.1	16.8	4.2	5.2	2.0	1.9
Completely Rural	19.2	15.3	3.7	3.4	1.6	1.2

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.