

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 1,036 | 956 | 603 | 648 | 460 | 492 |
| 12 | 28 | 25 | 20 | 20 | 15 | 14 |
| 13 | 35 | 31 | 27 | 26 | 19 | 20 |
| 14 | 40 | 41 | 34 | 33 | 24 | 23 |
| 15 | 50 | 46 | 46 | 39 | 32 | 25 |
| 16 | 57 | 55 | 49 | 49 | 37 | 36 |
| 17 | 58 | 61 | 49 | 53 | 38 | 40 |
| 18 | 100 | 104 | 85 | 95 | 65 | 65 |
| 19 | 96 | 98 | 82 | 82 | 61 | 67 |
| 20 | 110 | 105 | 91 | 92 | 68 | 64 |
| 21 | 88 | 96 | 67 | 79 | 53 | 58 |
| 22 | 103 | 94 | 80 | 77 | 57 | 58 |
| 23 | 84 | 92 | 63 | 69 | 48 | 46 |
| 24 | 88 | 82 | 61 | 60 | 46 | 47 |
| 25 | 94 | 86 | 59 | 60 | 50 | 45 |
| 26-29 | 345 | 348 | 223 | 217 | 173 | 162 |
| 30-34 | 375 | 372 | 207 | 211 | 154 | 163 |
| 35-39 | 367 | 370 | 183 | 181 | 138 | 134 |
| 40-44 | 374 | 346 | 161 | 187 | 119 | 131 |
| 45-49 | 384 | 378 | 167 | 177 | 124 | 137 |
| 50-54 | 594 | 550 | 220 | 244 | 173 | 201 |
| 55-59 | 454 | 519 | 191 | 228 | 153 | 174 |
| 60-64 | 389 | 445 | 132 | 124 | 87 | 97 |
| 65 or Older | 343 | 385 | 109 | 151 | 85 | 109 |

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.40 | 0.37 | 0.23 | 0.25 | 0.18 | 0.19 |
| 12 | 0.72 | 0.58 | 0.53 | 0.47 | 0.39 | 0.33 |
| 13 | 0.75 | 0.72 | 0.60 | 0.62 | 0.43 | 0.49 |
| 14 | 0.84 | 0.86 | 0.74 | 0.72 | 0.54 | 0.52 |
| 15 | 0.96 | 0.92 | 0.93 | 0.81 | 0.71 | 0.56 |
| 16 | 1.09 | 1.00 | 0.98 | 0.96 | 0.76 | 0.76 |
| 17 | 1.08 | 1.07 | 1.00 | 1.01 | 0.82 | 0.83 |
| 18 | 1.28 | 1.18 | 1.19 | 1.20 | 1.04 | 0.98 |
| 19 | 1.36 | 1.40 | 1.34 | 1.35 | 1.15 | 1.23 |
| 20 | 1.32 | 1.28 | 1.33 | 1.34 | 1.17 | 1.15 |
| 21 | 1.28 | 1.25 | 1.18 | 1.37 | 1.03 | 1.14 |
| 22 | 1.28 | 1.22 | 1.31 | 1.28 | 1.11 | 1.08 |
| 23 | 1.27 | 1.26 | 1.30 | 1.25 | 1.07 | 0.94 |
| 24 | 1.27 | 1.21 | 1.24 | 1.16 | 1.02 | 1.02 |
| 25 | 1.26 | 1.32 | 1.18 | 1.24 | 1.10 | 0.99 |
| 26-29 | 1.27 | 1.23 | 1.09 | 1.08 | 0.91 | 0.87 |
| 30-34 | 1.15 | 1.15 | 0.94 | 0.91 | 0.75 | 0.76 |
| 35-39 | 1.21 | 1.26 | 0.86 | 0.85 | 0.68 | 0.66 |
| 40-44 | 1.22 | 1.15 | 0.75 | 0.85 | 0.56 | 0.62 |
| 45-49 | 1.15 | 1.16 | 0.73 | 0.77 | 0.55 | 0.62 |
| 50-54 | 1.58 | 1.47 | 0.91 | 0.99 | 0.73 | 0.84 |
| 55-59 | 1.62 | 1.68 | 0.94 | 1.05 | 0.77 | 0.82 |
| 60-64 | 1.62 | 1.74 | 0.70 | 0.73 | 0.47 | 0.57 |
| 65 or Older | 0.85 | 0.92 | 0.27 | 0.36 | 0.21 | 0.26 |

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.0649 | | 0.0009 | | 0.0723 | |
| 12 | 0.2525 | | 0.7651 | | 0.6907 | |
| 13 | 0.2833 | | 0.8684 | | 0.3316 | |
| 14 | 0.3927 | | 0.2110 | | 0.9909 | |
| 15 | 0.0829 | | 0.0042 | | 0.0180 | |
| 16 | 0.3166 | | 0.5454 | | 0.3018 | |
| 17 | 0.8686 | | 0.8121 | | 0.9196 | |
| 18 | 0.2209 | | 0.3335 | | 0.9316 | |
| 19 | 0.9883 | | 0.6526 | | 0.7310 | |
| 20 | 0.9838 | | 0.3036 | | 0.9290 | |
| 21 | 0.2085 | | 0.1219 | | 0.9472 | |
| 22 | 0.7296 | | 0.5005 | | 0.3837 | |
| 23 | 0.7825 | | 0.4422 | | 0.0643 | |
| 24 | 0.5361 | | 0.1196 | | 0.3273 | |
| 25 | 0.6620 | | 0.4371 | | 0.4545 | |
| 26-29 | 0.1323 | | 0.0556 | | 0.8349 | |
| 30-34 | 1.0000 | | 0.1406 | | 0.0474 | |
| 35-39 | 0.5814 | | 0.3523 | | 0.4754 | |
| 40-44 | 0.5783 | | 0.0610 | | 0.2794 | |
| 45-49 | 0.3257 | | 0.0413 | | 0.2376 | |
| 50-54 | 0.5851 | | 0.3563 | | 0.6428 | |
| 55-59 | 0.7496 | | 0.3602 | | 0.5774 | |
| 60-64 | 0.0152 | | 0.9274 | | 0.2373 | |
| 65 or Older | 0.0285 | | 0.1680 | | 0.4229 | |

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 1,013 | 1,005 | 514 | 556 | 409 | 431 |
| 12 | 11 | 11 | 10 | 10 | 6 | 3 |
| 13 | 23 | 20 | 19 | 18 | 15 | 13 |
| 14 | 27 | 29 | 26 | 25 | 19 | 18 |
| 15 | 45 | 38 | 42 | 34 | 28 | 24 |
| 16 | 51 | 50 | 45 | 45 | 34 | 31 |
| 17 | 54 | 57 | 47 | 49 | 36 | 36 |
| 18 | 95 | 97 | 82 | 90 | 61 | 61 |
| 19 | 91 | 90 | 78 | 77 | 59 | 62 |
| 20 | 107 | 101 | 87 | 84 | 66 | 61 |
| 21 | 83 | 89 | 62 | 73 | 51 | 53 |
| 22 | 100 | 90 | 71 | 72 | 53 | 54 |
| 23 | 82 | 86 | 59 | 65 | 45 | 42 |
| 24 | 85 | 80 | 57 | 57 | 42 | 44 |
| 25 | 90 | 81 | 54 | 54 | 45 | 42 |
| 26-29 | 325 | 330 | 201 | 187 | 158 | 144 |
| 30-34 | 352 | 347 | 176 | 183 | 133 | 145 |
| 35-39 | 348 | 347 | 153 | 153 | 121 | 115 |
| 40-44 | 351 | 317 | 131 | 139 | 99 | 105 |
| 45-49 | 363 | 365 | 133 | 149 | 101 | 111 |
| 50-54 | 568 | 535 | 188 | 199 | 149 | 166 |
| 55-59 | 433 | 498 | 154 | 186 | 133 | 151 |
| 60-64 | 370 | 440 | 119 | 115 | 85 | 89 |
| 65 or Older | 316 | 338 | 89 | 108 | 77 | 99 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.39 | 0.39 | 0.20 | 0.21 | 0.16 | 0.17 |
| 12 | 0.31 | 0.26 | 0.28 | 0.24 | 0.15 | 0.08 |
| 13 | 0.53 | 0.49 | 0.45 | 0.43 | 0.36 | 0.32 |
| 14 | 0.61 | 0.64 | 0.58 | 0.57 | 0.45 | 0.41 |
| 15 | 0.92 | 0.80 | 0.88 | 0.73 | 0.62 | 0.54 |
| 16 | 1.01 | 0.96 | 0.91 | 0.91 | 0.72 | 0.67 |
| 17 | 1.05 | 1.06 | 0.96 | 0.98 | 0.80 | 0.78 |
| 18 | 1.29 | 1.23 | 1.18 | 1.24 | 0.99 | 0.97 |
| 19 | 1.37 | 1.40 | 1.31 | 1.31 | 1.11 | 1.15 |
| 20 | 1.36 | 1.30 | 1.34 | 1.30 | 1.15 | 1.11 |
| 21 | 1.27 | 1.27 | 1.14 | 1.29 | 1.00 | 1.06 |
| 22 | 1.30 | 1.27 | 1.24 | 1.25 | 1.04 | 1.05 |
| 23 | 1.32 | 1.24 | 1.25 | 1.21 | 1.02 | 0.88 |
| 24 | 1.34 | 1.25 | 1.18 | 1.15 | 0.93 | 0.98 |
| 25 | 1.30 | 1.34 | 1.14 | 1.18 | 1.02 | 0.95 |
| 26-29 | 1.28 | 1.26 | 1.01 | 0.97 | 0.85 | 0.79 |
| 30-34 | 1.17 | 1.17 | 0.83 | 0.84 | 0.65 | 0.69 |
| 35-39 | 1.23 | 1.29 | 0.73 | 0.74 | 0.59 | 0.57 |
| 40-44 | 1.20 | 1.16 | 0.62 | 0.66 | 0.47 | 0.51 |
| 45-49 | 1.11 | 1.20 | 0.59 | 0.67 | 0.46 | 0.51 |
| 50-54 | 1.59 | 1.52 | 0.79 | 0.83 | 0.63 | 0.69 |
| 55-59 | 1.62 | 1.67 | 0.78 | 0.89 | 0.68 | 0.72 |
| 60-64 | 1.58 | 1.82 | 0.64 | 0.68 | 0.46 | 0.53 |
| 65 or Older | 0.79 | 0.81 | 0.22 | 0.26 | 0.19 | 0.24 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.0949 | | 0.0450 | | 0.2989 | |
| 12 | 0.7531 | | 0.7973 | | 0.2770 | |
| 13 | 0.9597 | | 0.9262 | | 0.9413 | |
| 14 | 0.8903 | | 0.6343 | | 0.7710 | |
| 15 | 0.0768 | | 0.0059 | | 0.2389 | |
| 16 | 0.6498 | | 0.6373 | | 0.1289 | |
| 17 | 0.7718 | | 0.7746 | | 0.6019 | |
| 18 | 0.6736 | | 0.6258 | | 0.7755 | |
| 19 | 0.4558 | | 0.9011 | | 0.8578 | |
| 20 | 0.9739 | | 0.6485 | | 0.7675 | |
| 21 | 0.5448 | | 0.1654 | | 0.5291 | |
| 22 | 0.9073 | | 0.7452 | | 0.6027 | |
| 23 | 0.7328 | | 0.5724 | | 0.1580 | |
| 24 | 0.3680 | | 0.1374 | | 0.2067 | |
| 25 | 0.5769 | | 0.8618 | | 0.4701 | |
| 26-29 | 0.1083 | | 0.5581 | | 0.7227 | |
| 30-34 | 0.7678 | | 0.0487 | | 0.0110 | |
| 35-39 | 0.4172 | | 0.5124 | | 0.6483 | |
| 40-44 | 0.3015 | | 0.1889 | | 0.3860 | |
| 45-49 | 0.5360 | | 0.0507 | | 0.5485 | |
| 50-54 | 0.8019 | | 0.9286 | | 0.8108 | |
| 55-59 | 0.5826 | | 0.7265 | | 0.7845 | |
| 60-64 | 0.0188 | | 0.9836 | | 0.7248 | |
| 65 or Older | 0.1120 | | 0.6757 | | 0.5986 | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 701 | 703 | 168 | 217 | 106 | 142 |
| 12 | 2 | 1 | 1 | 1 | * | 1 |
| 13 | 3 | 4 | 3 | 4 | 1 | * |
| 14 | 7 | 6 | 6 | 5 | 1 | 1 |
| 15 | 17 | 9 | 10 | 6 | 7 | 4 |
| 16 | 12 | 11 | 10 | 10 | 6 | 2 |
| 17 | 14 | 17 | 12 | 14 | 7 | 5 |
| 18 | 29 | 31 | 22 | 24 | 15 | 15 |
| 19 | 32 | 35 | 24 | 24 | 12 | 10 |
| 20 | 39 | 43 | 29 | 31 | 19 | 13 |
| 21 | 37 | 39 | 26 | 23 | 15 | 9 |
| 22 | 42 | 43 | 25 | 33 | 11 | 17 |
| 23 | 45 | 45 | 22 | 25 | 11 | 14 |
| 24 | 41 | 41 | 23 | 21 | 12 | 8 |
| 25 | 47 | 42 | 28 | 24 | 10 | 10 |
| 26-29 | 183 | 178 | 60 | 77 | 35 | 40 |
| 30-34 | 203 | 201 | 68 | 87 | 30 | 64 |
| 35-39 | 204 | 200 | 56 | 68 | 43 | 48 |
| 40-44 | 205 | 199 | 50 | 59 | 27 | 31 |
| 45-49 | 247 | 245 | 60 | 75 | 36 | 58 |
| 50-54 | 386 | 377 | 60 | 56 | 43 | 38 |
| 55-59 | 274 | 312 | 28 | 79 | 24 | 54 |
| 60-64 | 195 | 206 | 34 | 33 | 26 | 44 |
| 65 or Older | 173 | 156 | 6 | * | 5 | * |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.27 | 0.27 | 0.07 | 0.08 | 0.04 | 0.05 |
| 12 | 0.06 | 0.03 | 0.03 | 0.03 | * | 0.02 |
| 13 | 0.08 | 0.10 | 0.06 | 0.10 | 0.02 | * |
| 14 | 0.15 | 0.14 | 0.14 | 0.11 | 0.02 | 0.03 |
| 15 | 0.40 | 0.20 | 0.24 | 0.14 | 0.17 | 0.10 |
| 16 | 0.28 | 0.27 | 0.24 | 0.24 | 0.14 | 0.05 |
| 17 | 0.33 | 0.38 | 0.28 | 0.32 | 0.16 | 0.11 |
| 18 | 0.59 | 0.63 | 0.45 | 0.50 | 0.32 | 0.33 |
| 19 | 0.74 | 0.80 | 0.54 | 0.56 | 0.27 | 0.23 |
| 20 | 0.83 | 0.91 | 0.63 | 0.67 | 0.42 | 0.30 |
| 21 | 0.78 | 0.84 | 0.56 | 0.52 | 0.33 | 0.21 |
| 22 | 0.87 | 0.89 | 0.54 | 0.70 | 0.25 | 0.37 |
| 23 | 1.03 | 0.94 | 0.52 | 0.55 | 0.28 | 0.32 |
| 24 | 0.92 | 0.90 | 0.54 | 0.50 | 0.30 | 0.20 |
| 25 | 1.07 | 0.95 | 0.68 | 0.59 | 0.26 | 0.25 |
| 26-29 | 0.97 | 0.89 | 0.35 | 0.44 | 0.21 | 0.23 |
| 30-34 | 0.94 | 0.90 | 0.34 | 0.43 | 0.16 | 0.32 |
| 35-39 | 0.94 | 0.93 | 0.28 | 0.35 | 0.22 | 0.24 |
| 40-44 | 0.93 | 0.90 | 0.24 | 0.29 | 0.13 | 0.16 |
| 45-49 | 0.98 | 0.99 | 0.28 | 0.34 | 0.17 | 0.27 |
| 50-54 | 1.40 | 1.35 | 0.26 | 0.24 | 0.18 | 0.16 |
| 55-59 | 1.28 | 1.36 | 0.15 | 0.38 | 0.13 | 0.26 |
| 60-64 | 1.04 | 1.12 | 0.19 | 0.20 | 0.15 | 0.26 |
| 65 or Older | 0.43 | 0.37 | 0.01 | * | 0.01 | * |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

130401

Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.6718 | | 0.0049 | | 0.1269 | |
| 12 | 0.6208 | | 0.5401 | | * | |
| 13 | 0.5357 | | 0.9481 | | * | |
| 14 | 0.8143 | | 0.5540 | | 0.6950 | |
| 15 | 0.0524 | | 0.0320 | | 0.1247 | |
| 16 | 0.1955 | | 0.4134 | | 0.0541 | |
| 17 | 0.8192 | | 0.7133 | | 0.1222 | |
| 18 | 0.5788 | | 0.4801 | | 0.9762 | |
| 19 | 0.8138 | | 0.8291 | | 0.1868 | |
| 20 | 0.6291 | | 0.6894 | | 0.0248 | |
| 21 | 0.9794 | | 0.3997 | | 0.2034 | |
| 22 | 0.3961 | | 0.1072 | | 0.1617 | |
| 23 | 0.4032 | | 0.6023 | | 0.8814 | |
| 24 | 0.1353 | | 0.6066 | | 0.2071 | |
| 25 | 0.0571 | | 0.7627 | | 0.7147 | |
| 26-29 | 0.1066 | | 0.0125 | | 0.6393 | |
| 30-34 | 0.3526 | | 0.5940 | | 0.1082 | |
| 35-39 | 0.8041 | | 0.3043 | | 0.8030 | |
| 40-44 | 0.4880 | | 0.2936 | | 0.3188 | |
| 45-49 | 0.6063 | | 0.2964 | | 0.2641 | |
| 50-54 | 0.2860 | | 0.6968 | | 0.8196 | |
| 55-59 | 0.7082 | | 0.0405 | | 0.3215 | |
| 60-64 | 0.5119 | | 0.3265 | | 0.4055 | |
| 65 or Older | 0.8928 | | * | | * | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 327 | 346 | 69 | 111 | 42 | 94 |
| 12 | 1 | * | 1 | * | * | * |
| 13 | * | 0 | * | 0 | * | * |
| 14 | 3 | 1 | 3 | 1 | * | * |
| 15 | 14 | 4 | 2 | 3 | 2 | * |
| 16 | 4 | 7 | 4 | 5 | 0 | 1 |
| 17 | 6 | 6 | 4 | 5 | 1 | * |
| 18 | 5 | 10 | 3 | 6 | 2 | 4 |
| 19 | 9 | 10 | 5 | 7 | 2 | 2 |
| 20 | 16 | 15 | 8 | 8 | 4 | 4 |
| 21 | 11 | 12 | 5 | 4 | 1 | 2 |
| 22 | 15 | 16 | 6 | 6 | 2 | 3 |
| 23 | 17 | 17 | 6 | 5 | 2 | 3 |
| 24 | 19 | 14 | 5 | 6 | 1 | 3 |
| 25 | 19 | 17 | 5 | 6 | 2 | 3 |
| 26-29 | 74 | 86 | 13 | 25 | 7 | 8 |
| 30-34 | 95 | 114 | 19 | 52 | 9 | 46 |
| 35-39 | 108 | 99 | 26 | 32 | 9 | 28 |
| 40-44 | 103 | 101 | 35 | 24 | 11 | 12 |
| 45-49 | 110 | 129 | 32 | 51 | 27 | 42 |
| 50-54 | 185 | 177 | 24 | 42 | 22 | 29 |
| 55-59 | 120 | 130 | 14 | 47 | 12 | 46 |
| 60-64 | 90 | 93 | 16 | 19 | * | 13 |
| 65 or Older | 43 | 48 | 5 | * | 5 | * |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.13 | 0.13 | 0.03 | 0.04 | 0.02 | 0.04 |
| 12 | 0.03 | * | 0.03 | * | * | * |
| 13 | * | 0.01 | * | 0.01 | * | * |
| 14 | 0.08 | 0.03 | 0.06 | 0.02 | * | * |
| 15 | 0.32 | 0.11 | 0.05 | 0.06 | 0.04 | * |
| 16 | 0.09 | 0.15 | 0.09 | 0.13 | 0.01 | 0.02 |
| 17 | 0.14 | 0.13 | 0.10 | 0.12 | 0.03 | * |
| 18 | 0.12 | 0.20 | 0.06 | 0.14 | 0.05 | 0.09 |
| 19 | 0.21 | 0.25 | 0.13 | 0.17 | 0.04 | 0.04 |
| 20 | 0.37 | 0.35 | 0.17 | 0.19 | 0.08 | 0.08 |
| 21 | 0.25 | 0.27 | 0.11 | 0.10 | 0.03 | 0.05 |
| 22 | 0.34 | 0.35 | 0.15 | 0.13 | 0.05 | 0.08 |
| 23 | 0.43 | 0.37 | 0.15 | 0.11 | 0.04 | 0.07 |
| 24 | 0.45 | 0.34 | 0.11 | 0.14 | 0.02 | 0.08 |
| 25 | 0.46 | 0.42 | 0.13 | 0.14 | 0.05 | 0.07 |
| 26-29 | 0.43 | 0.49 | 0.08 | 0.14 | 0.04 | 0.05 |
| 30-34 | 0.47 | 0.55 | 0.10 | 0.26 | 0.04 | 0.23 |
| 35-39 | 0.54 | 0.50 | 0.13 | 0.16 | 0.04 | 0.15 |
| 40-44 | 0.50 | 0.49 | 0.17 | 0.12 | 0.05 | 0.06 |
| 45-49 | 0.49 | 0.59 | 0.15 | 0.23 | 0.13 | 0.20 |
| 50-54 | 0.76 | 0.73 | 0.10 | 0.18 | 0.09 | 0.12 |
| 55-59 | 0.62 | 0.63 | 0.07 | 0.23 | 0.06 | 0.22 |
| 60-64 | 0.50 | 0.55 | 0.09 | 0.12 | * | 0.07 |
| 65 or Older | 0.11 | 0.11 | 0.01 | * | 0.01 | * |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.1175 | | 0.0321 | | 0.0405 | |
| 12 | * | | * | | * | |
| 13 | * | | * | | * | |
| 14 | 0.3474 | | 0.5542 | | * | |
| 15 | 0.1689 | | 0.8179 | | * | |
| 16 | 0.8013 | | 0.8455 | | 0.5203 | |
| 17 | 0.1429 | | 0.4407 | | * | |
| 18 | 0.0784 | | 0.1280 | | 0.4323 | |
| 19 | 0.8962 | | 0.6158 | | 0.7789 | |
| 20 | 0.4568 | | 0.4738 | | 0.9138 | |
| 21 | 0.8092 | | 0.9293 | | 0.2939 | |
| 22 | 0.5920 | | 0.9474 | | 0.4301 | |
| 23 | 0.5692 | | 0.9487 | | 0.7299 | |
| 24 | 0.3651 | | 0.3494 | | 0.0759 | |
| 25 | 0.3424 | | 0.7718 | | 0.7344 | |
| 26-29 | 0.4905 | | 0.1363 | | 0.4725 | |
| 30-34 | 0.0292 | | 0.1307 | | 0.1543 | |
| 35-39 | 0.9230 | | 0.9375 | | 0.2425 | |
| 40-44 | 0.5525 | | 0.6733 | | 0.9283 | |
| 45-49 | 0.1062 | | 0.4038 | | 0.7798 | |
| 50-54 | 0.6310 | | 0.1768 | | 0.5925 | |
| 55-59 | 0.6919 | | 0.3544 | | 0.3508 | |
| 60-64 | 0.3585 | | 0.8285 | | * | |
| 65 or Older | 0.6500 | | * | | * | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 628 | 674 | 143 | 175 | 60 | 90 |
| 12 | 5 | 4 | 2 | 3 | 1 | 3 |
| 13 | 11 | 8 | 5 | 7 | 2 | 4 |
| 14 | 14 | 14 | 12 | 11 | 9 | 6 |
| 15 | 20 | 15 | 16 | 11 | 10 | 6 |
| 16 | 25 | 21 | 22 | 19 | 15 | 10 |
| 17 | 24 | 26 | 19 | 19 | 10 | 10 |
| 18 | 40 | 42 | 32 | 31 | 14 | 22 |
| 19 | 42 | 47 | 33 | 36 | 16 | 18 |
| 20 | 48 | 57 | 36 | 44 | 14 | 15 |
| 21 | 49 | 49 | 27 | 27 | 17 | 13 |
| 22 | 51 | 55 | 30 | 29 | 15 | 16 |
| 23 | 48 | 52 | 27 | 26 | 10 | 15 |
| 24 | 48 | 47 | 22 | 21 | 11 | 8 |
| 25 | 49 | 54 | 25 | 26 | 13 | 9 |
| 26-29 | 201 | 204 | 58 | 74 | 18 | 32 |
| 30-34 | 219 | 224 | 49 | 69 | 26 | 45 |
| 35-39 | 181 | 216 | 51 | 57 | 13 | 26 |
| 40-44 | 191 | 183 | 34 | 29 | 5 | 10 |
| 45-49 | 214 | 210 | 32 | 38 | 15 | 35 |
| 50-54 | 315 | 294 | 26 | 26 | * | 26 |
| 55-59 | 243 | 292 | 6 | 11 | 2 | 4 |
| 60-64 | 192 | 210 | * | 44 | * | * |
| 65 or Older | 141 | 197 | 5 | 46 | * | 13 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.24 | 0.26 | 0.06 | 0.07 | 0.02 | 0.03 |
| 12 | 0.14 | 0.10 | 0.06 | 0.08 | 0.02 | 0.08 |
| 13 | 0.25 | 0.19 | 0.11 | 0.17 | 0.05 | 0.10 |
| 14 | 0.32 | 0.32 | 0.29 | 0.26 | 0.21 | 0.15 |
| 15 | 0.46 | 0.36 | 0.38 | 0.26 | 0.22 | 0.15 |
| 16 | 0.55 | 0.49 | 0.49 | 0.44 | 0.34 | 0.24 |
| 17 | 0.55 | 0.57 | 0.44 | 0.42 | 0.23 | 0.23 |
| 18 | 0.74 | 0.82 | 0.61 | 0.64 | 0.28 | 0.45 |
| 19 | 0.88 | 0.96 | 0.72 | 0.80 | 0.36 | 0.41 |
| 20 | 0.97 | 1.10 | 0.74 | 0.91 | 0.32 | 0.33 |
| 21 | 1.00 | 0.97 | 0.59 | 0.58 | 0.37 | 0.30 |
| 22 | 1.02 | 1.04 | 0.64 | 0.63 | 0.34 | 0.36 |
| 23 | 1.05 | 1.03 | 0.66 | 0.57 | 0.25 | 0.33 |
| 24 | 1.03 | 1.00 | 0.53 | 0.50 | 0.25 | 0.21 |
| 25 | 1.14 | 1.16 | 0.61 | 0.62 | 0.33 | 0.22 |
| 26-29 | 1.05 | 1.04 | 0.33 | 0.43 | 0.11 | 0.19 |
| 30-34 | 0.94 | 0.97 | 0.25 | 0.35 | 0.13 | 0.23 |
| 35-39 | 0.86 | 1.00 | 0.26 | 0.29 | 0.07 | 0.13 |
| 40-44 | 0.84 | 0.81 | 0.16 | 0.14 | 0.03 | 0.05 |
| 45-49 | 0.88 | 0.88 | 0.15 | 0.17 | 0.07 | 0.16 |
| 50-54 | 1.23 | 1.14 | 0.11 | 0.11 | * | 0.11 |
| 55-59 | 1.18 | 1.30 | 0.03 | 0.05 | 0.01 | 0.02 |
| 60-64 | 1.02 | 1.13 | * | 0.26 | * | * |
| 65 or Older | 0.35 | 0.47 | 0.01 | 0.11 | * | 0.03 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

130401

Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.1960 | | 0.3911 | | 0.1576 | |
| 12 | 0.8731 | | 0.4069 | | 0.1405 | |
| 13 | 0.4402 | | 0.3730 | | 0.1371 | |
| 14 | 0.4737 | | 0.6969 | | 0.6577 | |
| 15 | 0.0887 | | 0.0369 | | 0.0544 | |
| 16 | 0.0378 | | 0.0946 | | 0.0579 | |
| 17 | 0.8362 | | 0.5296 | | 0.9016 | |
| 18 | 0.5676 | | 0.9870 | | 0.0245 | |
| 19 | 0.9423 | | 0.5801 | | 0.7959 | |
| 20 | 1.0000 | | 0.8882 | | 0.7245 | |
| 21 | 0.4692 | | 0.4151 | | 0.2927 | |
| 22 | 0.2608 | | 0.3007 | | 0.8043 | |
| 23 | 0.1057 | | 0.3203 | | 0.2592 | |
| 24 | 0.3032 | | 0.9120 | | 0.2459 | |
| 25 | 0.8058 | | 0.5461 | | 0.2735 | |
| 26-29 | 0.1549 | | 0.0496 | | 0.2182 | |
| 30-34 | 0.6428 | | 0.5001 | | 0.9701 | |
| 35-39 | 0.0351 | | 0.6398 | | 0.2660 | |
| 40-44 | 0.3797 | | 0.5939 | | 0.2406 | |
| 45-49 | 0.7900 | | 0.5274 | | 0.4266 | |
| 50-54 | 0.1079 | | 0.9214 | | * | |
| 55-59 | 0.4051 | | 0.6101 | | 0.8059 | |
| 60-64 | 0.0781 | | * | | * | |
| 65 or Older | 0.2045 | | 0.2636 | | * | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 491 | 489 | 107 | 120 | 68 | 65 |
| 12 | 22 | 18 | 14 | 11 | 9 | 7 |
| 13 | 24 | 20 | 17 | 14 | 10 | 10 |
| 14 | 26 | 25 | 20 | 18 | 10 | 8 |
| 15 | 24 | 26 | 15 | 14 | 6 | 5 |
| 16 | 26 | 23 | 17 | 15 | 6 | 7 |
| 17 | 24 | 23 | 15 | 13 | 8 | 9 |
| 18 | 31 | 37 | 18 | 21 | 5 | 13 |
| 19 | 32 | 32 | 20 | 13 | 17 | 5 |
| 20 | 33 | 36 | 16 | 14 | 11 | 8 |
| 21 | 29 | 31 | 14 | 18 | 11 | 13 |
| 22 | 40 | 33 | 19 | 14 | 8 | 6 |
| 23 | 35 | 30 | 11 | 8 | 8 | 3 |
| 24 | 35 | 30 | 8 | 4 | 3 | 1 |
| 25 | 35 | 36 | 5 | 10 | 2 | 4 |
| 26-29 | 141 | 136 | 33 | 40 | 15 | 15 |
| 30-34 | 136 | 179 | 30 | 23 | 25 | 10 |
| 35-39 | 162 | 151 | 20 | 35 | 9 | 14 |
| 40-44 | 157 | 162 | 25 | 60 | 22 | 5 |
| 45-49 | 170 | 153 | 23 | 45 | 23 | 27 |
| 50-54 | 193 | 202 | 53 | 25 | 15 | 21 |
| 55-59 | 195 | 210 | 23 | 37 | 23 | 37 |
| 60-64 | 99 | 116 | 24 | 9 | 10 | * |
| 65 or Older | 109 | 144 | 11 | * | 11 | * |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.19 | 0.19 | 0.04 | 0.05 | 0.03 | 0.02 |
| 12 | 0.56 | 0.42 | 0.37 | 0.27 | 0.23 | 0.17 |
| 13 | 0.55 | 0.48 | 0.40 | 0.35 | 0.24 | 0.24 |
| 14 | 0.59 | 0.57 | 0.46 | 0.41 | 0.23 | 0.19 |
| 15 | 0.53 | 0.58 | 0.36 | 0.33 | 0.15 | 0.12 |
| 16 | 0.58 | 0.53 | 0.39 | 0.35 | 0.14 | 0.17 |
| 17 | 0.56 | 0.51 | 0.34 | 0.30 | 0.18 | 0.21 |
| 18 | 0.61 | 0.69 | 0.38 | 0.43 | 0.11 | 0.28 |
| 19 | 0.72 | 0.70 | 0.46 | 0.30 | 0.39 | 0.12 |
| 20 | 0.71 | 0.78 | 0.35 | 0.32 | 0.24 | 0.18 |
| 21 | 0.64 | 0.67 | 0.32 | 0.39 | 0.25 | 0.29 |
| 22 | 0.82 | 0.72 | 0.42 | 0.31 | 0.19 | 0.13 |
| 23 | 0.82 | 0.64 | 0.26 | 0.17 | 0.20 | 0.07 |
| 24 | 0.77 | 0.71 | 0.20 | 0.11 | 0.08 | 0.02 |
| 25 | 0.85 | 0.84 | 0.12 | 0.25 | 0.06 | 0.09 |
| 26-29 | 0.78 | 0.74 | 0.20 | 0.23 | 0.09 | 0.09 |
| 30-34 | 0.66 | 0.83 | 0.15 | 0.11 | 0.13 | 0.05 |
| 35-39 | 0.79 | 0.74 | 0.10 | 0.18 | 0.05 | 0.07 |
| 40-44 | 0.73 | 0.77 | 0.12 | 0.29 | 0.11 | 0.02 |
| 45-49 | 0.73 | 0.70 | 0.10 | 0.21 | 0.10 | 0.13 |
| 50-54 | 0.80 | 0.82 | 0.23 | 0.11 | 0.06 | 0.09 |
| 55-59 | 0.98 | 0.98 | 0.12 | 0.18 | 0.12 | 0.18 |
| 60-64 | 0.55 | 0.68 | 0.13 | 0.06 | 0.05 | * |
| 65 or Older | 0.27 | 0.34 | 0.03 | * | 0.03 | * |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

130401

Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.5982 | | 0.2371 | | 0.1275 | |
| 12 | 0.1440 | | 0.2117 | | 0.5226 | |
| 13 | 0.0091 | | 0.0482 | | 0.8666 | |
| 14 | 0.3299 | | 0.1131 | | 0.4565 | |
| 15 | 0.5711 | | 0.0883 | | 0.1615 | |
| 16 | 0.2794 | | 0.6351 | | 0.9928 | |
| 17 | 0.1694 | | 0.3302 | | 0.6917 | |
| 18 | 0.5438 | | 0.8557 | | 0.2263 | |
| 19 | 0.9866 | | 0.3812 | | 0.2210 | |
| 20 | 0.2979 | | 0.3841 | | 0.4106 | |
| 21 | 0.7883 | | 0.2991 | | 0.5894 | |
| 22 | 0.2263 | | 0.5006 | | 0.5438 | |
| 23 | 0.0243 | | 0.2563 | | 0.2596 | |
| 24 | 0.6910 | | 0.0546 | | 0.2043 | |
| 25 | 0.7725 | | 0.0060 | | 0.3272 | |
| 26-29 | 0.7991 | | 0.7763 | | 0.6422 | |
| 30-34 | 0.0102 | | 0.4761 | | 0.0598 | |
| 35-39 | 0.9212 | | 0.2570 | | 0.9804 | |
| 40-44 | 0.7554 | | 0.7196 | | 0.0985 | |
| 45-49 | 0.5746 | | 0.1537 | | 0.9650 | |
| 50-54 | 0.6282 | | 0.1813 | | 0.6620 | |
| 55-59 | 0.9237 | | 0.9254 | | 0.9254 | |
| 60-64 | 0.1069 | | 0.5879 | | * | |
| 65 or Older | 0.1721 | | * | | * | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 752 | 765 | 363 | 406 | 231 | 265 |
| 12 | 19 | 18 | 15 | 16 | 10 | 12 |
| 13 | 19 | 19 | 14 | 17 | 8 | 12 |
| 14 | 23 | 24 | 19 | 18 | 12 | 13 |
| 15 | 28 | 30 | 25 | 21 | 15 | 13 |
| 16 | 35 | 34 | 30 | 30 | 17 | 19 |
| 17 | 37 | 39 | 28 | 32 | 17 | 19 |
| 18 | 54 | 66 | 39 | 50 | 24 | 22 |
| 19 | 53 | 57 | 41 | 42 | 27 | 23 |
| 20 | 65 | 74 | 46 | 56 | 26 | 31 |
| 21 | 57 | 63 | 39 | 45 | 22 | 24 |
| 22 | 63 | 67 | 45 | 52 | 28 | 32 |
| 23 | 57 | 62 | 35 | 40 | 21 | 26 |
| 24 | 63 | 56 | 38 | 35 | 25 | 23 |
| 25 | 60 | 62 | 34 | 35 | 22 | 25 |
| 26-29 | 246 | 240 | 123 | 144 | 87 | 85 |
| 30-34 | 239 | 254 | 129 | 135 | 85 | 91 |
| 35-39 | 232 | 253 | 112 | 120 | 75 | 80 |
| 40-44 | 225 | 236 | 117 | 136 | 72 | 79 |
| 45-49 | 254 | 257 | 106 | 104 | 68 | 69 |
| 50-54 | 351 | 368 | 126 | 164 | 88 | 117 |
| 55-59 | 300 | 339 | 123 | 151 | 71 | 93 |
| 60-64 | 222 | 232 | 62 | 63 | 29 | 53 |
| 65 or Older | 194 | 260 | 65 | 113 | 35 | 45 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.29 | 0.29 | 0.14 | 0.16 | 0.09 | 0.10 |
| 12 | 0.49 | 0.43 | 0.38 | 0.38 | 0.26 | 0.29 |
| 13 | 0.44 | 0.48 | 0.33 | 0.41 | 0.20 | 0.31 |
| 14 | 0.53 | 0.54 | 0.43 | 0.42 | 0.29 | 0.30 |
| 15 | 0.61 | 0.66 | 0.56 | 0.49 | 0.34 | 0.30 |
| 16 | 0.73 | 0.74 | 0.65 | 0.66 | 0.39 | 0.44 |
| 17 | 0.80 | 0.82 | 0.62 | 0.70 | 0.40 | 0.44 |
| 18 | 0.92 | 1.01 | 0.72 | 0.81 | 0.49 | 0.45 |
| 19 | 1.05 | 1.09 | 0.86 | 0.86 | 0.59 | 0.52 |
| 20 | 1.16 | 1.29 | 0.95 | 1.09 | 0.57 | 0.68 |
| 21 | 1.07 | 1.18 | 0.82 | 0.92 | 0.48 | 0.52 |
| 22 | 1.19 | 1.18 | 0.92 | 1.05 | 0.61 | 0.68 |
| 23 | 1.22 | 1.18 | 0.85 | 0.84 | 0.53 | 0.57 |
| 24 | 1.28 | 1.12 | 0.87 | 0.79 | 0.59 | 0.56 |
| 25 | 1.25 | 1.26 | 0.81 | 0.81 | 0.53 | 0.59 |
| 26-29 | 1.16 | 1.10 | 0.69 | 0.79 | 0.51 | 0.48 |
| 30-34 | 1.00 | 1.02 | 0.63 | 0.63 | 0.43 | 0.45 |
| 35-39 | 1.02 | 1.09 | 0.56 | 0.59 | 0.38 | 0.40 |
| 40-44 | 1.00 | 1.01 | 0.56 | 0.65 | 0.35 | 0.39 |
| 45-49 | 0.98 | 1.03 | 0.48 | 0.47 | 0.31 | 0.32 |
| 50-54 | 1.31 | 1.32 | 0.53 | 0.68 | 0.38 | 0.50 |
| 55-59 | 1.34 | 1.44 | 0.63 | 0.72 | 0.37 | 0.45 |
| 60-64 | 1.14 | 1.22 | 0.34 | 0.37 | 0.16 | 0.31 |
| 65 or Older | 0.48 | 0.62 | 0.16 | 0.27 | 0.09 | 0.11 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.0121 | | 0.0004 | | 0.0593 | |
| 12 | 0.6668 | | 0.7729 | | 0.5262 | |
| 13 | 0.6048 | | 0.2398 | | 0.1750 | |
| 14 | 0.8587 | | 0.5648 | | 0.5674 | |
| 15 | 0.8867 | | 0.1085 | | 0.3328 | |
| 16 | 0.5206 | | 0.1194 | | 0.6291 | |
| 17 | 0.9840 | | 0.9503 | | 0.9865 | |
| 18 | 0.0595 | | 0.0436 | | 0.6160 | |
| 19 | 0.3727 | | 0.8015 | | 0.7441 | |
| 20 | 0.6384 | | 0.3247 | | 0.8112 | |
| 21 | 0.8464 | | 0.9008 | | 0.7687 | |
| 22 | 0.1389 | | 0.0224 | | 0.1482 | |
| 23 | 0.4075 | | 0.9416 | | 0.5783 | |
| 24 | 0.9677 | | 0.6402 | | 0.7687 | |
| 25 | 0.7942 | | 0.5314 | | 0.5780 | |
| 26-29 | 0.0338 | | 0.0370 | | 0.7900 | |
| 30-34 | 0.1879 | | 0.6148 | | 0.9063 | |
| 35-39 | 0.0381 | | 0.2723 | | 0.4453 | |
| 40-44 | 0.5258 | | 0.4217 | | 0.5614 | |
| 45-49 | 0.3635 | | 0.3486 | | 0.6254 | |
| 50-54 | 0.5648 | | 0.0758 | | 0.7248 | |
| 55-59 | 0.6053 | | 0.3505 | | 0.3504 | |
| 60-64 | 0.2080 | | 0.5453 | | 0.0751 | |
| 65 or Older | 0.0832 | | 0.1360 | | 0.3556 | |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 598 | 622 | 302 | 355 | 189 | 211 |
| 12 | 17 | 17 | 14 | 15 | 9 | 11 |
| 13 | 17 | 18 | 13 | 15 | 7 | 11 |
| 14 | 22 | 21 | 16 | 17 | 12 | 12 |
| 15 | 26 | 27 | 23 | 20 | 13 | 12 |
| 16 | 33 | 33 | 28 | 28 | 16 | 19 |
| 17 | 33 | 34 | 26 | 27 | 14 | 15 |
| 18 | 47 | 57 | 33 | 40 | 21 | 19 |
| 19 | 47 | 48 | 35 | 33 | 24 | 20 |
| 20 | 54 | 66 | 38 | 48 | 20 | 27 |
| 21 | 51 | 53 | 35 | 31 | 18 | 18 |
| 22 | 56 | 61 | 38 | 44 | 23 | 27 |
| 23 | 52 | 52 | 32 | 34 | 18 | 22 |
| 24 | 53 | 49 | 34 | 31 | 19 | 18 |
| 25 | 54 | 54 | 31 | 31 | 18 | 22 |
| 26-29 | 226 | 214 | 104 | 127 | 75 | 72 |
| 30-34 | 206 | 213 | 113 | 122 | 71 | 80 |
| 35-39 | 190 | 225 | 96 | 108 | 62 | 71 |
| 40-44 | 178 | 202 | 96 | 122 | 60 | 64 |
| 45-49 | 193 | 204 | 87 | 92 | 58 | 62 |
| 50-54 | 254 | 256 | 94 | 126 | 69 | 86 |
| 55-59 | 219 | 249 | 109 | 131 | 54 | 55 |
| 60-64 | 134 | 149 | 56 | 65 | 28 | 52 |
| 65 or Older | 149 | 159 | 61 | 104 | 32 | 42 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2011 and 2012

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.23 | 0.24 | 0.12 | 0.14 | 0.07 | 0.08 |
| 12 | 0.46 | 0.42 | 0.36 | 0.37 | 0.24 | 0.28 |
| 13 | 0.40 | 0.44 | 0.31 | 0.38 | 0.17 | 0.27 |
| 14 | 0.49 | 0.49 | 0.37 | 0.39 | 0.27 | 0.28 |
| 15 | 0.58 | 0.61 | 0.52 | 0.46 | 0.31 | 0.28 |
| 16 | 0.70 | 0.71 | 0.61 | 0.62 | 0.36 | 0.43 |
| 17 | 0.72 | 0.71 | 0.58 | 0.60 | 0.34 | 0.34 |
| 18 | 0.85 | 0.95 | 0.64 | 0.69 | 0.43 | 0.40 |
| 19 | 0.94 | 1.02 | 0.75 | 0.73 | 0.54 | 0.46 |
| 20 | 1.04 | 1.23 | 0.84 | 0.97 | 0.43 | 0.61 |
| 21 | 0.98 | 1.06 | 0.75 | 0.67 | 0.41 | 0.41 |
| 22 | 1.08 | 1.13 | 0.79 | 0.91 | 0.51 | 0.59 |
| 23 | 1.15 | 1.04 | 0.78 | 0.73 | 0.45 | 0.50 |
| 24 | 1.13 | 1.03 | 0.79 | 0.71 | 0.46 | 0.44 |
| 25 | 1.18 | 1.18 | 0.74 | 0.72 | 0.45 | 0.55 |
| 26-29 | 1.09 | 1.04 | 0.60 | 0.70 | 0.44 | 0.41 |
| 30-34 | 0.94 | 0.92 | 0.56 | 0.58 | 0.36 | 0.39 |
| 35-39 | 0.88 | 1.01 | 0.48 | 0.54 | 0.32 | 0.36 |
| 40-44 | 0.81 | 0.90 | 0.46 | 0.58 | 0.29 | 0.32 |
| 45-49 | 0.83 | 0.87 | 0.40 | 0.42 | 0.27 | 0.29 |
| 50-54 | 1.01 | 1.01 | 0.40 | 0.53 | 0.30 | 0.37 |
| 55-59 | 1.06 | 1.11 | 0.56 | 0.63 | 0.28 | 0.26 |
| 60-64 | 0.73 | 0.83 | 0.31 | 0.38 | 0.15 | 0.31 |
| 65 or Older | 0.37 | 0.38 | 0.15 | 0.25 | 0.08 | 0.10 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

| Age Category | Lifetime (2011) | Lifetime (2012) | Past Year (2011) | Past Year (2012) | Past Month (2011) | Past Month (2012) |
|--------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.0038 | | 0.0069 | | 0.2255 | |
| 12 | 0.9463 | | 0.7087 | | 0.5900 | |
| 13 | 0.5479 | | 0.3624 | | 0.1512 | |
| 14 | 0.7254 | | 0.8167 | | 0.4307 | |
| 15 | 0.7656 | | 0.1040 | | 0.2438 | |
| 16 | 0.2915 | | 0.0528 | | 0.6592 | |
| 17 | 0.6673 | | 0.2134 | | 0.6525 | |
| 18 | 0.0712 | | 0.1429 | | 0.3974 | |
| 19 | 0.9282 | | 0.5043 | | 0.3492 | |
| 20 | 0.6691 | | 0.3187 | | 0.3051 | |
| 21 | 0.6425 | | 0.1746 | | 0.6551 | |
| 22 | 0.2919 | | 0.0833 | | 0.4166 | |
| 23 | 0.3773 | | 0.8109 | | 0.4667 | |
| 24 | 0.8067 | | 0.9766 | | 0.7021 | |
| 25 | 0.4390 | | 0.6817 | | 0.3818 | |
| 26-29 | 0.0835 | | 0.0842 | | 0.8740 | |
| 30-34 | 0.1434 | | 0.5717 | | 0.9616 | |
| 35-39 | 0.0106 | | 0.2947 | | 0.4986 | |
| 40-44 | 0.1335 | | 0.2670 | | 0.5307 | |
| 45-49 | 0.4139 | | 0.2114 | | 0.6400 | |
| 50-54 | 0.9738 | | 0.0324 | | 0.7103 | |
| 55-59 | 0.4417 | | 0.3872 | | 0.8360 | |
| 60-64 | 0.1220 | | 0.6400 | | 0.3975 | |
| 65 or Older | 0.6609 | | 0.2886 | | 0.4118 | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.