

Table 1.49C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	752	765	363	406	231	265
AGE						
12-17	64	64	53	53	35	37
18-25	145	151	110	111	69	74
26 or Older	740	740	338	380	215	247
GENDER						
Male	518	557	250	293	165	181
Female	496	509	242	277	152	173
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	702	711	336	358	207	240
White	634	662	307	329	186	221
Black or African American	208	218	106	143	52	70
American Indian or Alaska Native	51	60	35	39	25	22
Native Hawaiian or Other Pacific Islander	*	51	16	12	15	5
Asian	111	142	53	71	38	41
Two or More Races	107	88	33	33	25	17
Hispanic or Latino	258	313	135	186	96	101

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.49D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.29	0.29	0.14	0.16	0.09	0.10
AGE						
12-17	0.26	0.26	0.21	0.21	0.14	0.15
18-25	0.42	0.44	0.32	0.32	0.20	0.21
26 or Older	0.37	0.37	0.17	0.19	0.11	0.12
GENDER						
Male	0.42	0.44	0.20	0.23	0.13	0.14
Female	0.37	0.38	0.18	0.21	0.11	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.32	0.32	0.15	0.16	0.09	0.11
White	0.37	0.39	0.18	0.19	0.11	0.13
Black or African American	0.69	0.71	0.35	0.46	0.17	0.23
American Indian or Alaska Native	3.19	3.60	2.34	2.52	1.63	1.52
Native Hawaiian or Other Pacific Islander	*	4.70	1.75	1.20	1.65	0.48
Asian	0.87	1.05	0.42	0.54	0.30	0.33
Two or More Races	2.31	2.08	0.84	0.83	0.63	0.41
Hispanic or Latino	0.65	0.78	0.34	0.46	0.24	0.25

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.49P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.0121		0.0004		0.0593	
AGE						
12-17	0.6854		0.1606		0.9119	
18-25	0.2126		0.0320		0.2813	
26 or Older	0.0206		0.0009		0.0991	
GENDER						
Male	0.0541		0.0178		0.2199	
Female	0.0823		0.0061		0.1207	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2252		0.0041		0.0779	
White	0.4487		0.0812		0.1828	
Black or African American	0.3787		0.0048		0.0162	
American Indian or Alaska Native	0.4872		0.9189		0.5389	
Native Hawaiian or Other Pacific Islander	*		0.6622		0.1417	
Asian	0.0667		0.1767		0.4835	
Two or More Races	0.8160		0.8395		0.2906	
Hispanic or Latino	0.0004		0.0314		0.4722	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.50C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	64	64	53	53	35	37
GENDER						
Male	46	46	37	38	24	26
Female	46	46	38	38	24	27
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	54	53	45	45	30	31
White	50	51	42	43	24	28
Black or African American	28	26	21	21	13	15
American Indian or Alaska Native	6	5	6	3	4	3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	13	10	10	7	6	5
Two or More Races	14	13	11	12	8	8
Hispanic or Latino	35	37	28	27	16	19
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	34	34	28	27	17	18
Female, White, Not Hispanic	34	38	30	32	18	21
Male, Black, Not Hispanic	18	18	15	13	8	7
Female, Black, Not Hispanic	20	19	13	17	10	13
Male, Hispanic	25	28	19	23	11	16
Female, Hispanic	27	27	21	19	12	12

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.50D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.26	0.26	0.21	0.21	0.14	0.15
GENDER						
Male	0.36	0.36	0.29	0.29	0.18	0.20
Female	0.37	0.38	0.31	0.31	0.20	0.22
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.28	0.27	0.23	0.23	0.16	0.16
White	0.32	0.32	0.28	0.28	0.17	0.19
Black or African American	0.73	0.70	0.55	0.57	0.37	0.40
American Indian or Alaska Native	3.46	3.01	3.40	1.94	2.83	1.61
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.02	0.84	0.80	0.54	0.48	0.38
Two or More Races	1.81	1.66	1.52	1.53	1.07	1.02
Hispanic or Latino	0.65	0.67	0.51	0.50	0.30	0.35
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.45	0.45	0.37	0.36	0.23	0.25
Female, White, Not Hispanic	0.46	0.50	0.42	0.44	0.26	0.30
Male, Black, Not Hispanic	0.98	0.96	0.83	0.73	0.46	0.39
Female, Black, Not Hispanic	1.05	1.02	0.70	0.91	0.53	0.69
Male, Hispanic	0.85	0.91	0.67	0.77	0.39	0.57
Female, Hispanic	0.93	0.99	0.76	0.68	0.45	0.44

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.50P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.6854		0.1606		0.9119	
GENDER						
Male	0.9545		0.3475		0.8587	
Female	0.6066		0.2918		0.9822	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7606		0.3062		0.8840	
White	0.7406		0.2646		0.8947	
Black or African American	0.8305		0.5228		0.4228	
American Indian or Alaska Native	0.3860		0.0802		0.3801	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6021		0.4442		0.8115	
Two or More Races	0.8448		0.7072		0.4481	
Hispanic or Latino	0.7925		0.3215		0.9854	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.5559		0.2640		0.8878	
Female, White, Not Hispanic	0.9021		0.6318		0.9691	
Male, Black, Not Hispanic	0.7435		0.9811		0.9973	
Female, Black, Not Hispanic	0.9880		0.3931		0.3218	
Male, Hispanic	0.8899		0.9037		0.5508	
Female, Hispanic	0.6054		0.1770		0.4880	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.51C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	145	151	110	111	69	74
GENDER						
Male	104	105	79	84	50	52
Female	99	101	71	72	45	49
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	129	135	98	99	63	67
White	151	168	101	107	61	65
Black or African American	51	55	30	41	17	24
American Indian or Alaska Native	14	*	8	*	4	4
Native Hawaiian or Other Pacific Islander	*	*	*	5	*	4
Asian	31	31	18	23	11	18
Two or More Races	22	30	16	24	12	9
Hispanic or Latino	69	69	46	53	27	31
EDUCATION						
< High School	67	72	46	54	31	28
High School Graduate	99	109	68	78	41	44
Some College	121	125	78	86	46	51
College Graduate	68	66	37	43	23	22
CURRENT EMPLOYMENT						
Full-Time	112	114	70	74	41	48
Part-Time	91	103	61	71	40	39
Unemployed	59	58	44	43	30	32
Other ¹	79	92	55	68	34	31

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.51D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.42	0.44	0.32	0.32	0.20	0.21
GENDER						
Male	0.61	0.60	0.46	0.48	0.29	0.30
Female	0.58	0.59	0.42	0.42	0.26	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.49	0.36	0.36	0.23	0.24
White	0.56	0.57	0.43	0.43	0.28	0.30
Black or African American	0.92	0.99	0.59	0.77	0.36	0.46
American Indian or Alaska Native	4.43	*	3.36	*	1.78	1.41
Native Hawaiian or Other Pacific Islander	*	*	*	2.62	*	1.99
Asian	1.55	1.54	0.93	1.25	0.57	0.98
Two or More Races	2.74	2.87	2.12	2.75	1.63	1.20
Hispanic or Latino	0.98	0.97	0.65	0.74	0.38	0.44
EDUCATION						
< High School	0.94	1.10	0.73	0.91	0.52	0.51
High School Graduate	0.69	0.68	0.52	0.53	0.33	0.36
Some College	0.70	0.72	0.53	0.58	0.35	0.39
College Graduate	1.13	1.01	0.69	0.74	0.45	0.41
CURRENT EMPLOYMENT						
Full-Time	0.72	0.64	0.50	0.50	0.30	0.34
Part-Time	0.74	0.84	0.57	0.66	0.41	0.39
Unemployed	1.15	1.19	0.89	0.98	0.64	0.75
Other ¹	0.78	0.87	0.61	0.71	0.40	0.39

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.51P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.2126		0.0320		0.2813	
GENDER						
Male	0.7419		0.6552		0.6069	
Female	0.1265		0.0056		0.0264	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3530		0.0304		0.6362	
White	0.5667		0.2105		0.9079	
Black or African American	0.6163		0.1372		0.2578	
American Indian or Alaska Native	*		*		0.9721	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7905		0.1407		0.2771	
Two or More Races	0.0095		0.4242		0.5160	
Hispanic or Latino	0.3859		0.7007		0.1231	
EDUCATION						
< High School	0.1600		0.1312		0.9849	
High School Graduate	0.6104		0.4030		0.1363	
Some College	0.8209		0.5011		0.7179	
College Graduate	0.1174		0.0362		0.7667	
CURRENT EMPLOYMENT						
Full-Time	0.3082		0.2611		0.0228	
Part-Time	0.5637		0.5439		0.4182	
Unemployed	0.4071		0.0427		0.3431	
Other ¹	0.7979		0.3357		0.6560	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	740	740	338	380	215	247
GENDER						
Male	514	546	234	280	155	172
Female	484	485	225	260	141	162
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	687	688	309	333	192	225
White	743	772	293	310	178	210
Black or African American	223	243	101	147	48	67
American Indian or Alaska Native	48	50	33	24	23	22
Native Hawaiian or Other Pacific Islander	*	*	*	10	*	3
Asian	103	142	48	67	36	37
Two or More Races	103	82	25	21	18	12
Hispanic or Latino	245	301	126	174	91	94
EDUCATION						
< High School	241	313	121	168	88	89
High School Graduate	392	418	167	214	117	133
Some College	452	446	183	210	111	123
College Graduate	496	516	192	175	104	129
CURRENT EMPLOYMENT						
Full-Time	635	647	247	282	161	172
Part-Time	266	301	141	124	78	92
Unemployed	178	195	96	108	67	88
Other ¹	384	396	153	198	101	104

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.37	0.37	0.17	0.19	0.11	0.12
GENDER						
Male	0.54	0.57	0.25	0.29	0.16	0.18
Female	0.47	0.46	0.22	0.25	0.14	0.15
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.40	0.40	0.18	0.19	0.11	0.13
White	0.46	0.47	0.21	0.22	0.13	0.15
Black or African American	0.93	0.95	0.44	0.63	0.22	0.30
American Indian or Alaska Native	4.07	4.34	2.99	2.28	2.11	2.09
Native Hawaiian or Other Pacific Islander	*	*	*	1.37	*	0.46
Asian	1.09	1.35	0.51	0.66	0.38	0.38
Two or More Races	3.44	3.02	0.97	0.77	0.72	0.44
Hispanic or Latino	0.91	1.09	0.47	0.63	0.34	0.34
EDUCATION						
< High School	0.82	0.95	0.43	0.56	0.32	0.31
High School Graduate	0.62	0.64	0.28	0.36	0.20	0.23
Some College	0.78	0.74	0.36	0.40	0.22	0.24
College Graduate	0.69	0.68	0.30	0.28	0.16	0.20
CURRENT EMPLOYMENT						
Full-Time	0.50	0.52	0.23	0.26	0.15	0.16
Part-Time	1.01	1.09	0.59	0.51	0.33	0.39
Unemployed	1.69	1.74	1.02	1.08	0.71	0.88
Other ¹	0.58	0.57	0.24	0.31	0.16	0.16

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.0206		0.0009		0.0991	
GENDER						
Male	0.0608		0.0110		0.1507	
Female	0.1344		0.0244		0.3287	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2934		0.0106		0.0905	
White	0.4922		0.0883		0.1720	
Black or African American	0.4631		0.0154		0.0454	
American Indian or Alaska Native	0.3723		0.4846		0.6405	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0582		0.2639		0.6827	
Two or More Races	0.3923		0.8177		0.4853	
Hispanic or Latino	0.0003		0.0152		0.7723	
EDUCATION						
< High School	0.0807		0.2999		0.2778	
High School Graduate	0.1452		0.0732		0.4050	
Some College	0.7507		0.0146		0.5108	
College Graduate	0.1485		0.2307		0.0120	
CURRENT EMPLOYMENT						
Full-Time	0.0026		0.0011		0.1246	
Part-Time	0.8399		1.0000		0.3179	
Unemployed	0.5240		0.6975		0.2417	
Other ¹	0.8254		0.1221		0.6568	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.53C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	752	759	357	398	228	263
GENDER						
Male	518	553	246	290	164	180
Female	496	507	238	273	150	171
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	702	707	330	352	204	240
White	771	811	314	339	187	226
Black or African American	230	252	107	151	51	70
American Indian or Alaska Native	50	59	35	38	24	22
Native Hawaiian or Other Pacific Islander	*	*	16	11	15	5
Asian	109	142	52	71	37	41
Two or More Races	105	87	31	31	23	15
Hispanic or Latino	254	309	132	182	95	98
EDUCATION						
< High School	252	322	130	173	94	93
High School Graduate	407	429	182	228	124	142
Some College	478	471	202	226	124	130
College Graduate	506	529	196	184	106	132
CURRENT EMPLOYMENT						
Full-Time	646	659	254	290	165	180
Part-Time	278	318	151	147	87	101
Unemployed	188	202	106	115	73	92
Other ¹	393	404	164	209	108	110

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.53D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2011 and 2012

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.32	0.32	0.15	0.17	0.10	0.11
GENDER						
Male	0.46	0.49	0.22	0.26	0.15	0.16
Female	0.41	0.42	0.20	0.22	0.12	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.35	0.35	0.17	0.18	0.10	0.12
White	0.41	0.42	0.19	0.21	0.12	0.14
Black or African American	0.77	0.79	0.38	0.52	0.19	0.25
American Indian or Alaska Native	3.52	4.00	2.56	2.80	1.76	1.69
Native Hawaiian or Other Pacific Islander	*	*	1.86	1.24	1.76	0.54
Asian	0.95	1.15	0.46	0.59	0.33	0.36
Two or More Races	2.76	2.55	0.94	0.93	0.71	0.45
Hispanic or Latino	0.75	0.89	0.39	0.52	0.28	0.28
EDUCATION						
< High School	0.71	0.82	0.38	0.49	0.28	0.27
High School Graduate	0.53	0.55	0.26	0.31	0.18	0.20
Some College	0.65	0.62	0.31	0.34	0.20	0.20
College Graduate	0.65	0.64	0.28	0.27	0.15	0.19
CURRENT EMPLOYMENT						
Full-Time	0.45	0.46	0.21	0.24	0.14	0.15
Part-Time	0.74	0.82	0.45	0.43	0.27	0.30
Unemployed	1.19	1.25	0.75	0.80	0.52	0.65
Other ¹	0.53	0.52	0.23	0.29	0.15	0.15

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.53P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Demographic Characteristic	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOTAL	0.0117		0.0002		0.0592	
GENDER						
Male	0.0552		0.0110		0.2238	
Female	0.0767		0.0031		0.1156	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2276		0.0022		0.0803	
White	0.4516		0.0566		0.1890	
Black or African American	0.3932		0.0055		0.0215	
American Indian or Alaska Native	0.4245		0.9188		0.6249	
Native Hawaiian or Other Pacific Islander	*		0.5691		0.1383	
Asian	0.0585		0.1419		0.4592	
Two or More Races	0.8267		0.7082		0.3894	
Hispanic or Latino	0.0003		0.0166		0.4578	
EDUCATION						
< High School	0.0686		0.2339		0.2226	
High School Graduate	0.1191		0.0515		0.2138	
Some College	0.7839		0.0100		0.4371	
College Graduate	0.1012		0.1072		0.0112	
CURRENT EMPLOYMENT						
Full-Time	0.0012		0.0003		0.0297	
Part-Time	0.7249		0.8257		0.5586	
Unemployed	0.4080		0.3520		0.1803	
Other ¹	0.7491		0.1128		0.5360	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.