

**Table 1.78C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	752	765	363	406	231	265
<b>GEOGRAPHIC DIVISION</b>						
Northeast	321	297	139	151	91	96
New England	147	117	66	64	42	41
Middle Atlantic	284	272	122	137	81	86
Midwest	273	282	133	157	95	85
East North Central	223	242	115	135	83	72
West North Central	158	152	70	80	49	43
South	447	464	234	250	130	168
South Atlantic	354	346	183	179	95	113
East South Central	164	172	87	97	56	64
West South Central	231	250	114	138	76	102
West	437	431	202	222	135	152
Mountain	195	200	93	95	53	59
Pacific	391	380	181	203	124	141
<b>COUNTY TYPE</b>						
Large Metro	702	771	290	347	174	222
Small Metro	494	513	209	212	132	131
250K – 1 Mil. Pop.	414	440	164	171	108	114
< 250K Pop.	300	278	130	134	76	66
Nonmetro	334	304	146	161	93	73
Urbanized	252	220	117	119	69	50
Less Urbanized	205	196	82	105	59	48
Completely Rural	110	116	35	34	25	23

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.78D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.29	0.29	0.14	0.16	0.09	0.10
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.69	0.63	0.30	0.32	0.19	0.20
New England	1.20	0.94	0.54	0.52	0.34	0.33
Middle Atlantic	0.82	0.78	0.35	0.39	0.24	0.25
Midwest	0.49	0.50	0.24	0.28	0.17	0.15
East North Central	0.58	0.62	0.30	0.35	0.21	0.19
West North Central	0.93	0.89	0.41	0.47	0.29	0.25
South	0.47	0.48	0.25	0.26	0.14	0.17
South Atlantic	0.71	0.68	0.36	0.35	0.19	0.22
East South Central	1.07	1.12	0.57	0.63	0.37	0.41
West South Central	0.78	0.83	0.39	0.46	0.25	0.34
West	0.73	0.71	0.34	0.37	0.22	0.25
Mountain	1.07	1.09	0.51	0.51	0.29	0.32
Pacific	0.94	0.90	0.43	0.48	0.30	0.33
<b>COUNTY TYPE</b>						
Large Metro	0.44	0.44	0.21	0.23	0.13	0.16
Small Metro	0.46	0.49	0.24	0.25	0.16	0.16
250K – 1 Mil. Pop.	0.57	0.62	0.28	0.31	0.20	0.21
< 250K Pop.	0.81	0.78	0.47	0.45	0.28	0.24
Nonmetro	0.66	0.59	0.32	0.35	0.22	0.17
Urbanized	1.08	1.00	0.58	0.61	0.36	0.27
Less Urbanized	0.80	0.78	0.40	0.48	0.30	0.24
Completely Rural	1.96	1.87	0.78	0.66	0.57	0.46

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.78P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.0121		0.0004		0.0593	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.2175		0.2099		0.9917	
New England	0.2443		0.9120		0.9531	
Middle Atlantic	0.4297		0.1468		0.9699	
Midwest	0.4280		0.3149		0.4541	
East North Central	0.1326		0.2582		0.9711	
West North Central	0.3789		0.9261		0.1628	
South	0.5333		0.0319		0.0159	
South Atlantic	0.4569		0.3343		0.0623	
East South Central	0.4811		0.3104		0.3794	
West South Central	0.5507		0.0610		0.2017	
West	0.0185		0.0149		0.2010	
Mountain	0.5688		0.4178		0.3869	
Pacific	0.0178		0.0230		0.3020	
<b>COUNTY TYPE</b>						
Large Metro	0.0027		0.0001		0.0044	
Small Metro	0.3538		0.3190		0.7545	
250K – 1 Mil. Pop.	0.6360		0.2305		0.3690	
< 250K Pop.	0.3120		0.9716		0.3495	
Nonmetro	0.3952		0.9814		0.1355	
Urbanized	0.7288		0.3361		0.0947	
Less Urbanized	0.7066		0.2066		0.7612	
Completely Rural	0.1962		0.9019		0.6809	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.79C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	64	64	53	53	35	37
<b>GEOGRAPHIC DIVISION</b>						
Northeast	24	22	19	19	14	11
New England	10	11	9	9	6	5
Middle Atlantic	22	19	17	17	12	9
Midwest	26	24	21	21	15	13
East North Central	20	20	17	17	12	11
West North Central	16	13	13	12	8	8
South	39	43	33	34	21	26
South Atlantic	26	30	22	24	15	19
East South Central	15	14	13	12	7	8
West South Central	24	26	20	21	13	15
West	37	38	32	31	19	19
Mountain	18	17	13	16	9	9
Pacific	33	34	29	26	17	17
<b>COUNTY TYPE</b>						
Large Metro	56	56	43	47	26	32
Small Metro	41	36	35	28	21	18
250K – 1 Mil. Pop.	31	31	27	24	16	15
< 250K Pop.	26	22	20	16	14	10
Nonmetro	27	26	22	19	13	12
Urbanized	20	20	17	14	9	9
Less Urbanized	21	15	15	11	9	6
Completely Rural	8	8	6	7	3	5

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.79D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.26	0.26	0.21	0.21	0.14	0.15
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.57	0.52	0.45	0.46	0.32	0.26
New England	0.94	1.01	0.77	0.85	0.57	0.49
Middle Atlantic	0.69	0.61	0.54	0.54	0.38	0.30
Midwest	0.47	0.44	0.38	0.38	0.27	0.24
East North Central	0.53	0.54	0.44	0.46	0.32	0.28
West North Central	0.99	0.80	0.78	0.72	0.51	0.48
South	0.42	0.46	0.35	0.37	0.23	0.28
South Atlantic	0.56	0.65	0.48	0.52	0.33	0.41
East South Central	1.05	0.98	0.90	0.81	0.49	0.57
West South Central	0.77	0.81	0.63	0.65	0.40	0.49
West	0.62	0.63	0.53	0.52	0.32	0.33
Mountain	0.97	0.90	0.72	0.86	0.47	0.48
Pacific	0.79	0.82	0.70	0.64	0.42	0.42
<b>COUNTY TYPE</b>						
Large Metro	0.37	0.37	0.29	0.32	0.19	0.22
Small Metro	0.47	0.44	0.40	0.36	0.26	0.23
250K – 1 Mil. Pop.	0.54	0.53	0.47	0.44	0.29	0.27
< 250K Pop.	0.86	0.82	0.71	0.63	0.53	0.43
Nonmetro	0.60	0.59	0.52	0.46	0.33	0.29
Urbanized	0.94	0.91	0.82	0.70	0.50	0.46
Less Urbanized	0.94	0.82	0.73	0.64	0.48	0.38
Completely Rural	1.99	1.71	1.48	1.56	0.96	1.11

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.79P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.6854		0.1606		0.9119	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.1349		0.0638		0.4169	
New England	0.3442		0.6771		0.8103	
Middle Atlantic	0.2350		0.0585		0.2857	
Midwest	0.4375		0.3485		0.3140	
East North Central	0.7229		0.6371		0.5000	
West North Central	0.0873		0.3754		0.4550	
South	0.5111		0.7074		0.6176	
South Atlantic	0.1929		0.6223		0.3244	
East South Central	0.6405		0.5802		0.2589	
West South Central	0.9000		0.3739		0.8004	
West	0.9808		0.7475		0.4144	
Mountain	0.4778		0.1936		0.2524	
Pacific	0.6818		0.3100		0.7252	
<b>COUNTY TYPE</b>						
Large Metro	0.3686		0.2886		0.4594	
Small Metro	0.4117		0.6691		0.5438	
250K – 1 Mil. Pop.	0.6685		0.6952		0.7687	
< 250K Pop.	0.3781		0.8527		0.5270	
Nonmetro	0.6767		0.3921		0.7594	
Urbanized	0.4067		0.6409		0.7967	
Less Urbanized	0.1558		0.2976		0.7779	
Completely Rural	0.8179		0.6250		0.8867	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.80C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	145	151	110	111	69	74
<b>GEOGRAPHIC DIVISION</b>						
Northeast	53	62	39	47	26	38
New England	25	33	20	24	13	21
Middle Atlantic	47	53	34	40	22	32
Midwest	58	59	41	45	27	27
East North Central	49	46	35	34	23	24
West North Central	31	39	24	30	14	12
South	93	97	68	66	44	43
South Atlantic	65	70	50	52	32	33
East South Central	36	33	25	21	21	15
West South Central	53	55	38	33	23	22
West	81	85	60	66	39	39
Mountain	41	36	29	25	20	15
Pacific	70	77	52	61	33	36
<b>COUNTY TYPE</b>						
Large Metro	141	170	93	111	53	66
Small Metro	129	130	77	82	49	39
250K – 1 Mil. Pop.	93	87	59	56	36	29
< 250K Pop.	91	96	50	61	32	25
Nonmetro	72	57	49	35	25	21
Urbanized	61	43	41	23	18	13
Less Urbanized	40	38	27	24	17	16
Completely Rural	21	17	8	11	4	6

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.80D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.42	0.44	0.32	0.32	0.20	0.21
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.86	1.02	0.64	0.76	0.42	0.62
New England	1.53	2.03	1.26	1.48	0.80	1.28
Middle Atlantic	1.03	1.18	0.75	0.89	0.49	0.71
Midwest	0.78	0.81	0.57	0.61	0.36	0.36
East North Central	0.97	0.91	0.69	0.67	0.46	0.47
West North Central	1.35	1.72	1.05	1.31	0.62	0.53
South	0.73	0.76	0.54	0.52	0.35	0.33
South Atlantic	1.01	1.07	0.78	0.80	0.50	0.50
East South Central	1.79	1.65	1.23	1.05	1.02	0.75
West South Central	1.27	1.31	0.92	0.78	0.56	0.53
West	0.99	1.02	0.73	0.79	0.47	0.46
Mountain	1.68	1.43	1.16	1.00	0.81	0.61
Pacific	1.22	1.32	0.89	1.04	0.58	0.62
<b>COUNTY TYPE</b>						
Large Metro	0.59	0.65	0.44	0.47	0.27	0.33
Small Metro	0.77	0.74	0.55	0.56	0.38	0.32
250K – 1 Mil. Pop.	0.98	0.84	0.73	0.66	0.47	0.40
< 250K Pop.	1.16	1.42	0.84	1.07	0.63	0.54
Nonmetro	1.02	0.91	0.84	0.64	0.46	0.42
Urbanized	1.67	1.42	1.43	0.89	0.71	0.56
Less Urbanized	1.29	1.34	1.08	0.96	0.71	0.70
Completely Rural	3.53	2.78	1.61	2.18	0.89	1.25

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.80P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.2126		0.0320		0.2813	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0329		0.0466		0.0985	
New England	0.3436		0.7986		0.5861	
Middle Atlantic	0.0552		0.0275		0.1169	
Midwest	0.6730		0.7371		0.0927	
East North Central	0.6180		0.8447		0.4143	
West North Central	0.9801		0.7644		0.0459	
South	0.7856		0.1866		0.2067	
South Atlantic	0.7644		0.1295		0.1838	
East South Central	0.1787		0.3606		0.7497	
West South Central	0.2591		0.3111		0.4585	
West	0.9420		0.2175		0.6819	
Mountain	0.3722		0.1721		0.8921	
Pacific	0.5750		0.0422		0.6961	
<b>COUNTY TYPE</b>						
Large Metro	0.0038		0.0007		0.0672	
Small Metro	0.3673		0.9131		0.6304	
250K – 1 Mil. Pop.	0.1815		0.7971		0.7202	
< 250K Pop.	0.8565		0.6245		0.2139	
Nonmetro	0.1850		0.1059		0.7457	
Urbanized	0.1136		0.0250		0.2620	
Less Urbanized	0.7067		0.7283		0.7657	
Completely Rural	0.6881		0.2851		0.5462	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.81C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	740	740	338	380	215	247
<b>GEOGRAPHIC DIVISION</b>						
Northeast	316	288	129	137	85	87
New England	143	116	61	57	38	36
Middle Atlantic	281	263	114	124	76	79
Midwest	261	268	124	148	89	78
East North Central	211	231	108	128	77	66
West North Central	154	142	65	71	47	37
South	437	444	214	236	119	161
South Atlantic	347	331	169	169	85	108
East South Central	156	167	78	92	49	60
West South Central	221	236	106	132	69	98
West	434	423	187	209	128	143
Mountain	188	194	84	88	48	54
Pacific	390	374	168	190	119	132
<b>COUNTY TYPE</b>						
Large Metro	671	730	266	317	160	207
Small Metro	457	464	181	185	118	121
250K – 1 Mil. Pop.	392	408	144	152	102	105
< 250K Pop.	264	234	109	110	60	59
Nonmetro	307	281	131	150	89	65
Urbanized	236	201	106	111	67	45
Less Urbanized	179	180	71	96	53	41
Completely Rural	96	104	30	29	24	20

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.81D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.37	0.37	0.17	0.19	0.11	0.12
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.87	0.78	0.35	0.37	0.23	0.24
New England	1.49	1.19	0.64	0.59	0.40	0.37
Middle Atlantic	1.04	0.97	0.43	0.46	0.28	0.29
Midwest	0.61	0.62	0.29	0.34	0.21	0.18
East North Central	0.71	0.77	0.36	0.43	0.26	0.22
West North Central	1.18	1.07	0.50	0.54	0.36	0.28
South	0.60	0.60	0.29	0.32	0.16	0.22
South Atlantic	0.89	0.83	0.43	0.43	0.22	0.27
East South Central	1.32	1.40	0.66	0.77	0.41	0.51
West South Central	0.99	1.04	0.47	0.58	0.31	0.43
West	0.95	0.91	0.41	0.45	0.28	0.31
Mountain	1.35	1.38	0.61	0.63	0.35	0.38
Pacific	1.23	1.16	0.53	0.59	0.37	0.41
<b>COUNTY TYPE</b>						
Large Metro	0.56	0.56	0.25	0.28	0.15	0.19
Small Metro	0.59	0.61	0.28	0.30	0.19	0.20
250K – 1 Mil. Pop.	0.72	0.77	0.33	0.36	0.24	0.25
< 250K Pop.	1.04	0.97	0.55	0.53	0.31	0.30
Nonmetro	0.80	0.72	0.38	0.43	0.27	0.19
Urbanized	1.39	1.21	0.73	0.74	0.47	0.32
Less Urbanized	0.95	0.97	0.44	0.59	0.35	0.26
Completely Rural	2.18	2.14	0.85	0.73	0.69	0.51

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.81P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.0206		0.0009		0.0991	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.3473		0.3268		0.6216	
New England	0.3058		0.8692		0.7418	
Middle Atlantic	0.6014		0.2433		0.6964	
Midwest	0.4302		0.1857		0.9204	
East North Central	0.1560		0.1856		0.7332	
West North Central	0.4507		0.7008		0.4562	
South	0.6117		0.0512		0.0371	
South Atlantic	0.5691		0.6406		0.1774	
East South Central	0.7150		0.1311		0.2036	
West South Central	0.6989		0.0606		0.2703	
West	0.0191		0.0223		0.2660	
Mountain	0.4868		0.2722		0.5020	
Pacific	0.0227		0.0470		0.3519	
<b>COUNTY TYPE</b>						
Large Metro	0.0102		0.0014		0.0206	
Small Metro	0.3029		0.2481		0.5419	
250K – 1 Mil. Pop.	0.4913		0.1390		0.3746	
< 250K Pop.	0.3806		0.9620		0.7318	
Nonmetro	0.5799		0.5054		0.1613	
Urbanized	0.9707		0.8093		0.1764	
Less Urbanized	0.8081		0.1190		0.6622	
Completely Rural	0.1521		0.6383		0.5763	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.82C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	752	759	357	398	228	263
<b>GEOGRAPHIC DIVISION</b>						
Northeast	322	293	136	149	89	95
New England	147	114	66	64	42	41
Middle Atlantic	286	269	120	135	80	85
Midwest	270	280	131	155	94	84
East North Central	221	241	113	133	82	72
West North Central	154	149	69	78	48	41
South	444	458	228	244	128	166
South Atlantic	351	343	179	177	93	111
East South Central	163	171	85	95	56	63
West South Central	230	245	111	134	73	101
West	439	430	198	219	133	152
Mountain	192	198	90	93	52	58
Pacific	393	380	178	200	123	140
<b>COUNTY TYPE</b>						
Large Metro	695	761	285	338	170	218
Small Metro	487	505	201	209	130	130
250K – 1 Mil. Pop.	411	432	158	168	107	113
< 250K Pop.	290	272	124	130	74	64
Nonmetro	325	298	138	158	91	71
Urbanized	245	215	110	116	68	48
Less Urbanized	197	194	78	104	58	48
Completely Rural	107	112	34	31	24	21

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.82D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.32	0.32	0.15	0.17	0.10	0.11
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.76	0.68	0.32	0.35	0.21	0.22
New England	1.31	1.01	0.58	0.57	0.37	0.36
Middle Atlantic	0.91	0.85	0.38	0.43	0.25	0.27
Midwest	0.54	0.55	0.26	0.31	0.19	0.17
East North Central	0.63	0.69	0.32	0.38	0.24	0.21
West North Central	1.00	0.96	0.45	0.50	0.31	0.26
South	0.52	0.53	0.27	0.28	0.15	0.19
South Atlantic	0.77	0.74	0.39	0.38	0.20	0.24
East South Central	1.18	1.23	0.62	0.68	0.41	0.45
West South Central	0.87	0.91	0.42	0.50	0.28	0.37
West	0.81	0.79	0.37	0.40	0.25	0.28
Mountain	1.18	1.20	0.55	0.56	0.32	0.35
Pacific	1.05	1.00	0.47	0.53	0.33	0.37
<b>COUNTY TYPE</b>						
Large Metro	0.48	0.48	0.23	0.25	0.14	0.17
Small Metro	0.51	0.54	0.26	0.28	0.18	0.18
250K – 1 Mil. Pop.	0.62	0.68	0.30	0.34	0.21	0.23
< 250K Pop.	0.89	0.85	0.50	0.49	0.30	0.26
Nonmetro	0.72	0.64	0.34	0.38	0.24	0.18
Urbanized	1.19	1.08	0.63	0.66	0.41	0.29
Less Urbanized	0.88	0.86	0.42	0.53	0.33	0.26
Completely Rural	2.10	1.98	0.84	0.68	0.61	0.46

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.82P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.0117		0.0002		0.0592	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.1830		0.1368		0.9052	
New England	0.2153		0.9563		0.9208	
Middle Atlantic	0.3880		0.0922		0.8618	
Midwest	0.3936		0.2553		0.5531	
East North Central	0.1423		0.2308		0.9498	
West North Central	0.4622		0.8098		0.1908	
South	0.5795		0.0237		0.0173	
South Atlantic	0.5323		0.3600		0.0826	
East South Central	0.5023		0.2646		0.2912	
West South Central	0.5397		0.0366		0.2066	
West	0.0189		0.0112		0.2404	
Mountain	0.6130		0.5465		0.4874	
Pacific	0.0168		0.0132		0.3233	
<b>COUNTY TYPE</b>						
Large Metro	0.0020		0.0001		0.0055	
Small Metro	0.4246		0.2773		0.6792	
250K – 1 Mil. Pop.	0.6856		0.1970		0.3433	
< 250K Pop.	0.3803		0.9449		0.4038	
Nonmetro	0.3987		0.8793		0.1436	
Urbanized	0.6520		0.3526		0.0996	
Less Urbanized	0.7995		0.1483		0.7970	
Completely Rural	0.1955		0.8279		0.6543	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.