

Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	173,890	173,654	81,859	82,829	68,225	69,497
Cigarettes	161,799	161,057	67,126	67,802	56,819	57,525
Smokeless Tobacco	46,269	45,936	11,746	12,040	8,243	9,036
Cigars	88,505	88,596	24,980	25,647	12,865	13,417
Pipe Tobacco ²	34,532	34,604	--	--	2,137	2,530
ALCOHOL	211,747 ^a	214,013	170,422 ^a	173,573	133,385	135,502
Binge Alcohol Use ³	--	--	--	--	58,341	59,685
Heavy Alcohol Use ³	--	--	--	--	15,865	17,021

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	67.5	66.8	31.8	31.9	26.5	26.7
Cigarettes	62.8	61.9	26.1	26.1	22.1	22.1
Smokeless Tobacco	18.0	17.7	4.6	4.6	3.2	3.5
Cigars	34.4	34.1	9.7	9.9	5.0	5.2
Pipe Tobacco ²	13.4	13.3	--	--	0.8	1.0
ALCOHOL	82.2	82.3	66.2	66.7	51.8	52.1
Binge Alcohol Use ³	--	--	--	--	22.6	23.0
Heavy Alcohol Use ³	--	--	--	--	6.2	6.5

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	602	512	366 ^a	293	154	128
Cigarettes	444	383	257	222	108	93
Smokeless Tobacco	155	154	89	81	36	36
Cigars	222 ^b	136	142 ^b	81	49	31
Pipe Tobacco ²	45	56	--	--	16	24
ALCOHOL	922 ^a	794	608	546	196	179
Binge Alcohol Use ³	--	--	--	--	83	73
Heavy Alcohol Use ³	--	--	--	--	4 ^a	16

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	7.6 ^a	6.4	4.6 ^a	3.6	1.9	1.6
Cigarettes	5.6	4.8	3.3	2.8	1.4	1.2
Smokeless Tobacco	2.0	1.9	1.1	1.0	0.5	0.4
Cigars	2.8 ^b	1.7	1.8 ^b	1.0	0.6	0.4
Pipe Tobacco ²	0.6	0.7	--	--	0.2	0.3
ALCOHOL	11.7 ^a	9.9	7.7	6.8	2.5	2.2
Binge Alcohol Use ³	--	--	--	--	1.1	0.9
Heavy Alcohol Use ³	--	--	--	--	0.1 ^a	0.2

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	1,816	1,677	1,279 ^a	1,140	671 ^b	529
Cigarettes	1,468	1,339	966	859	506 ^b	389
Smokeless Tobacco	501	466	337	280	144	143
Cigars	772 ^a	645	538	463	214 ^b	141
Pipe Tobacco ²	176	165	--	--	53	41
ALCOHOL	2,842	2,658	2,227	2,116	959	936
Binge Alcohol Use ³	--	--	--	--	481	454
Heavy Alcohol Use ³	--	--	--	--	81 ^a	50

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	21.4	20.0	15.1 ^a	13.6	7.9 ^b	6.3
Cigarettes	17.3	16.0	11.4	10.2	6.0 ^b	4.6
Smokeless Tobacco	5.9	5.6	4.0	3.3	1.7	1.7
Cigars	9.1 ^a	7.7	6.3	5.5	2.5 ^b	1.7
Pipe Tobacco ²	2.1	2.0	--	--	0.6	0.5
ALCOHOL	33.5	31.7	26.3	25.2	11.3	11.1
Binge Alcohol Use ³	--	--	--	--	5.7	5.4
Heavy Alcohol Use ³	--	--	--	--	0.9 ^a	0.6

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	3,364	3,165	2,558	2,368	1,665 ^a	1,496
Cigarettes	2,861 ^a	2,616	2,064 ^a	1,868	1,325 ^a	1,152
Smokeless Tobacco	1,031	966	699	611	337	336
Cigars	1,681	1,608	1,259	1,139	595 ^b	473
Pipe Tobacco ²	353	399	--	--	112	100
ALCOHOL	4,847	4,615	4,105	3,893	2,171	2,102
Binge Alcohol Use ³	--	--	--	--	1,285	1,273
Heavy Alcohol Use ³	--	--	--	--	294	260

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	39.2	37.3	29.8	27.9	19.4 ^a	17.6
Cigarettes	33.3 ^a	30.8	24.0 ^a	22.0	15.4 ^a	13.6
Smokeless Tobacco	12.0	11.4	8.1	7.2	3.9	4.0
Cigars	19.6	19.0	14.7	13.4	6.9 ^b	5.6
Pipe Tobacco ²	4.1	4.7	--	--	1.3	1.2
ALCOHOL	56.4	54.4	47.8	45.9	25.3	24.8
Binge Alcohol Use ³	--	--	--	--	15.0	15.0
Heavy Alcohol Use ³	--	--	--	--	3.4	3.1

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	8,305	7,865	6,626	6,248	5,054 ^a	4,634
Cigarettes	7,268	6,878	5,552 ^a	5,131	4,272 ^b	3,745
Smokeless Tobacco	2,550 ^a	2,313	1,430 ^a	1,255	748	710
Cigars	4,702	4,557	3,313	3,188	1,683	1,582
Pipe Tobacco ²	1,189	1,182	--	--	291	276
ALCOHOL	10,059	9,849	8,998	8,817	6,328	6,077
Binge Alcohol Use ³	--	--	--	--	4,226	4,053
Heavy Alcohol Use ³	--	--	--	--	1,319	1,325

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	61.4 ^a	59.3	49.0	47.1	37.4 ^a	34.9
Cigarettes	53.7	51.8	41.0 ^a	38.7	31.6 ^b	28.2
Smokeless Tobacco	18.9	17.4	10.6	9.5	5.5	5.4
Cigars	34.8	34.3	24.5	24.0	12.4	11.9
Pipe Tobacco ²	8.8	8.9	--	--	2.2	2.1
ALCOHOL	74.4	74.2	66.5	66.4	46.8	45.8
Binge Alcohol Use ³	--	--	--	--	31.2	30.5
Heavy Alcohol Use ³	--	--	--	--	9.8	10.0

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	14,998	15,061	10,533	10,643	8,479	8,533
Cigarettes	13,661	13,714	8,964	9,063	7,216	7,263
Smokeless Tobacco	4,452	4,578	1,789	1,854	1,119	1,201
Cigars	8,838	8,933	4,256	4,332	2,041	2,134
Pipe Tobacco ²	2,141	2,239	--	--	344	349
ALCOHOL	18,842	19,345	17,432	17,961	14,486	14,750
Binge Alcohol Use ³	--	--	--	--	9,427	9,609
Heavy Alcohol Use ³	--	--	--	--	2,846	3,061

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	72.2	70.7	50.7	49.9	40.8	40.0
Cigarettes	65.7	64.3	43.1	42.5	34.7	34.1
Smokeless Tobacco	21.4	21.5	8.6	8.7	5.4	5.6
Cigars	42.5	41.9	20.5	20.3	9.8	10.0
Pipe Tobacco ²	10.3	10.5	--	--	1.7	1.6
ALCOHOL	90.7	90.8	83.9	84.3	69.7	69.2
Binge Alcohol Use ³	--	--	--	--	45.4	45.1
Heavy Alcohol Use ³	--	--	--	--	13.7	14.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	26,784	27,131	16,159	16,479	13,509	13,824
Cigarettes	25,161	25,468	13,669	14,110	11,495	12,033
Smokeless Tobacco	8,316	8,157	2,264	2,290	1,570	1,722
Cigars	15,773	15,869	5,207	5,396	2,753	2,709
Pipe Tobacco ²	2,816 ^a	3,370	--	--	284	402
ALCOHOL	32,779 ^a	33,463	28,831 ^a	29,722	23,230	23,792
Binge Alcohol Use ³	--	--	--	--	13,004	13,126
Heavy Alcohol Use ³	--	--	--	--	3,823	3,465

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	73.6	73.5	44.4	44.6	37.1	37.5
Cigarettes	69.1	69.0	37.5	38.2	31.6	32.6
Smokeless Tobacco	22.8	22.1	6.2	6.2	4.3	4.7
Cigars	43.3	43.0	14.3	14.6	7.6	7.3
Pipe Tobacco ²	7.7 ^a	9.1	--	--	0.8	1.1
ALCOHOL	90.0	90.7	79.2	80.5	63.8	64.5
Binge Alcohol Use ³	--	--	--	--	35.7	35.6
Heavy Alcohol Use ³	--	--	--	--	10.5	9.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	118,020	118,243	44,337	45,659	38,694	40,355
Cigarettes	110,935	110,657	35,655	36,549	31,897	32,849
Smokeless Tobacco	29,263	29,302	5,138	5,670	4,289	4,888
Cigars	56,517	56,847	10,265	11,047	5,530	6,346
Pipe Tobacco ²	27,811	27,193	--	--	1,036	1,338
ALCOHOL	141,457 ^a	143,288	108,222	110,518	86,014	87,666
Binge Alcohol Use ³	--	--	--	--	29,834	31,098
Heavy Alcohol Use ³	--	--	--	--	7,497 ^b	8,844

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	72.9	72.3	27.4	27.9	23.9	24.7
Cigarettes	68.5	67.6	22.0	22.3	19.7	20.1
Smokeless Tobacco	18.1	17.9	3.2	3.5	2.6	3.0
Cigars	34.9	34.7	6.3	6.8	3.4	3.9
Pipe Tobacco ²	17.2	16.6	--	--	0.6	0.8
ALCOHOL	87.4	87.6	66.8	67.5	53.1	53.6
Binge Alcohol Use ³	--	--	--	--	18.4	19.0
Heavy Alcohol Use ³	--	--	--	--	4.6 ^b	5.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	93,794	93,877	48,509	49,600	40,232	41,522
Cigarettes	84,626	84,479	35,937	36,941	30,331	31,021
Smokeless Tobacco	39,296	38,912	10,711	11,130	7,676 ^a	8,496
Cigars	64,738	65,531	19,392	19,909	10,179	10,756
Pipe Tobacco ²	29,886	30,116	--	--	1,728	2,038
ALCOHOL	106,455	107,153	86,725	88,046	70,736	71,187
Binge Alcohol Use ³	--	--	--	--	37,375	38,277
Heavy Alcohol Use ³	--	--	--	--	11,663	12,469

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	75.3	74.6	38.9	39.4	32.3	33.0
Cigarettes	67.9	67.1	28.8	29.3	24.3	24.6
Smokeless Tobacco	31.5	30.9	8.6	8.8	6.2	6.7
Cigars	51.9	52.1	15.6	15.8	8.2	8.5
Pipe Tobacco ²	24.0	23.9	--	--	1.4	1.6
ALCOHOL	85.4	85.1	69.6	69.9	56.8	56.5
Binge Alcohol Use ³	--	--	--	--	30.0	30.4
Heavy Alcohol Use ³	--	--	--	--	9.4	9.9

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	80,096	79,777	33,349	33,229	27,993	27,976
Cigarettes	77,173	76,578	31,189	30,861	26,488	26,504
Smokeless Tobacco	6,973	7,025	1,035	910	568	540
Cigars	23,767	23,065	5,588	5,738	2,686	2,661
Pipe Tobacco ²	4,645	4,488	--	--	410	492
ALCOHOL	105,292 ^a	106,860	83,697 ^a	85,526	62,649	64,315
Binge Alcohol Use ³	--	--	--	--	20,966	21,409
Heavy Alcohol Use ³	--	--	--	--	4,202	4,553

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2011 and 2012

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
TOBACCO PRODUCTS¹	60.2	59.5	25.1	24.8	21.1	20.9
Cigarettes	58.0	57.1	23.5	23.0	19.9	19.8
Smokeless Tobacco	5.2	5.2	0.8	0.7	0.4	0.4
Cigars	17.9	17.2	4.2	4.3	2.0	2.0
Pipe Tobacco ²	3.5	3.3	--	--	0.3	0.4
ALCOHOL	79.2	79.6	62.9	63.7	47.1	47.9
Binge Alcohol Use ³	--	--	--	--	15.8	16.0
Heavy Alcohol Use ³	--	--	--	--	3.2	3.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.