

**Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	926	939	900	933	846	882
Cigarettes	927	956	859	890	816	839
Smokeless Tobacco	752	705	326	325	278	298
Cigars	920	882	504	511	373	388
Pipe Tobacco <sup>2</sup>	698	679	--	--	144	184
<b>ALCOHOL</b>	684	675	984	955	1,006	1,016
Binge Alcohol Use <sup>3</sup>	--	--	--	--	734	797
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	423	448

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2011 and 2012**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.36	0.36	0.35	0.36	0.33	0.34
Cigarettes	0.36	0.37	0.33	0.34	0.32	0.32
Smokeless Tobacco	0.29	0.27	0.13	0.13	0.11	0.11
Cigars	0.36	0.34	0.20	0.20	0.14	0.15
Pipe Tobacco <sup>2</sup>	0.27	0.26	--	--	0.06	0.07
<b>ALCOHOL</b>	0.27	0.26	0.38	0.37	0.39	0.39
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.29	0.31
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.16	0.17

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1528		0.8801		0.6014	
Cigarettes	0.0847		0.9761		0.8854	
Smokeless Tobacco	0.4376		0.6922		0.0804	
Cigars	0.5603		0.5579		0.4289	
Pipe Tobacco <sup>2</sup>	0.7960		--		0.1109	
<b>ALCOHOL</b>	0.7993		0.2667		0.5527	
Binge Alcohol Use <sup>3</sup>	--		--		0.4605	
Heavy Alcohol Use <sup>3</sup>	--		--		0.0933	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2011 and 2012**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	35	31	27	24	19	15
Cigarettes	30	26	23	20	16	12
Smokeless Tobacco	17	17	13	12	10	7
Cigars	22	16	17	11	11	7
Pipe Tobacco <sup>2</sup>	10	10	--	--	6	7
<b>ALCOHOL</b>	44	42	36	36	20	20
Binge Alcohol Use <sup>3</sup>	--	--	--	--	13	13
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	2	5

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.43	0.37	0.33	0.29	0.23	0.18
Cigarettes	0.36	0.32	0.28	0.25	0.20	0.16
Smokeless Tobacco	0.22	0.21	0.17	0.15	0.12	0.09
Cigars	0.28	0.20	0.21	0.14	0.14	0.08
Pipe Tobacco <sup>2</sup>	0.12	0.12	--	--	0.08	0.08
<b>ALCOHOL</b>	0.51	0.49	0.43	0.42	0.25	0.24
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.16	0.16
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.02	0.07

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0285		0.0244		0.2373	
Cigarettes	0.0807		0.1830		0.4122	
Smokeless Tobacco	0.8478		0.5631		0.9110	
Cigars	0.0009		0.0014		0.1365	
Pipe Tobacco <sup>2</sup>	0.5077		--		0.4116	
<b>ALCOHOL</b>	0.0100		0.1257		0.4643	
Binge Alcohol Use <sup>3</sup>	--		--		0.5135	
Heavy Alcohol Use <sup>3</sup>	--		--		0.0457	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2011 and 2012**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	62	56	50	45	39	30
Cigarettes	57	50	45	40	34	25
Smokeless Tobacco	32	30	26	22	15	16
Cigars	40	35	34	30	22	15
Pipe Tobacco <sup>2</sup>	18	18	--	--	10	8
<b>ALCOHOL</b>	80	77	70	67	45	44
Binge Alcohol Use <sup>3</sup>	--	--	--	--	31	29
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	11	8

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2011 and 2012**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.66	0.58	0.56	0.50	0.45	0.35
Cigarettes	0.63	0.54	0.51	0.45	0.39	0.29
Smokeless Tobacco	0.36	0.35	0.30	0.27	0.18	0.19
Cigars	0.45	0.40	0.39	0.35	0.25	0.18
Pipe Tobacco <sup>2</sup>	0.21	0.21	--	--	0.11	0.09
<b>ALCOHOL</b>	0.74	0.72	0.69	0.67	0.50	0.47
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.35	0.33
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.13	0.10

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0905		0.0383		0.0037	
Cigarettes	0.0888		0.0750		0.0053	
Smokeless Tobacco	0.4616		0.1016		0.9806	
Cigars	0.0151		0.1111		0.0064	
Pipe Tobacco <sup>2</sup>	0.7102		--		0.3196	
<b>ALCOHOL</b>	0.0787		0.2587		0.8095	
Binge Alcohol Use <sup>3</sup>	--		--		0.5928	
Heavy Alcohol Use <sup>3</sup>	--		--		0.0343	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	80	83	70	70	57	54
Cigarettes	73	77	62	62	51	49
Smokeless Tobacco	43	43	36	32	23	24
Cigars	58	56	48	46	33	28
Pipe Tobacco <sup>2</sup>	25	32	--	--	14	14
<b>ALCOHOL</b>	102	102	93	92	64	67
Binge Alcohol Use <sup>3</sup>	--	--	--	--	46	51
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	21	23

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.75	0.73	0.69	0.68	0.58	0.56
Cigarettes	0.71	0.72	0.63	0.63	0.54	0.51
Smokeless Tobacco	0.47	0.47	0.40	0.36	0.27	0.28
Cigars	0.59	0.58	0.51	0.50	0.37	0.32
Pipe Tobacco <sup>2</sup>	0.28	0.36	--	--	0.16	0.16
<b>ALCOHOL</b>	0.77	0.78	0.79	0.77	0.63	0.64
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.50	0.52
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.24	0.27

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0845		0.0552		0.0319	
Cigarettes	0.0147		0.0228		0.0138	
Smokeless Tobacco	0.3519		0.0833		0.9007	
Cigars	0.4716		0.0866		0.0062	
Pipe Tobacco <sup>2</sup>	0.1849		--		0.5519	
<b>ALCOHOL</b>	0.0684		0.0916		0.5997	
Binge Alcohol Use <sup>3</sup>	--		--		0.9458	
Heavy Alcohol Use <sup>3</sup>	--		--		0.3230	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	219	220	183	187	145	152
Cigarettes	193	202	157	162	128	135
Smokeless Tobacco	86	88	61	64	44	44
Cigars	153	151	117	122	73	75
Pipe Tobacco <sup>2</sup>	60	66	--	--	31	24
<b>ALCOHOL</b>	274	277	262	270	218	222
Binge Alcohol Use <sup>3</sup>	--	--	--	--	168	176
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	92	85

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.76	0.80	0.78	0.78	0.76	0.75
Cigarettes	0.79	0.83	0.78	0.78	0.72	0.73
Smokeless Tobacco	0.58	0.59	0.44	0.45	0.32	0.32
Cigars	0.74	0.73	0.64	0.66	0.47	0.45
Pipe Tobacco <sup>2</sup>	0.42	0.41	--	--	0.23	0.17
<b>ALCOHOL</b>	0.71	0.73	0.83	0.81	0.91	0.90
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.85	0.87
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.58	0.54

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0472		0.0833		0.0197	
Cigarettes	0.0855		0.0314		0.0008	
Smokeless Tobacco	0.0904		0.0693		0.6929	
Cigars	0.6670		0.6032		0.4393	
Pipe Tobacco <sup>2</sup>	0.8473		--		0.7990	
<b>ALCOHOL</b>	0.8604		0.9265		0.3900	
Binge Alcohol Use <sup>3</sup>	--		--		0.5195	
Heavy Alcohol Use <sup>3</sup>	--		--		0.7256	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	285	266	219	212	184	186
Cigarettes	258	244	193	188	168	165
Smokeless Tobacco	121	123	76	70	59	58
Cigars	202	197	130	122	86	79
Pipe Tobacco <sup>2</sup>	93	90	--	--	36	31
<b>ALCOHOL</b>	350	313	343	301	304	269
Binge Alcohol Use <sup>3</sup>	--	--	--	--	225	208
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	100	107

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.55	0.54	0.62	0.60	0.59	0.60
Cigarettes	0.60	0.58	0.61	0.59	0.57	0.58
Smokeless Tobacco	0.46	0.46	0.33	0.31	0.26	0.26
Cigars	0.61	0.60	0.49	0.45	0.36	0.33
Pipe Tobacco <sup>2</sup>	0.39	0.37	--	--	0.17	0.14
<b>ALCOHOL</b>	0.38	0.35	0.50	0.44	0.59	0.56
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.66	0.60
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.41	0.42

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0505		0.3663		0.3513	
Cigarettes	0.0978		0.4643		0.4129	
Smokeless Tobacco	0.9390		0.8381		0.5018	
Cigars	0.4623		0.8148		0.7110	
Pipe Tobacco <sup>2</sup>	0.7105		--		0.9350	
<b>ALCOHOL</b>	0.9033		0.5936		0.5046	
Binge Alcohol Use <sup>3</sup>	--		--		0.7286	
Heavy Alcohol Use <sup>3</sup>	--		--		0.2532	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	283	289	316	300	298	293
Cigarettes	298	293	305	288	286	278
Smokeless Tobacco	259	261	131	143	108	129
Cigars	314	309	219	222	172	168
Pipe Tobacco <sup>2</sup>	162	180	--	--	49	60
<b>ALCOHOL</b>	195	195	263	256	304	312
Binge Alcohol Use <sup>3</sup>	--	--	--	--	298	313
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	200	188

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.78	0.78	0.87	0.81	0.82	0.79
Cigarettes	0.82	0.79	0.84	0.78	0.78	0.75
Smokeless Tobacco	0.71	0.71	0.36	0.39	0.30	0.35
Cigars	0.86	0.84	0.60	0.60	0.47	0.45
Pipe Tobacco <sup>2</sup>	0.44	0.49	--	--	0.13	0.16
<b>ALCOHOL</b>	0.53	0.53	0.72	0.69	0.84	0.85
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.82	0.85
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.55	0.51

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.9556		0.8241		0.7639	
Cigarettes	0.9256		0.5489		0.3522	
Smokeless Tobacco	0.4656		0.9811		0.4481	
Cigars	0.7805		0.7135		0.7383	
Pipe Tobacco <sup>2</sup>	0.0339		--		0.1447	
<b>ALCOHOL</b>	0.4055		0.1853		0.5954	
Binge Alcohol Use <sup>3</sup>	--		--		0.8963	
Heavy Alcohol Use <sup>3</sup>	--		--		0.1394	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	819	821	756	799	716	765
Cigarettes	825	852	709	755	685	721
Smokeless Tobacco	656	638	266	278	241	261
Cigars	811	788	396	424	302	324
Pipe Tobacco <sup>2</sup>	661	644	--	--	126	165
<b>ALCOHOL</b>	609	610	901	874	921	899
Binge Alcohol Use <sup>3</sup>	--	--	--	--	604	664
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	324	374

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2011 and 2012**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.51	0.50	0.47	0.49	0.44	0.47
Cigarettes	0.51	0.52	0.44	0.46	0.42	0.44
Smokeless Tobacco	0.41	0.39	0.16	0.17	0.15	0.16
Cigars	0.50	0.48	0.24	0.26	0.19	0.20
Pipe Tobacco <sup>2</sup>	0.41	0.39	--	--	0.08	0.10
<b>ALCOHOL</b>	0.38	0.37	0.56	0.53	0.57	0.55
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.37	0.41
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.20	0.23

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.3910		0.4307		0.2236	
Cigarettes	0.2231		0.6101		0.5304	
Smokeless Tobacco	0.7627		0.2054		0.1136	
Cigars	0.8152		0.2616		0.0940	
Pipe Tobacco <sup>2</sup>	0.3394		--		0.1563	
<b>ALCOHOL</b>	0.6944		0.3587		0.5631	
Binge Alcohol Use <sup>3</sup>	--		--		0.2853	
Heavy Alcohol Use <sup>3</sup>	--		--		0.0093	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	558	536	605	655	581	618
Cigarettes	588	577	560	598	537	556
Smokeless Tobacco	652	609	311	307	264	281
Cigars	684	667	436	451	336	351
Pipe Tobacco <sup>2</sup>	623	615	--	--	135	170
<b>ALCOHOL</b>	408	380	630	605	674	675
Binge Alcohol Use <sup>3</sup>	--	--	--	--	567	625
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	355	382

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.45	0.43	0.49	0.52	0.47	0.49
Cigarettes	0.47	0.46	0.45	0.47	0.43	0.44
Smokeless Tobacco	0.52	0.48	0.25	0.24	0.21	0.22
Cigars	0.55	0.53	0.35	0.36	0.27	0.28
Pipe Tobacco <sup>2</sup>	0.50	0.49	--	--	0.11	0.14
<b>ALCOHOL</b>	0.33	0.30	0.51	0.48	0.54	0.54
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.45	0.50
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.28	0.30

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.2679		0.5065		0.3000	
Cigarettes	0.2262		0.4330		0.6251	
Smokeless Tobacco	0.3587		0.4787		0.0551	
Cigars	0.8925		0.6159		0.3368	
Pipe Tobacco <sup>2</sup>	0.9330		--		0.1824	
<b>ALCOHOL</b>	0.5017		0.6235		0.7798	
Binge Alcohol Use <sup>3</sup>	--		--		0.5309	
Heavy Alcohol Use <sup>3</sup>	--		--		0.1842	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Substance</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	660	683	564	578	542	554
Cigarettes	664	688	556	561	532	539
Smokeless Tobacco	281	296	97	109	82	103
Cigars	479	472	197	211	142	146
Pipe Tobacco <sup>2</sup>	234	222	--	--	46	61
<b>ALCOHOL</b>	530	516	690	660	684	686
Binge Alcohol Use <sup>3</sup>	--	--	--	--	441	427
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	207	203

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2011 and 2012**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.50	0.51	0.42	0.43	0.41	0.41
Cigarettes	0.50	0.51	0.42	0.42	0.40	0.40
Smokeless Tobacco	0.21	0.22	0.07	0.08	0.06	0.08
Cigars	0.36	0.35	0.15	0.16	0.11	0.11
Pipe Tobacco <sup>2</sup>	0.18	0.17	--	--	0.03	0.05
<b>ALCOHOL</b>	0.40	0.38	0.52	0.49	0.51	0.51
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.33	0.32
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.16	0.15

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

Substance	Lifetime (2011)	Lifetime (2012)	Past Year (2011)	Past Year (2012)	Past Month (2011)	Past Month (2012)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.2801		0.5989		0.7270	
Cigarettes	0.1796		0.4331		0.7659	
Smokeless Tobacco	0.9787		0.3693		0.8057	
Cigars	0.1749		0.7285		0.8054	
Pipe Tobacco <sup>2</sup>	0.5430		--		0.2995	
<b>ALCOHOL</b>	0.3915		0.2450		0.2373	
Binge Alcohol Use <sup>3</sup>	--		--		0.6774	
Heavy Alcohol Use <sup>3</sup>	--		--		0.2658	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.