

**Table 2.57C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:  
Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	752	705	326	325	278	298
<b>GEOGRAPHIC DIVISION</b>						
Northeast	244	235	93	91	80	77
New England	135	114	47	28	38	23
Middle Atlantic	202	207	81	87	70	74
Midwest	279	308	165	156	150	151
East North Central	221	243	139	129	127	122
West North Central	159	191	92	86	82	84
South	495	460	208	228	175	205
South Atlantic	372	344	149	146	128	133
East South Central	179	177	93	104	80	97
West South Central	245	260	118	137	94	123
West	404	361	143	152	124	139
Mountain	191	185	77	93	72	88
Pacific	353	321	120	120	101	106
<b>COUNTY TYPE</b>						
Large Metro	638	611	214	223	174	195
Small Metro	515	473	216	189	190	168
250K – 1 Mil. Pop.	402	363	168	138	146	125
< 250K Pop.	315	316	129	129	118	114
Nonmetro	367	423	152	190	126	178
Urbanized	219	237	102	107	65	99
Less Urbanized	286	317	105	139	99	128
Completely Rural	138	175	52	87	50	83

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.57D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:  
Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.29	0.27	0.13	0.13	0.11	0.11
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.52	0.50	0.20	0.19	0.17	0.16
New England	1.09	0.91	0.38	0.23	0.31	0.19
Middle Atlantic	0.59	0.59	0.23	0.25	0.20	0.21
Midwest	0.50	0.55	0.30	0.28	0.27	0.27
East North Central	0.57	0.63	0.36	0.33	0.33	0.32
West North Central	0.94	1.12	0.54	0.50	0.48	0.49
South	0.52	0.48	0.22	0.24	0.18	0.21
South Atlantic	0.74	0.68	0.30	0.29	0.26	0.26
East South Central	1.17	1.15	0.61	0.67	0.52	0.63
West South Central	0.82	0.86	0.40	0.46	0.32	0.41
West	0.67	0.60	0.24	0.25	0.21	0.23
Mountain	1.05	1.00	0.42	0.50	0.39	0.48
Pacific	0.85	0.76	0.29	0.28	0.24	0.25
<b>COUNTY TYPE</b>						
Large Metro	0.40	0.38	0.15	0.15	0.12	0.14
Small Metro	0.51	0.45	0.25	0.22	0.22	0.20
250K – 1 Mil. Pop.	0.61	0.54	0.30	0.25	0.26	0.23
< 250K Pop.	0.86	0.85	0.43	0.45	0.41	0.41
Nonmetro	0.68	0.75	0.34	0.41	0.30	0.39
Urbanized	1.07	1.00	0.54	0.53	0.39	0.50
Less Urbanized	0.99	1.19	0.51	0.65	0.48	0.59
Completely Rural	2.23	2.15	1.00	1.45	0.99	1.42

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.57P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:  
P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.4376		0.6922		0.0804	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.9344		0.8569		0.5774	
New England	0.9275		0.0736		0.2172	
Middle Atlantic	0.9664		0.5109		0.2474	
Midwest	0.3553		0.5626		0.7537	
East North Central	0.3897		0.4946		0.9406	
West North Central	0.6954		0.9861		0.6588	
South	0.2435		0.4493		0.0957	
South Atlantic	0.5507		0.8396		0.8868	
East South Central	0.6321		0.5000		0.2432	
West South Central	0.3256		0.2892		0.0656	
West	0.5002		0.6446		0.4356	
Mountain	0.6452		0.3638		0.3053	
Pacific	0.3096		0.9587		0.8220	
<b>COUNTY TYPE</b>						
Large Metro	0.1031		0.7369		0.5103	
Small Metro	0.4616		0.3756		0.7872	
250K – 1 Mil. Pop.	0.3067		0.1861		0.4808	
< 250K Pop.	0.8856		0.8092		0.6711	
Nonmetro	0.0525		0.0259		0.0035	
Urbanized	0.3560		0.1436		0.0017	
Less Urbanized	0.1901		0.2076		0.2138	
Completely Rural	0.1050		0.1910		0.2962	

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.58C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	56	53	46	42	30	31
<b>GEOGRAPHIC DIVISION</b>						
Northeast	24	16	20	14	14	9
New England	9	7	7	6	4	4
Middle Atlantic	23	15	18	13	14	9
Midwest	25	21	21	17	15	12
East North Central	19	16	16	12	12	9
West North Central	16	13	13	11	8	8
South	36	39	30	30	18	24
South Atlantic	27	27	22	21	13	17
East South Central	15	17	12	13	9	11
West South Central	19	22	16	17	10	13
West	24	24	20	18	12	12
Mountain	11	11	9	9	6	7
Pacific	21	21	18	15	11	9
<b>COUNTY TYPE</b>						
Large Metro	38	38	30	28	19	19
Small Metro	34	32	28	24	19	17
250K – 1 Mil. Pop.	25	24	22	18	12	14
< 250K Pop.	21	21	18	15	15	11
Nonmetro	32	30	23	22	15	18
Urbanized	18	16	14	12	8	9
Less Urbanized	25	24	17	19	12	15
Completely Rural	11	12	9	8	6	6

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.58D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.22	0.21	0.19	0.17	0.12	0.12
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.57	0.38	0.46	0.33	0.34	0.22
New England	0.81	0.63	0.65	0.54	0.38	0.34
Middle Atlantic	0.72	0.47	0.58	0.40	0.44	0.28
Midwest	0.45	0.39	0.38	0.31	0.27	0.22
East North Central	0.50	0.43	0.43	0.33	0.32	0.24
West North Central	0.95	0.81	0.78	0.67	0.52	0.46
South	0.39	0.41	0.32	0.32	0.20	0.25
South Atlantic	0.58	0.58	0.47	0.46	0.28	0.37
East South Central	1.00	1.16	0.84	0.89	0.59	0.73
West South Central	0.60	0.70	0.50	0.53	0.31	0.40
West	0.39	0.40	0.33	0.30	0.20	0.20
Mountain	0.59	0.60	0.51	0.50	0.31	0.40
Pacific	0.50	0.52	0.42	0.37	0.26	0.22
<b>COUNTY TYPE</b>						
Large Metro	0.27	0.26	0.22	0.20	0.14	0.14
Small Metro	0.41	0.40	0.35	0.31	0.24	0.22
250K – 1 Mil. Pop.	0.44	0.43	0.40	0.34	0.22	0.26
< 250K Pop.	0.85	0.80	0.74	0.61	0.61	0.46
Nonmetro	0.73	0.69	0.56	0.55	0.39	0.46
Urbanized	0.92	0.76	0.73	0.61	0.46	0.47
Less Urbanized	1.13	1.26	0.87	1.05	0.65	0.88
Completely Rural	2.52	2.21	2.24	1.70	1.45	1.33

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.58P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.1889		0.0126		0.9810	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0070		0.0070		0.1875	
New England	0.0754		0.1103		0.5457	
Middle Atlantic	0.0296		0.0239		0.2363	
Midwest	0.3140		0.0826		0.9761	
East North Central	0.2524		0.2133		0.9601	
West North Central	0.8439		0.2007		0.9786	
South	0.9775		0.6512		0.4464	
South Atlantic	0.7495		0.8180		0.3969	
East South Central	0.9841		0.5434		0.8522	
West South Central	0.5790		0.9803		0.6236	
West	0.7407		0.3518		0.8767	
Mountain	0.3131		0.4721		0.2225	
Pacific	0.3545		0.5073		0.2582	
<b>COUNTY TYPE</b>						
Large Metro	0.0630		0.0366		0.7551	
Small Metro	0.7291		0.0739		0.8602	
250K – 1 Mil. Pop.	0.4445		0.1215		0.2875	
< 250K Pop.	0.6965		0.3584		0.4296	
Nonmetro	0.9708		0.7407		0.8648	
Urbanized	0.5863		0.4932		0.9951	
Less Urbanized	0.7948		0.8500		0.7074	
Completely Rural	0.4906		0.9724		0.7659	

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.59C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	127	132	93	87	70	70
<b>GEOGRAPHIC DIVISION</b>						
Northeast	50	54	32	34	22	22
New England	26	24	18	18	12	11
Middle Atlantic	43	48	27	29	19	20
Midwest	54	54	38	40	31	36
East North Central	45	41	32	29	25	26
West North Central	29	34	23	28	19	26
South	80	78	60	54	47	46
South Atlantic	58	59	45	38	35	32
East South Central	31	28	21	23	17	21
West South Central	45	46	33	33	26	24
West	67	76	45	45	31	31
Mountain	32	34	21	23	15	18
Pacific	60	65	41	39	27	26
<b>COUNTY TYPE</b>						
Large Metro	105	115	70	64	50	47
Small Metro	105	109	62	64	44	46
250K – 1 Mil. Pop.	79	66	45	43	34	34
< 250K Pop.	70	86	42	45	28	31
Nonmetro	64	65	45	44	34	37
Urbanized	43	43	32	28	21	23
Less Urbanized	45	46	30	32	24	28
Completely Rural	22	22	15	15	13	11

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.59D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.37	0.38	0.27	0.25	0.20	0.20
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.82	0.87	0.53	0.55	0.36	0.36
New England	1.61	1.48	1.13	1.13	0.72	0.65
Middle Atlantic	0.96	1.06	0.59	0.63	0.42	0.44
Midwest	0.73	0.73	0.52	0.54	0.42	0.49
East North Central	0.88	0.80	0.64	0.56	0.50	0.51
West North Central	1.29	1.51	1.00	1.23	0.82	1.13
South	0.64	0.61	0.48	0.43	0.37	0.36
South Atlantic	0.90	0.90	0.70	0.58	0.54	0.50
East South Central	1.53	1.40	1.05	1.13	0.84	1.01
West South Central	1.09	1.08	0.80	0.79	0.62	0.58
West	0.82	0.91	0.55	0.54	0.38	0.38
Mountain	1.29	1.37	0.85	0.92	0.62	0.74
Pacific	1.04	1.11	0.70	0.67	0.47	0.44
<b>COUNTY TYPE</b>						
Large Metro	0.49	0.53	0.36	0.32	0.26	0.24
Small Metro	0.69	0.63	0.48	0.46	0.37	0.38
250K – 1 Mil. Pop.	0.86	0.73	0.57	0.51	0.46	0.46
< 250K Pop.	1.16	1.11	0.86	0.87	0.63	0.69
Nonmetro	0.96	0.97	0.76	0.76	0.64	0.65
Urbanized	1.42	1.46	1.15	1.03	0.86	0.87
Less Urbanized	1.52	1.44	1.14	1.24	1.00	1.08
Completely Rural	2.92	2.86	2.29	2.76	2.23	2.16

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.



**Table 2.59P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.3421		0.2695		0.7737	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.7023		0.4711		0.2289	
New England	0.4208		0.8682		1.0000	
Middle Atlantic	0.3728		0.3133		0.1618	
Midwest	0.3083		0.3262		0.7154	
East North Central	0.3833		0.5273		0.6363	
West North Central	0.5775		0.4703		0.9771	
South	0.9345		0.7591		0.7553	
South Atlantic	0.4645		0.7374		0.8466	
East South Central	0.2952		0.7686		0.7005	
West South Central	0.9022		0.9234		0.5827	
West	0.1785		0.1428		0.3199	
Mountain	0.8969		1.0000		0.8518	
Pacific	0.1060		0.0962		0.1857	
<b>COUNTY TYPE</b>						
Large Metro	0.4733		0.0342		0.5787	
Small Metro	0.4274		0.4578		0.3430	
250K – 1 Mil. Pop.	0.0304		0.7807		0.7891	
< 250K Pop.	0.1881		0.4378		0.2579	
Nonmetro	0.6952		0.8293		0.6525	
Urbanized	0.8093		0.7820		0.5398	
Less Urbanized	0.9660		0.6835		0.4887	
Completely Rural	0.3699		0.4130		0.1713	

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.60C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:  
Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	726	692	305	309	266	286
<b>GEOGRAPHIC DIVISION</b>						
Northeast	235	223	84	82	76	72
New England	129	114	44	26	36	20
Middle Atlantic	196	193	71	78	66	69
Midwest	273	293	157	144	146	141
East North Central	217	233	135	122	123	117
West North Central	158	185	83	77	79	75
South	470	452	191	219	164	197
South Atlantic	352	346	129	139	117	127
East South Central	176	171	91	102	77	95
West South Central	231	250	114	133	90	120
West	396	352	133	144	120	134
Mountain	187	177	75	89	71	84
Pacific	346	313	109	113	97	102
<b>COUNTY TYPE</b>						
Large Metro	604	570	189	204	161	183
Small Metro	483	440	197	171	179	158
250K – 1 Mil. Pop.	385	342	158	124	140	114
< 250K Pop.	293	288	114	117	109	108
Nonmetro	337	398	139	179	119	169
Urbanized	203	223	91	96	59	92
Less Urbanized	260	294	95	132	94	122
Completely Rural	122	160	45	81	44	78

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.60D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:  
Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.37	0.34	0.15	0.15	0.13	0.14
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.65	0.61	0.23	0.22	0.21	0.19
New England	1.34	1.18	0.45	0.27	0.38	0.21
Middle Atlantic	0.73	0.71	0.27	0.29	0.25	0.25
Midwest	0.64	0.68	0.37	0.33	0.34	0.33
East North Central	0.73	0.78	0.45	0.41	0.41	0.39
West North Central	1.21	1.40	0.63	0.58	0.60	0.57
South	0.64	0.61	0.26	0.30	0.22	0.27
South Atlantic	0.90	0.87	0.33	0.35	0.30	0.32
East South Central	1.49	1.44	0.77	0.85	0.65	0.80
West South Central	1.03	1.10	0.51	0.59	0.40	0.53
West	0.87	0.76	0.29	0.31	0.26	0.29
Mountain	1.35	1.26	0.54	0.63	0.51	0.60
Pacific	1.09	0.97	0.34	0.35	0.30	0.32
<b>COUNTY TYPE</b>						
Large Metro	0.50	0.47	0.18	0.19	0.15	0.17
Small Metro	0.64	0.58	0.31	0.27	0.28	0.25
250K – 1 Mil. Pop.	0.77	0.68	0.37	0.29	0.33	0.27
< 250K Pop.	1.11	1.10	0.53	0.56	0.52	0.52
Nonmetro	0.83	0.93	0.41	0.51	0.37	0.48
Urbanized	1.32	1.26	0.64	0.64	0.46	0.61
Less Urbanized	1.23	1.48	0.60	0.81	0.58	0.75
Completely Rural	2.77	2.63	1.23	1.75	1.20	1.68

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.60P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:  
P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.5784		0.2728		0.0778	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.9473		0.9168		0.6379	
New England	0.8324		0.1249		0.2075	
Middle Atlantic	0.9565		0.3674		0.2809	
Midwest	0.2303		0.8836		0.8133	
East North Central	0.2642		0.6735		0.9818	
West North Central	0.6151		0.6969		0.6347	
South	0.2146		0.3428		0.1171	
South Atlantic	0.4654		0.9275		0.9186	
East South Central	0.7609		0.4046		0.2659	
West South Central	0.2880		0.2873		0.0915	
West	0.6329		0.2726		0.2804	
Mountain	0.6160		0.3004		0.3623	
Pacific	0.4138		0.5304		0.4914	
<b>COUNTY TYPE</b>						
Large Metro	0.1506		0.5724		0.3811	
Small Metro	0.5279		0.3885		0.5797	
250K – 1 Mil. Pop.	0.5291		0.2122		0.3584	
< 250K Pop.	0.8119		0.8603		0.8160	
Nonmetro	0.0592		0.0205		0.0038	
Urbanized	0.3260		0.0851		0.0013	
Less Urbanized	0.2089		0.2680		0.3096	
Completely Rural	0.1459		0.1250		0.1638	

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.61C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:  
Standard Errors of Numbers in Thousands, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	746	707	319	324	275	298
<b>GEOGRAPHIC DIVISION</b>						
Northeast	242	234	90	89	78	76
New England	134	113	47	27	38	23
Middle Atlantic	201	206	77	85	68	73
Midwest	274	303	162	154	148	149
East North Central	219	239	138	128	126	122
West North Central	156	190	88	84	80	82
South	491	458	203	229	173	205
South Atlantic	369	344	143	147	126	133
East South Central	180	174	92	105	80	98
West South Central	240	257	118	136	93	123
West	404	363	142	151	124	138
Mountain	192	182	78	93	72	88
Pacific	353	324	119	119	101	105
<b>COUNTY TYPE</b>						
Large Metro	629	607	208	218	171	193
Small Metro	510	466	211	186	187	167
250K – 1 Mil. Pop.	397	358	164	136	144	124
< 250K Pop.	312	311	126	127	116	114
Nonmetro	356	416	148	187	125	176
Urbanized	214	233	98	104	64	98
Less Urbanized	276	309	101	137	97	126
Completely Rural	133	170	51	86	49	82

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.61D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:  
Standard Errors of Percentages, 2011 and 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.32	0.30	0.14	0.14	0.12	0.13
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.57	0.55	0.21	0.21	0.18	0.18
New England	1.20	1.00	0.42	0.24	0.34	0.20
Middle Atlantic	0.64	0.65	0.25	0.27	0.22	0.23
Midwest	0.55	0.60	0.32	0.31	0.29	0.30
East North Central	0.63	0.68	0.40	0.37	0.36	0.35
West North Central	1.02	1.23	0.57	0.54	0.52	0.53
South	0.57	0.53	0.24	0.26	0.20	0.24
South Atlantic	0.81	0.74	0.31	0.32	0.28	0.29
East South Central	1.30	1.25	0.67	0.75	0.58	0.70
West South Central	0.90	0.95	0.44	0.51	0.35	0.45
West	0.75	0.66	0.26	0.28	0.23	0.25
Mountain	1.18	1.10	0.48	0.56	0.44	0.53
Pacific	0.94	0.85	0.32	0.31	0.27	0.28
<b>COUNTY TYPE</b>						
Large Metro	0.44	0.42	0.16	0.17	0.14	0.15
Small Metro	0.56	0.50	0.27	0.24	0.24	0.22
250K – 1 Mil. Pop.	0.67	0.60	0.33	0.27	0.29	0.25
< 250K Pop.	0.95	0.93	0.46	0.49	0.44	0.45
Nonmetro	0.74	0.82	0.37	0.45	0.33	0.42
Urbanized	1.18	1.10	0.58	0.58	0.43	0.55
Less Urbanized	1.08	1.30	0.55	0.71	0.52	0.66
Completely Rural	2.45	2.36	1.11	1.61	1.10	1.57

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 2.61P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:  
P Values from Tests of Differences of Percentages, 2012 Versus 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2011)</b>	<b>Lifetime (2012)</b>	<b>Past Year (2011)</b>	<b>Past Year (2012)</b>	<b>Past Month (2011)</b>	<b>Past Month (2012)</b>
<b>TOTAL</b>	0.4758		0.4631		0.0797	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.8793		0.7182		0.4335	
New England	0.9785		0.1238		0.2402	
Middle Atlantic	0.8727		0.2461		0.1650	
Midwest	0.3206		0.7019		0.7576	
East North Central	0.3478		0.5834		0.9471	
West North Central	0.6877		0.8883		0.6512	
South	0.2325		0.4108		0.1153	
South Atlantic	0.5562		0.8615		0.9646	
East South Central	0.6283		0.4526		0.2396	
West South Central	0.2944		0.2896		0.0746	
West	0.4757		0.5641		0.4314	
Mountain	0.6043		0.3229		0.3598	
Pacific	0.2782		0.9763		0.7518	
<b>COUNTY TYPE</b>						
Large Metro	0.1294		0.9358		0.4858	
Small Metro	0.4411		0.5177		0.7570	
250K – 1 Mil. Pop.	0.3198		0.2589		0.4072	
< 250K Pop.	0.9635		0.7098		0.6143	
Nonmetro	0.0543		0.0220		0.0037	
Urbanized	0.3375		0.1091		0.0017	
Less Urbanized	0.2081		0.2164		0.2351	
Completely Rural	0.1193		0.1974		0.2901	

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.