

Table 6.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	4,962	5,281	7,285	7,900	5,022	5,407	5,185	5,275
Marijuana and Hashish	4,149	4,476	6,302	6,656	4,106	4,252	3,513	3,471
Cocaine	646	678	479	561	125	199	120	212
Crack	86	169	112	146	5	14	26 ^a	115
Heroin	53	64	65	75	51	51	112	145
Hallucinogens	314 ^a	469	370	429	160	102	128	134
LSD	56	79	55	75	15	14	25	16
PCP	6	16	18	8	*	4	*	*
Ecstasy	187	287	205	229	108	68	44	44
Inhalants	133	91	152	195	110	68	229 ^a	139
Nonmedical Use of Psychotherapeutics ^{3,4}	1,416	1,473	1,624 ^a	2,001	1,215	1,456	1,864	1,901
Pain Relievers	954	1,007	1,241	1,499	828	923	1,448	1,434
OxyContin [®]	126	125	161	91	49	41	98	102
Tranquilizers	499	491	529	663	395	486	418	499
Stimulants ⁴	359	354	232 ^a	382	150	244	228	191
Methamphetamine ⁴	155	88	60	161	72	87	152	104
Sedatives	48	35	32 ^a	118	49	54	101	63
ILLICIT DRUGS OTHER THAN MARIJUANA²	2,025	2,196	2,270 ^a	2,737	1,484	1,717	2,241	2,233

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	31.3	31.0	17.2	18.5	6.7	7.1	4.2	4.2
Marijuana and Hashish	26.2	26.3	14.8	15.6	5.5	5.6	2.8	2.8
Cocaine	4.1	4.0	1.1	1.3	0.2	0.3	0.1	0.2
Crack	0.5	1.0	0.3	0.3	0.0	0.0	0.0 ^a	0.1
Heroin	0.3	0.4	0.2	0.2	0.1	0.1	0.1	0.1
Hallucinogens	2.0	2.8	0.9	1.0	0.2	0.1	0.1	0.1
LSD	0.4	0.5	0.1	0.2	0.0	0.0	0.0	0.0
PCP	0.0	0.1	0.0	0.0	*	0.0	*	*
Ecstasy	1.2	1.7	0.5	0.5	0.1	0.1	0.0	0.0
Inhalants	0.8	0.5	0.4	0.5	0.1	0.1	0.2 ^a	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	8.9	8.7	3.8 ^a	4.7	1.6	1.9	1.5	1.5
Pain Relievers	6.0	5.9	2.9	3.5	1.1	1.2	1.2	1.2
OxyContin [®]	0.8	0.7	0.4	0.2	0.1	0.1	0.1	0.1
Tranquilizers	3.1	2.9	1.2	1.6	0.5	0.6	0.3	0.4
Stimulants ⁴	2.3	2.1	0.5 ^a	0.9	0.2	0.3	0.2	0.2
Methamphetamine ⁴	1.0	0.5	0.1	0.4	0.1	0.1	0.1	0.1
Sedatives	0.3	0.2	0.1 ^a	0.3	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	12.8	12.9	5.3 ^a	6.4	2.0	2.3	1.8	1.8

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	267	221	658	621	441	406	1,157	1,128
Marijuana and Hashish	250	196	589	557	397	346	731	701
Cocaine	22 ^a	6	21	15	14	3	10	7
Crack	0	0	3	1	*	*	*	*
Heroin	1	*	2	*	10	*	1	*
Hallucinogens	46	46	76	53	35 ^a	11	58	50
LSD	8	7	21	9	1	5	4	6
PCP	2	*	2	1	*	2	*	2
Ecstasy	31	22	37	27	23	9	17	16
Inhalants	13	26	43	34	20	27	147 ^a	101
Nonmedical Use of Psychotherapeutics ^{3,4}	94	69	176	163	81	76	338	384
Pain Relievers	77 ^a	44	140	132	53	63	293	319
OxyContin [®]	17 ^a	5	23	30	2	4	9	9
Tranquilizers	42	27	62	42	23	11	35	60
Stimulants ⁴	29	21	21 ^a	44	22	12	25	40
Methamphetamine ⁴	12	6	9	12	8	3	7	9
Sedatives	2	3	7	8	5	0	9	15
ILLICIT DRUGS OTHER THAN MARIJUANA²	119	104	257	208	127	104	522	501

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	70.4	67.9	44.7	42.1	29.8	28.6	5.3	5.2
Marijuana and Hashish	66.1	60.2	40.1	37.8	26.9	24.4	3.4	3.2
Cocaine	5.8 ^a	1.7	1.4	1.0	1.0	0.2	0.0	0.0
Crack	0.0	0.0	0.2	0.0	*	*	*	*
Heroin	0.3	*	0.1	*	0.7	*	0.0	*
Hallucinogens	12.1	14.2	5.2	3.6	2.4 ^a	0.8	0.3	0.2
LSD	2.0	2.1	1.4	0.6	0.1	0.3	0.0	0.0
PCP	0.4	*	0.1	0.0	*	0.2	*	0.0
Ecstasy	8.3	6.6	2.5	1.8	1.5	0.6	0.1	0.1
Inhalants	3.4	7.9	2.9	2.3	1.4	1.9	0.7 ^a	0.5
Nonmedical Use of Psychotherapeutics ^{3,4}	24.7	21.3	12.0	11.1	5.5	5.4	1.6	1.8
Pain Relievers	20.3	13.5	9.5	9.0	3.6	4.4	1.4	1.5
OxyContin [®]	4.4	1.7	1.6	2.0	0.1	0.3	0.0	0.0
Tranquilizers	11.2	8.4	4.2	2.8	1.6	0.8	0.2	0.3
Stimulants ⁴	7.6	6.4	1.5 ^a	3.0	1.5	0.8	0.1	0.2
Methamphetamine ⁴	3.1	1.8	0.6	0.8	0.6	0.2	0.0	0.0
Sedatives	0.6	0.8	0.5	0.5	0.4	0.0	0.0	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	31.5	31.9	17.5	14.1	8.6	7.3	2.4	2.3

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	2,019	2,061	2,903	2,744	1,391	1,306	1,038 ^b	1,270
Marijuana and Hashish	1,809	1,869	2,659	2,468	1,226	1,169	831	980
Cocaine	247	205	170	140	27	22	20	31
Crack	7	18	12	7	1	2	1 ^a	10
Heroin	21	22	33	36	31	15	22 ^a	65
Hallucinogens	198	272	229	229	79	46	52	52
LSD	47	39	34	60	8	9	4	10
PCP	1	6	10	4	1	2	*	*
Ecstasy	108 ^a	190	135	127	46	26	27	18
Inhalants	49	41	49	45	36	11	11	25
Nonmedical Use of Psychotherapeutics ^{3,4}	574	597	569	630	290	253	279	352
Pain Relievers	394	391	436	440	201	184	189 ^b	295
OxyContin [®]	52	48	61	37	27	26	18	34
Tranquilizers	182	167	173	194	105	95	85	81
Stimulants ⁴	166	186	95	140	42	36	41	37
Methamphetamine ⁴	25	36	17	31	14	17	23	13
Sedatives	14	19	9	21	12	6	8	6
ILLCIT DRUGS OTHER THAN MARIJUANA²	824	832	840	872	378	303	342	428

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	48.5	47.0	30.6	29.6	19.4	18.2	7.7 ^b	9.2
Marijuana and Hashish	43.4	42.6	28.0	26.6	17.1	16.3	6.2	7.1
Cocaine	5.9	4.7	1.8	1.5	0.4	0.3	0.2	0.2
Crack	0.2	0.4	0.1	0.1	0.0	0.0	0.0 ^a	0.1
Heroin	0.5	0.5	0.3	0.4	0.4	0.2	0.2 ^a	0.5
Hallucinogens	4.7	6.2	2.4	2.5	1.1	0.6	0.4	0.4
LSD	1.1	0.9	0.4	0.7	0.1	0.1	0.0	0.1
PCP	0.0	0.1	0.1	0.0	0.0	0.0	*	*
Ecstasy	2.6 ^a	4.3	1.4	1.4	0.6	0.4	0.2	0.1
Inhalants	1.2	0.9	0.5	0.5	0.5	0.2	0.1	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	13.8	13.6	6.0	6.8	4.0	3.5	2.1	2.6
Pain Relievers	9.5	8.9	4.6	4.7	2.8	2.6	1.4 ^a	2.1
OxyContin [®]	1.2	1.1	0.6	0.4	0.4	0.4	0.1	0.2
Tranquilizers	4.4	3.8	1.8	2.1	1.5	1.3	0.6	0.6
Stimulants ⁴	4.0	4.2	1.0 ^a	1.5	0.6	0.5	0.3	0.3
Methamphetamine ⁴	0.6	0.8	0.2	0.3	0.2	0.2	0.2	0.1
Sedatives	0.3	0.4	0.1	0.2	0.2	0.1	0.1	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	19.8	19.0	8.8	9.4	5.3	4.2	2.5	3.1

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	2,676	2,999	3,724 ^a	4,534	3,190	3,695	2,990	2,877
Marijuana and Hashish	2,090	2,411	3,053 ^a	3,631	2,483	2,738	1,952	1,791
Cocaine	377	468	288	407	84	174	89	174
Crack	79	150	97	138	4	12	24 ^a	104
Heroin	32	42	31	39	10	35	88	80
Hallucinogens	70	151	65	147	46	45	18	32
LSD	1	33	*	6	7	*	17	*
PCP	3	11	7	4	*	*	*	*
Ecstasy	48	76	33	74	39	33	*	11
Inhalants	71 ^a	24	60	116	54	30	71	13
Nonmedical Use of Psychotherapeutics ^{3,4}	749	807	879 ^a	1,207	845	1,126	1,247	1,165
Pain Relievers	483	571	665	926	574	676	966	820
OxyContin [®]	57	72	77	25	20	10	71	59
Tranquilizers	274	296	294	427	267	380	298	358
Stimulants ⁴	165	148	116	197	86	196	162	114
Methamphetamine ⁴	118	46	34	118	49	67	122	81
Sedatives	32	13	17 ^a	90	32	48	84	41
ILLICIT DRUGS OTHER THAN MARIJUANA²	1,081	1,260	1,173 ^a	1,657	978	1,310	1,376	1,304

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	23.6	24.4	11.8 ^a	14.2	4.8	5.5	3.4	3.2
Marijuana and Hashish	18.5	19.6	9.7	11.4	3.7	4.1	2.2	2.0
Cocaine	3.3	3.8	0.9	1.3	0.1	0.3	0.1	0.2
Crack	0.7	1.2	0.3	0.4	0.0	0.0	0.0 ^a	0.1
Heroin	0.3	0.3	0.1	0.1	0.0	0.1	0.1	0.1
Hallucinogens	0.6	1.2	0.2	0.5	0.1	0.1	0.0	0.0
LSD	0.0	0.3	*	0.0	0.0	*	0.0	*
PCP	0.0	0.1	0.0	0.0	*	*	*	*
Ecstasy	0.4	0.6	0.1	0.2	0.1	0.0	*	0.0
Inhalants	0.6 ^a	0.2	0.2	0.4	0.1	0.0	0.1	0.0
Nonmedical Use of Psychotherapeutics ^{3,4}	6.6	6.6	2.8 ^a	3.8	1.3	1.7	1.4	1.3
Pain Relievers	4.3	4.6	2.1	2.9	0.9	1.0	1.1	0.9
OxyContin [®]	0.5	0.6	0.2	0.1	0.0	0.0	0.1	0.1
Tranquilizers	2.4	2.4	0.9	1.3	0.4	0.6	0.3	0.4
Stimulants ⁴	1.5	1.2	0.4	0.6	0.1	0.3	0.2	0.1
Methamphetamine ⁴	1.0	0.4	0.1	0.4	0.1	0.1	0.1	0.1
Sedatives	0.3	0.1	0.1 ^a	0.3	0.0	0.1	0.1	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	9.6	10.2	3.7 ^a	5.2	1.5	1.9	1.5	1.5

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS¹	17,269 ^a	18,588	3,985	3,683	13,284 ^b	14,904
Marijuana or Hashish	14,557	15,384	3,674 ^a	3,338	10,884 ^a	12,046
Cocaine (Including Crack)	1,250	1,438	253 ^b	154	997	1,285
Heroin	169	190	43	30	126	160
Hallucinogens	844	999	381	388	463	611
Inhalants	395	353	146	138	249	216
Pain Relievers ²	3,023	3,428	653	611	2,370 ^a	2,817
Tranquilizers ²	1,423	1,640	328 ^a	234	1,095	1,406
Stimulants ^{2,3}	742 ^a	980	194	203	548 ^a	777
Methamphetamine ³	287	336	47	48	240	288
Sedatives ²	130	207	30	28	100	180
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5,779 ^a	6,650	1,326	1,193	4,453 ^b	5,456

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS¹	12.9	13.7	41.3	39.6	10.7 ^a	11.8
Marijuana or Hashish	10.9	11.4	38.1	35.9	8.8	9.5
Cocaine (Including Crack)	0.9	1.1	2.6 ^b	1.7	0.8	1.0
Heroin	0.1	0.1	0.4	0.3	0.1	0.1
Hallucinogens	0.6	0.7	4.0	4.2	0.4	0.5
Inhalants	0.3	0.3	1.5	1.5	0.2	0.2
Pain Relievers ²	2.3	2.5	6.8	6.6	1.9	2.2
Tranquilizers ²	1.1	1.2	3.4 ^a	2.5	0.9	1.1
Stimulants ^{2,3}	0.6 ^a	0.7	2.0	2.2	0.4	0.6
Methamphetamine ³	0.2	0.2	0.5	0.5	0.2	0.2
Sedatives ²	0.1	0.2	0.3	0.3	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4.3 ^a	4.9	13.7	12.8	3.6 ^b	4.3

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS²	7,929 ^b	9,188	1,904	1,881	6,025 ^b	7,307
Marijuana or Hashish	7,276 ^b	8,367	1,812	1,803	5,464 ^b	6,564
Cocaine (Including Crack)	279 ^a	502	52	35	227 ^a	468
Heroin	55	56	14	6	42	51
Hallucinogens	98	111	30	33	68	78
Inhalants	35	13	17	11	18	*
Pain Relievers ³	470	573	78	59	392	515
Tranquilizers ³	84	142	26	17	58	125
Stimulants ³	66	100	4	9	63	91
Methamphetamine ⁴	80	184	6	13	74	171
Sedatives ³	35	14	3	1	31	13
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,081 ^a	1,398	210	161	871 ^a	1,237

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS²	6.0 ^b	6.8	20.1	20.6	4.9 ^b	5.8
Marijuana or Hashish	5.5 ^b	6.2	19.2	19.8	4.4 ^b	5.2
Cocaine (Including Crack)	0.2 ^a	0.4	0.5	0.4	0.2 ^a	0.4
Heroin	0.0	0.0	0.1	0.1	0.0	0.0
Hallucinogens	0.1	0.1	0.3	0.4	0.1	0.1
Inhalants	0.0	0.0	0.2	0.1	0.0	*
Pain Relievers ³	0.4	0.4	0.8	0.6	0.3	0.4
Tranquilizers ³	0.1	0.1	0.3	0.2	0.0	0.1
Stimulants ³	0.0	0.1	0.0	0.1	0.1	0.1
Methamphetamine ⁴	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives ³	0.0	0.0	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.8	1.0	2.2	1.7	0.7 ^a	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.17A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLICIT DRUGS¹	17,269 ^a	18,588	1,365	1,248	6,314	6,111	9,590 ^b	11,228
Marijuana or Hashish	14,557	15,384	1,237 ^a	1,099	5,695	5,505	7,626 ^a	8,780
Cocaine (Including Crack)	1,250	1,438	57 ^b	23	444	367	749	1,049
Heroin	169	190	13	*	85	73	72	117
Hallucinogens	844	999	157 ^a	110	505	547	182	342
Inhalants	395	353	76	87	133	98	186	169
Pain Relievers ²	3,023	3,428	270	239	1,031	1,016	1,722 ^a	2,173
Tranquilizers ²	1,423	1,640	127 ^a	80	461	456	834	1,104
Stimulants ^{2,3}	742 ^a	980	73	77	303	362	366	540
Methamphetamine ³	287	336	29	21	56	84	201	231
Sedatives ²	130	207	15	11	35	45	80	151
ILLICIT DRUGS OTHER THAN MARIJUANA¹	5,779 ^a	6,650	503 ^a	417	2,042	2,007	3,233 ^b	4,226

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.17B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS¹	12.9	13.7	41.0	38.8	30.3	29.3	8.8 ^b	10.1
Marijuana or Hashish	10.9	11.4	37.2	34.2	27.4	26.4	7.0 ^a	7.9
Cocaine (Including Crack)	0.9	1.1	1.7 ^b	0.7	2.1	1.8	0.7	0.9
Heroin	0.1	0.1	0.4	*	0.4	0.4	0.1	0.1
Hallucinogens	0.6	0.7	4.7	3.4	2.4	2.6	0.2	0.3
Inhalants	0.3	0.3	2.3	2.7	0.6	0.5	0.2	0.2
Pain Relievers ²	2.3	2.5	8.1	7.4	5.0	4.9	1.6	2.0
Tranquilizers ²	1.1	1.2	3.8 ^a	2.5	2.2	2.2	0.8	1.0
Stimulants ^{2,3}	0.6 ^a	0.7	2.2	2.4	1.5	1.7	0.3	0.5
Methamphetamine ³	0.2	0.2	0.9	0.7	0.3	0.4	0.2	0.2
Sedatives ²	0.1	0.2	0.4	0.3	0.2	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4.3 ^a	4.9	15.1	13.0	9.8	9.6	3.0 ^b	3.8

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS²	7,929 ^b	9,188	619	576	3,162	3,200	4,147 ^b	5,411
Marijuana or Hashish	7,276 ^b	8,367	582	546	2,993	3,048	3,701 ^b	4,773
Cocaine (Including Crack)	279 ^a	502	12	5	95	83	171 ^a	414
Heroin	55	56	5	*	29	11	22	46
Hallucinogens	98	111	11	9	62	56	25	46
Inhalants	35	13	1	5	16	8	18	*
Pain Relievers ³	470	573	31	23	161	164	277	387
Tranquilizers ³	84	142	10	6	40	42	34	94
Stimulants ³	66	100	2	6	23	24	41	71
Methamphetamine ⁴	80	184	6	3	5 ^a	31	70	150
Sedatives ³	35	14	3	1	*	2	30	11
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,081 ^a	1,398	71	55	393	355	617 ^a	988

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS²	6.0 ^b	6.8	19.2	18.5	15.3	15.5	3.8 ^b	4.9
Marijuana or Hashish	5.5 ^b	6.2	18.2	17.5	14.5	14.8	3.4 ^b	4.3
Cocaine (Including Crack)	0.2 ^a	0.4	0.4	0.2	0.5	0.4	0.2 ^a	0.4
Heroin	0.0	0.0	0.1	*	0.1	0.1	0.0	0.0
Hallucinogens	0.1	0.1	0.3	0.3	0.3	0.3	0.0	0.0
Inhalants	0.0	0.0	0.0	0.2	0.1	0.0	0.0	*
Pain Relievers ³	0.4	0.4	1.0	0.7	0.8	0.8	0.3	0.3
Tranquilizers ³	0.1	0.1	0.3	0.2	0.2	0.2	0.0	0.1
Stimulants ³	0.0	0.1	0.1	0.2	0.1	0.1	0.0	0.1
Methamphetamine ⁴	0.1	0.1	0.2	0.1	0.0 ^a	0.1	0.1	0.1
Sedatives ³	0.0	0.0	0.1	0.0	*	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.8	1.0	2.2	1.7	1.9	1.7	0.6 ^a	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.