

Table 6.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	11,406	11,973
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,256	3,483	4,987	5,553
Cigars	6,691	7,261	6,174	6,156
Pipe Tobacco	1,149	1,283	988	1,247
ALCOHOL	37,764	37,643	95,621	97,859
Binge Alcohol Use ²	24,154	25,082	34,187	34,604
Heavy Alcohol Use ²	8,716	9,084	7,148	7,937

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	5.7	5.9
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	5.7	6.1	2.5	2.7
Cigars	11.8	12.6	3.1	3.0
Pipe Tobacco	2.0	2.2	0.5	0.6
ALCOHOL	66.5	65.4	47.6	48.3
Binge Alcohol Use ²	42.5	43.6	17.0	17.1
Heavy Alcohol Use ²	15.3	15.8	3.6	3.9

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	551	518
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	278	259	239	257
Cigars	549 ^b	393	309	252
Pipe Tobacco	120	104	62	61
ALCOHOL	1,171 ^a	1,003	2,156	2,214
Binge Alcohol Use ²	864 ^a	742	985	1,057
Heavy Alcohol Use ²	249	199	129	127

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	2.4	2.2
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	14.3	15.8	1.0	1.1
Cigars	28.3 ^a	24.1	1.3 ^a	1.1
Pipe Tobacco	6.2	6.4	0.3	0.3
ALCOHOL	60.4	61.4	9.4	9.5
Binge Alcohol Use ²	44.6	45.4	4.3	4.5
Heavy Alcohol Use ²	12.9	12.2	0.6	0.5

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	2,044	2,159
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,159	1,137	708	774
Cigars	2,439	2,321	1,285	1,395
Pipe Tobacco	399	386	236	239
ALCOHOL	8,880	8,417	11,934	12,411
Binge Alcohol Use ²	6,948	6,658	6,705	7,004
Heavy Alcohol Use ²	2,646	2,641	1,519 ^a	1,746

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	9.0	9.2
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	10.1	10.3	3.1	3.3
Cigars	21.2	21.1	5.6	5.9
Pipe Tobacco	3.5	3.5	1.0	1.0
ALCOHOL	77.3	76.5	52.3	52.6
Binge Alcohol Use ²	60.5	60.5	29.4	29.7
Heavy Alcohol Use ²	23.0	24.0	6.7	7.4

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	8,810	9,296
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,819	2,087	4,040	4,522
Cigars	3,702 ^b	4,546	4,581	4,509
Pipe Tobacco	630	794	690	947
ALCOHOL	27,713	28,224	81,532	83,234
Binge Alcohol Use ²	16,342 ^a	17,682	26,496	26,542
Heavy Alcohol Use ²	5,821	6,245	5,500	6,064

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	5.7	6.0
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.2	4.7	2.6	2.9
Cigars	8.5 ^a	10.1	3.0	2.9
Pipe Tobacco	1.5	1.8	0.4	0.6
ALCOHOL	63.9	62.9	52.6	53.5
Binge Alcohol Use ²	37.7	39.4	17.1	17.1
Heavy Alcohol Use ²	13.4	13.9	3.5	3.9

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.