

**Table 6.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	10,416	10,959	19,080	19,743	16,795	15,765	21,934	23,031
Cigarettes	8,716	9,084	15,437	15,998	13,611	12,561	19,054	19,882
Smokeless Tobacco	1,850	2,121	2,598	2,806	1,434	1,585	2,360	2,524
Cigars	2,410 <sup>a</sup>	2,946	4,268	4,542	3,399	3,201	2,787	2,727
Pipe Tobacco	316	330	662	642	554	794	606	765

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	65.7	64.4	44.9	46.3	22.4 <sup>a</sup>	20.8	17.7	18.5
Cigarettes	54.9	53.4	36.3	37.5	18.1 <sup>a</sup>	16.6	15.3	16.0
Smokeless Tobacco	11.7	12.5	6.1	6.6	1.9	2.1	1.9	2.0
Cigars	15.2	17.3	10.0	10.6	4.5	4.2	2.2	2.2
Pipe Tobacco	2.0	1.9	1.6	1.5	0.7	1.0	0.5	0.6

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	298	239	765	683	390	346	1,037 <sup>a</sup>	885
Cigarettes	249	199	615	543	307	260	768 <sup>b</sup>	632
Smokeless Tobacco	88	63	157	162	65	48	207	242
Cigars	127 <sup>a</sup>	89	298	245	136	110	297 <sup>b</sup>	201
Pipe Tobacco	29	17	82	52	17 <sup>a</sup>	44	55	53

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	78.7	73.2	52.0 <sup>a</sup>	46.4	26.4	24.4	4.8 <sup>b</sup>	4.1
Cigarettes	65.8	61.0	41.8 <sup>a</sup>	36.9	20.8	18.4	3.5 <sup>b</sup>	2.9
Smokeless Tobacco	23.1	19.5	10.7	11.0	4.4	3.4	1.0	1.1
Cigars	33.4	27.2	20.3	16.7	9.2	7.8	1.4 <sup>b</sup>	0.9
Pipe Tobacco	7.6	5.2	5.6 <sup>a</sup>	3.5	1.1 <sup>a</sup>	3.1	0.3	0.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	3,058	3,121	5,107	4,807	2,311	2,214	3,056	3,024
Cigarettes	2,646	2,641	4,301	4,017	1,933	1,759	2,607	2,591
Smokeless Tobacco	605 <sup>a</sup>	722	758	647	164	199	341	344
Cigars	1,017	1,088	1,331	1,356	663	622	714	650
Pipe Tobacco	148	156	249	215	128	129	110	124

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	73.4	71.2	53.8	51.8	32.3	30.9	22.7	22.0
Cigarettes	63.5	60.2	45.3	43.3	27.0 <sup>a</sup>	24.5	19.3	18.8
Smokeless Tobacco	14.5	16.5	8.0	7.0	2.3	2.8	2.5	2.5
Cigars	24.4	24.8	14.0	14.6	9.3	8.7	5.3	4.7
Pipe Tobacco	3.5	3.6	2.6	2.3	1.8	1.8	0.8	0.9

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	7,059	7,599	13,209	14,253	14,094	13,204	17,841	19,122
Cigarettes	5,821	6,245	10,521	11,437	11,371	10,542	15,679	16,659
Smokeless Tobacco	1,158	1,336	1,683	1,998	1,205	1,337	1,812	1,938
Cigars	1,267 <sup>a</sup>	1,769	2,639	2,941	2,600	2,469	1,777	1,877
Pipe Tobacco	140	157	331	374	409	621	441	588

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 6.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2011 and 2012**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
<b>TOBACCO PRODUCTS<sup>2</sup></b>	62.4	61.7	41.9	44.7	21.2	19.6	20.0	21.5
Cigarettes	51.4	50.7	33.4	35.8	17.1	15.7	17.6	18.7
Smokeless Tobacco	10.2	10.9	5.3	6.3	1.8	2.0	2.0	2.2
Cigars	11.2 <sup>a</sup>	14.4	8.4	9.2	3.9	3.7	2.0	2.1
Pipe Tobacco	1.2	1.3	1.0	1.2	0.6	0.9	0.5	0.7

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.