

Table 6.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1,978	2,133	253	241	497	481	1,228	1,412
From More Than One Doctor	236	198	27	25	65	46	144	127
Wrote Fake Prescription	21	23	6	8	2 ^a	10	13	5
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	28 ^a	91	19	12	3 ^a	17	6 ^a	62
From Friend or Relative for Free	6,278	5,838	575	515	2,020 ^b	1,688	3,683	3,636
Bought from Friend or Relative	1,287	1,179	111	103	535	499	640	577
Took from Friend or Relative without Asking	548 ^a	431	139 ^a	107	169	145	239	180
Bought from Drug Dealer or Other Stranger	502	468	50	37	239	223	213	208
Bought on the Internet	45	21	0	1	3	5	42	14
Some Other Way	483	433	96 ^a	70	146	114	241	249

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	17.3 ^a	19.7	19.8	21.5	13.5	14.9	19.0	21.8
From More Than One Doctor	2.1	1.8	2.1	2.2	1.8	1.4	2.2	2.0
Wrote Fake Prescription	0.2	0.2	0.5	0.7	0.1 ^a	0.3	0.2	0.1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2 ^a	0.8	1.5	1.0	0.1 ^a	0.5	0.1 ^a	1.0
From Friend or Relative for Free	55.0	54.0	45.0	46.1	54.9	52.3	57.1	56.2
Bought from Friend or Relative	11.3	10.9	8.7	9.2	14.6	15.5	9.9	8.9
Took from Friend or Relative without Asking	4.8	4.0	10.9	9.6	4.6	4.5	3.7	2.8
Bought from Drug Dealer or Other Stranger	4.4	4.3	3.9	3.3	6.5	6.9	3.3	3.2
Bought on the Internet	0.4	0.2	0.0	0.1	0.1	0.2	0.6	0.2
Some Other Way	4.2	4.0	7.5	6.3	4.0	3.5	3.7	3.9

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	4,469	4,331	287	262	1,290 ^b	1,124	2,892	2,945
From More Than One Doctor	201	188	28	31	69	71	104	86
Wrote Fake Prescription	13	17	1	*	1	3	11	14
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	14	19	5	3	8	14	2	2
From Friend or Relative for Free	357	286	47	37	150 ^a	107	161	142
Bought from Friend or Relative	253	216	30	31	129	109	95	77
Took from Friend or Relative without Asking	110 ^a	68	51 ^a	28	38	32	21	8
Bought from Drug Dealer or Other Stranger	130 ^a	74	17	13	69 ^a	39	44	21
Bought on the Internet	11	11	1	0	3	1	7	10
Some Other Way	71	60	22	17	22	14	27	28

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	79.4	82.2	58.8	62.0	72.5	74.2	86.0	88.3
From More Than One Doctor	3.6	3.6	5.7	7.3	3.9	4.7	3.1	2.6
Wrote Fake Prescription	0.2	0.3	0.2	*	0.1	0.2	0.3	0.4
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.4	1.0	0.7	0.4	0.9	0.0	0.1
From Friend or Relative for Free	6.3	5.4	9.6	8.7	8.4	7.0	4.8	4.3
Bought from Friend or Relative	4.5	4.1	6.1	7.3	7.2	7.2	2.8	2.3
Took from Friend or Relative without Asking	2.0	1.3	10.5 ^a	6.6	2.1	2.1	0.6	0.2
Bought from Drug Dealer or Other Stranger	2.3 ^a	1.4	3.5	3.2	3.9	2.6	1.3	0.6
Bought on the Internet	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.3
Some Other Way	1.3	1.1	4.5	4.1	1.2	0.9	0.8	0.9

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1,280	1,152	125	114	293	256	863	782
From More Than One Doctor	232	163	25	16	63	44	144	103
Wrote Fake Prescription	25	30	5	7	2	2	17	21
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	30	66	12	9	6	12	11	44
From Friend or Relative for Free	2,851 ^a	2,537	245	216	944 ^b	734	1,661	1,587
Bought from Friend or Relative	1,254	1,093	87	72	491 ^b	370	676	651
Took from Friend or Relative without Asking	525	427	87 ^a	63	185 ^a	134	253	230
Bought from Drug Dealer or Other Stranger	541	515	55	40	270	246	216	229
Bought on the Internet	48	46	*	3	13	15	35	29
Some Other Way	303 ^a	196	41	37	97	77	164	82

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	27.1	27.3	24.9	25.9	20.3	21.9	31.1	29.9
From More Than One Doctor	4.9	3.9	4.9	3.7	4.4	3.7	5.2	3.9
Wrote Fake Prescription	0.5	0.7	1.1	1.6	0.2	0.2	0.6	0.8
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6	1.6	2.4	2.0	0.4	1.1	0.4	1.7
From Friend or Relative for Free	60.3	60.0	48.7	49.0	65.4	62.7	59.8	60.6
Bought from Friend or Relative	26.6	25.9	17.5	16.4	34.0	31.7	24.4	24.9
Took from Friend or Relative without Asking	11.1	10.1	17.4	14.4	12.8	11.5	9.1	8.8
Bought from Drug Dealer or Other Stranger	11.5	12.2	11.0	9.2	18.7	21.0	7.8	8.7
Bought on the Internet	1.0	1.1	*	0.6	0.9	1.3	1.3	1.1
Some Other Way	6.6	4.7	8.6	8.7	6.9	6.7	6.0	3.2

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1,978	1,874	116	113	632 ^b	504	1,230	1,257
From More Than One Doctor	223	175	24	23	81	75	119	76
Wrote Fake Prescription	31	44	1	2	9	3	21	39
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	23	27	3	3	16	8	4	16
From Friend or Relative for Free	376	318	40	26	159 ^b	105	177	187
Bought from Friend or Relative	344	288	38	25	159 ^a	115	147	148
Took from Friend or Relative without Asking	112	99	28	18	53	45	31	37
Bought from Drug Dealer or Other Stranger	220	176	23	15	117 ^a	81	80	80
Bought on the Internet	9	22	1	*	5	5	3	16
Some Other Way	75	64	18	10	36	22	20	33

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	76.8	79.2	54.7	62.3	72.2	74.5	82.6	83.3
From More Than One Doctor	8.8	7.4	11.6	12.9	9.3	11.2	8.0	5.1
Wrote Fake Prescription	1.2	1.9	0.5	0.9	1.0	0.5	1.4	2.6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9	1.2	1.4	1.8	1.9	1.2	0.2	1.1
From Friend or Relative for Free	14.8	13.5	19.1	14.5	18.3	15.7	12.1	12.5
Bought from Friend or Relative	13.5	12.3	18.2	13.9	18.4	17.2	10.0	9.9
Took from Friend or Relative without Asking	4.4	4.2	13.5	9.8	6.1	6.8	2.1	2.4
Bought from Drug Dealer or Other Stranger	8.6	7.5	10.8	8.6	13.5	12.0	5.5	5.3
Bought on the Internet	0.4	0.9	0.7	*	0.6	0.8	0.2	1.1
Some Other Way	3.0	2.8	8.9	5.7	4.3	3.3	1.4	2.2

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	629	659	33	29	128	111	469	519
From More Than One Doctor	58	38	3	3	15	18	40	17
Wrote Fake Prescription	3	3	2	2	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	11	3	1	7	6	*	4
From Friend or Relative for Free	3,044	3,157	190	189	892 ^a	779	1,962	2,190
Bought from Friend or Relative	674	708	57	67	321	309	295	332
Took from Friend or Relative without Asking	190	159	37	38	75	65	78	56
Bought from Drug Dealer or Other Stranger	292	303	35	35	149	166	108	102
Bought on the Internet	21	1	1	*	3	1	17	*
Some Other Way	174	147	21	13	54	52	99	82

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	12.4	12.7	8.6	7.6	7.8	7.3	15.3	15.7
From More Than One Doctor	1.1	0.7	0.9	0.8	0.9	1.2	1.3	0.5
Wrote Fake Prescription	0.1	0.1	0.5	0.6	0.1	0.0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.2	0.8	0.2	0.5	0.4	*	0.1
From Friend or Relative for Free	59.7	60.9	49.8	50.0	54.2	51.7	63.9	66.3
Bought from Friend or Relative	13.2	13.7	14.9	17.7	19.5	20.5	9.6	10.1
Took from Friend or Relative without Asking	3.7	3.1	9.6	10.0	4.6	4.3	2.5	1.7
Bought from Drug Dealer or Other Stranger	5.7	5.8	9.3	9.4	9.0	11.0	3.5	3.1
Bought on the Internet	0.4	0.0	0.2	*	0.2	0.0	0.6	*
Some Other Way	3.4	2.8	5.4	3.6	3.3	3.4	3.2	2.5

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	2,122	2,340	89	74	520	500	1,513	1,765
From More Than One Doctor	85	95	7	7	35	29	43	59
Wrote Fake Prescription	5	1	*	*	1	1	2	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	10	*	1	1	3	5	6
From Friend or Relative for Free	237	209	23	24	113 ^b	68	102	117
Bought from Friend or Relative	83	122	12	21	56	41	15 ^b	60
Took from Friend or Relative without Asking	25	33	12	8	11	13	2	11
Bought from Drug Dealer or Other Stranger	76 ^a	43	9	11	42 ^a	21	25	11
Bought on the Internet	8	1	*	*	1	*	7	*
Some Other Way	24	25	7	3	4	15	13	8

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	79.4	81.3	55.3	49.6	66.3 ^a	72.2	87.7	86.7
From More Than One Doctor	3.2	3.3	4.5	4.5	4.4	4.2	2.5	2.9
Wrote Fake Prescription	0.2	0.0	*	*	0.1	0.2	0.1	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.3	*	0.5	0.1	0.5	0.3	0.3
From Friend or Relative for Free	8.9	7.2	14.3	15.8	14.3 ^a	9.8	5.9	5.7
Bought from Friend or Relative	3.1	4.2	7.3	13.9	7.2	5.9	0.9 ^a	2.9
Took from Friend or Relative without Asking	0.9	1.1	7.3	5.6	1.4	1.9	0.1	0.5
Bought from Drug Dealer or Other Stranger	2.8 ^a	1.5	5.4	7.5	5.4 ^a	3.1	1.4	0.5
Bought on the Internet	0.3	0.0	*	*	0.1	*	0.4	*
Some Other Way	0.9	0.9	4.1	2.0	0.6	2.1	0.7	0.4

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	314	353	13	12	68	57	233	285
From More Than One Doctor	17	15	*	2	4	10	11	3
Wrote Fake Prescription	2	3	1	2	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	21	*	*	*	4	*	*
From Friend or Relative for Free	1,116	1,092	42	60	277	250	798	783
Bought from Friend or Relative	421	349	26	28	159	124	236	198
Took from Friend or Relative without Asking	95	73	13	12	43	31	39	30
Bought from Drug Dealer or Other Stranger	144	156	17	18	77	74	50	64
Bought on the Internet	13	*	1	*	4	0	8	*
Some Other Way	58	57	5	7	15	16	38	34

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	16.6	18.8	12.1	9.4	12.4	11.5	18.9	22.6
From More Than One Doctor	0.9	0.8	*	1.7	0.7	2.0	0.9	0.2
Wrote Fake Prescription	0.1	0.2	1.2	1.3	0.1	0.3	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1	1.1	*	*	*	0.9	*	*
From Friend or Relative for Free	59.0	58.3	38.1	48.1	50.8	50.6	64.5	62.3
Bought from Friend or Relative	22.3	18.6	24.2	22.2	29.1	25.1	19.1	15.7
Took from Friend or Relative without Asking	5.0	3.9	12.4	9.3	7.9	6.4	3.2	2.4
Bought from Drug Dealer or Other Stranger	7.6	8.3	15.3	14.8	14.1	15.0	4.1	5.1
Bought on the Internet	0.7	*	1.0	*	0.8	0.0	0.6	*
Some Other Way	3.1	3.1	5.2	5.8	2.8	3.3	3.1	2.7

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	787	773	*	*	168	162	604	586
From More Than One Doctor	53	66	*	*	22	10	30	54
Wrote Fake Prescription	2	*	*	*	1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	7	*	*	1	3	*	*
From Friend or Relative for Free	131	92	*	*	49 ^a	29	76	52
Bought from Friend or Relative	67	56	*	*	37	27	25	17
Took from Friend or Relative without Asking	21	22	*	*	2	8	*	11
Bought from Drug Dealer or Other Stranger	56 ^a	23	*	*	19	14	33	5
Bought on the Internet	6	1	*	*	*	*	*	*
Some Other Way	8	16	*	*	4	*	2	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	76.6	76.7	*	*	65.6	70.1	81.9	80.5
From More Than One Doctor	5.2	6.6	*	*	8.5	4.2	4.1	7.4
Wrote Fake Prescription	0.1	*	*	*	0.6	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.7	*	*	0.6	1.3	*	*
From Friend or Relative for Free	12.7	9.2	*	*	19.2	12.5	10.3	7.2
Bought from Friend or Relative	6.6	5.6	*	*	14.4	11.7	3.4	2.4
Took from Friend or Relative without Asking	2.0	2.2	*	*	0.7	3.4	*	1.5
Bought from Drug Dealer or Other Stranger	5.4 ^a	2.3	*	*	7.5	6.2	4.5	0.7
Bought on the Internet	0.5	0.1	*	*	*	*	*	*
Some Other Way	0.8	1.6	*	*	1.5	*	0.3	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	228	241	21	23	85	64	122	153
From More Than One Doctor	23	27	4	3	5	19	14	6
Wrote Fake Prescription	10	3	3	*	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	21	10	5	2	2	5	14	3
From Friend or Relative for Free	1,130	1,151	97	108	533	497	501	546
Bought from Friend or Relative	422	477	45	42	274	284	103	152
Took from Friend or Relative without Asking	106	71	21	19	34	19	50	34
Bought from Drug Dealer or Other Stranger	142	136	19	16	63	77	60	43
Bought on the Internet	34	11	3	*	9	3	23	8
Some Other Way	87	59	8	4	29	30	50	25

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	10.4	11.0	9.3	10.7	8.2	6.5	13.0	15.8
From More Than One Doctor	1.1	1.2	1.8	1.2	0.5	1.9	1.5	0.6
Wrote Fake Prescription	0.4	0.1	1.2	*	0.0	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.0	0.4	2.3	0.9	0.2	0.5	1.4	0.3
From Friend or Relative for Free	51.3	52.7	42.8	49.6	51.5	49.8	53.0	56.3
Bought from Friend or Relative	19.1	21.8	20.0	19.1	26.5	28.4	10.9	15.7
Took from Friend or Relative without Asking	4.8	3.3	9.4	8.6	3.3	1.9	5.3	3.5
Bought from Drug Dealer or Other Stranger	6.5	6.2	8.5	7.4	6.1	7.7	6.4	4.4
Bought on the Internet	1.6	0.5	1.1	*	0.9	0.3	2.4	0.8
Some Other Way	3.9	2.7	3.5	1.8	2.8	3.0	5.3	2.6

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Stimulants ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	756	846	53	54	355	346	347	446
From More Than One Doctor	40	33	7	7	17	22	16	4
Wrote Fake Prescription	1	*	*	*	0	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	2	*	*	5	2	*	*
From Friend or Relative for Free	77	76	9	12	42	35	*	29
Bought from Friend or Relative	47	55	6	6	36	35	5	14
Took from Friend or Relative without Asking	7	12	3	5	3	7	1	*
Bought from Drug Dealer or Other Stranger	44	23	4	5	19	15	*	3
Bought on the Internet	8	3	*	1	*	0	*	*
Some Other Way	15	3	1	1	1	2	*	0

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Stimulants ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	75.5	80.3	63.0	59.9	74.2	74.6	79.4	89.4
From More Than One Doctor	4.0	3.1	8.3	7.7	3.5	4.8	3.7	0.8
Wrote Fake Prescription	0.1	*	*	*	0.1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5	0.2	*	*	1.0	0.4	*	*
From Friend or Relative for Free	7.7	7.2	10.5	12.8	8.7	7.6	*	5.9
Bought from Friend or Relative	4.7	5.2	7.2	6.5	7.6	7.5	1.1	2.8
Took from Friend or Relative without Asking	0.7	1.1	3.7	5.4	0.6	1.4	0.2	*
Bought from Drug Dealer or Other Stranger	4.4	2.1	4.9	5.2	4.1	3.3	*	0.6
Bought on the Internet	0.8	0.3	*	1.1	*	0.1	*	*
Some Other Way	1.5	0.3	0.6	1.4	0.3	0.4	*	0.0

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	172	140	12	13	57 ^b	25	103	*
From More Than One Doctor	24	19	2	3	5	4	*	*
Wrote Fake Prescription	3	*	2	*	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	12	*	1	1	3	*	*
From Friend or Relative for Free	407	356	34	39	183	163	190	*
Bought from Friend or Relative	209	236	15	23	123	115	71	*
Took from Friend or Relative without Asking	80	45	9	7	23	18	47	*
Bought from Drug Dealer or Other Stranger	89	103	16	11	38	44	35	47
Bought on the Internet	17	18	1	*	5	2	*	*
Some Other Way	32	39	4	3	7	12	21	25

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	20.3	18.9	14.8	17.7	16.3 ^b	8.3	24.7	*
From More Than One Doctor	2.9	2.5	2.8	4.2	1.6	1.4	*	*
Wrote Fake Prescription	0.3	*	2.7	*	0.2	0.0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	1.6	*	1.3	0.2	0.9	*	*
From Friend or Relative for Free	47.8	47.9	41.0	52.3	52.1	53.5	45.5	*
Bought from Friend or Relative	24.6	31.9	18.2 ^a	31.4	35.3	38.1	17.0	*
Took from Friend or Relative without Asking	9.4	6.1	11.4	9.8	6.7	6.1	11.3	*
Bought from Drug Dealer or Other Stranger	10.5	13.9	19.0	15.3	10.9	14.5	8.3	13.0
Bought on the Internet	2.0	2.5	1.2	*	1.5	0.7	*	*
Some Other Way	3.9	5.4	5.0	3.7	2.2	4.0	5.2	6.8

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Stimulants ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	267	271	*	*	120	131	*	*
From More Than One Doctor	20	19	*	*	12	4	*	*
Wrote Fake Prescription	0	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0	*	*	*	*	*	*	*
From Friend or Relative for Free	63	33	*	*	23	13	*	*
Bought from Friend or Relative	30	31	*	*	17	11	*	*
Took from Friend or Relative without Asking	11	17	*	*	*	5	*	*
Bought from Drug Dealer or Other Stranger	14	36	*	*	9	8	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	3	*	*	1	1	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specific questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Stimulants ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	73.2	83.4	*	*	75.6 ^a	86.2	*	*
From More Than One Doctor	5.6	5.9	*	*	7.5	2.8	*	*
Wrote Fake Prescription	0.0	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1	*	*	*	*	*	*	*
From Friend or Relative for Free	17.8	10.2	*	*	14.7	8.5	*	*
Bought from Friend or Relative	8.5	9.6	*	*	10.5	7.1	*	*
Took from Friend or Relative without Asking	3.2	5.4	*	*	*	3.5	*	*
Bought from Drug Dealer or Other Stranger	4.0 ^a	11.1	*	*	6.0	5.4	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	1.0	*	*	0.3	0.6	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.59A Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	392 ^b	182	*	*	129 ^b	54	226	*
Bought from Friend or Relative	288	*	9	*	55 ^b	10	224	*
Took from Friend or Relative without Asking	40	16	6	*	5	*	29	*
Bought from Drug Dealer or Other Stranger	174	*	10	*	58 ^b	22	105	*
Bought on the Internet	7	4	*	*	3	3	*	*
Some Other Way	40	16	*	*	6	*	28	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.59B Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	41.7	40.9	*	*	50.4	52.1	36.6	*
Bought from Friend or Relative	30.6	*	13.2	*	21.5 ^b	9.4	36.4	*
Took from Friend or Relative without Asking	4.2	3.7	9.1	*	2.0	*	4.6	*
Bought from Drug Dealer or Other Stranger	18.5	*	15.1	*	22.7	21.2	17.1	*
Bought on the Internet	0.7	0.8	*	*	1.0	2.7	*	*
Some Other Way	4.2	3.5	*	*	2.4	*	4.6	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	141	*	*	*	*	*	*	*
Bought from Friend or Relative	103	*	*	*	33	*	*	*
Took from Friend or Relative without Asking	11	1	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	52	*	*	*	19	*	*	*
Bought on the Internet	*	2	*	*	*	*	*	*
Some Other Way	9	15	*	*	7	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	44.3	*	*	*	*	*	*	*
Bought from Friend or Relative	32.5	*	*	*	31.0	*	*	*
Took from Friend or Relative without Asking	3.3	0.7	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	16.2	*	*	*	18.1	*	*	*
Bought on the Internet	*	1.3	*	*	*	*	*	*
Some Other Way	2.9 ^a	11.2	*	*	6.6	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	167	*	*	*	*	*	*	*
Bought from Friend or Relative	143	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	108	*	*	*	*	*	*	*
Bought on the Internet	4	*	*	*	*	*	*	*
Some Other Way	20	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	44.2	*	*	*	*	*	*	*
Bought from Friend or Relative	37.9	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	28.7	*	*	*	*	*	*	*
Bought on the Internet	1.0	*	*	*	*	*	*	*
Some Other Way	5.5	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	112	122	16	15	28	14	69	*
From More Than One Doctor	20	9	1	*	4	3	*	*
Wrote Fake Prescription	1	1	0	*	0	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	5	*	3	*	2	*	*
From Friend or Relative for Free	466 ^b	271	42	26	71	64	353	*
Bought from Friend or Relative	72	41	15	7	19	13	38	*
Took from Friend or Relative without Asking	40	17	9	5	17	*	14	*
Bought from Drug Dealer or Other Stranger	31	12	2	2	5	8	24	*
Bought on the Internet	*	0	*	0	*	0	*	*
Some Other Way	33	18	3	0	5	4	25	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	14.2 ^a	24.6	17.8	24.2	18.5	12.3	12.5	*
From More Than One Doctor	2.6	1.8	1.4	*	2.5	2.2	*	*
Wrote Fake Prescription	0.1	0.3	0.5	*	0.3	0.2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6	1.0	*	5.3	*	1.4	*	*
From Friend or Relative for Free	59.0	54.5	46.3	40.8	47.7	55.7	64.1	*
Bought from Friend or Relative	9.1	8.2	16.4	10.7	12.9	11.4	6.9	*
Took from Friend or Relative without Asking	5.1	3.4	10.3	8.4	11.6	*	2.5	*
Bought from Drug Dealer or Other Stranger	3.9	2.5	2.7	3.5	3.1	6.7	4.3	*
Bought on the Internet	*	0.0	*	0.1	*	0.1	*	*
Some Other Way	4.2	3.7	3.5	0.7	3.0	3.9	4.6	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Sedatives ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	371 ^a	227	*	*	*	*	*	*
From More Than One Doctor	4	7	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	*	*	*	*	*	*	*
From Friend or Relative for Free	24	12	*	*	*	*	*	*
Bought from Friend or Relative	5	3	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	6	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	8	5	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	3	1	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Sedatives ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	86.1	86.8	*	*	*	*	*	*
From More Than One Doctor	1.0	2.8	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.0	*	*	*	*	*	*	*
From Friend or Relative for Free	5.5	4.6	*	*	*	*	*	*
Bought from Friend or Relative	1.2	1.0	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.1	2.2	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	2.0	1.9	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.6	0.4	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	81	*	*	*	*	*	*	*
From More Than One Doctor	7	*	*	*	*	*	*	*
Wrote Fake Prescription	1	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	2	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	15	10	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	0	0	*	*	*	*	*	*
Some Other Way	19	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	23.7	*	*	*	*	*	*	*
From More Than One Doctor	2.1	*	*	*	*	*	*	*
Wrote Fake Prescription	0.2	0.5	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.6	1.1	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	4.2	5.0	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	0.0	0.0	*	*	*	*	*	*
Some Other Way	5.7	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Sedatives ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Source Where Friend or Relative Obtained Sedatives ¹	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.