

Table 6.71A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
ILLCIT DRUGS²	6,548	6,574	108	130	6,394	6,415
Marijuana and Hashish	4,910	5,122	89	115	4,792	4,985
Cocaine	387	344	3	4	380	338
Crack	99 ^a	47	2	*	96 ^a	45
Heroin	68	83	4	4	62	79
Hallucinogens	437 ^a	335	7	2	425 ^a	330
LSD	44	37	4	*	39	37
PCP	19	*	4	0	15	*
Ecstasy	291	226	7	1	280	225
Inhalants	145	102	6	1	137	101
Nonmedical Use of Psychotherapeutics ^{3,4}	2,269	2,162	27	33	2,218	2,116
Pain Relievers	1,657	1,473	22	19	1,612	1,442
OxyContin [®]	171 ^b	93	3	3	166 ^b	88
Tranquilizers	710	737	13	7	690	728
Stimulants ⁴	460	383	7	9	444	374
Methamphetamine ⁴	144	145	5	6	138	138
Sedatives	75	72	4	0	70	72
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,858	2,599	37	38	2,793	2,547

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.71B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
ILLCIT DRUGS²	10.7	10.5	4.4	5.9	10.9	10.7
Marijuana and Hashish	8.0	8.2	3.6	5.2	8.2	8.3
Cocaine	0.6	0.6	0.1	0.2	0.6	0.6
Crack	0.2 ^a	0.1	0.1	*	0.2 ^a	0.1
Heroin	0.1	0.1	0.2	0.2	0.1	0.1
Hallucinogens	0.7 ^b	0.5	0.3	0.1	0.7 ^a	0.5
LSD	0.1	0.1	0.2	*	0.1	0.1
PCP	0.0	*	0.2	0.0	0.0	*
Ecstasy	0.5 ^a	0.4	0.3	0.0	0.5	0.4
Inhalants	0.2	0.2	0.2	0.0	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	3.7	3.5	1.1	1.5	3.8	3.5
Pain Relievers	2.7 ^a	2.4	0.9	0.9	2.7 ^a	2.4
OxyContin [®]	0.3 ^b	0.1	0.1	0.1	0.3 ^b	0.1
Tranquilizers	1.2	1.2	0.5	0.3	1.2	1.2
Stimulants ⁴	0.7	0.6	0.3	0.4	0.8	0.6
Methamphetamine ⁴	0.2	0.2	0.2	0.3	0.2	0.2
Sedatives	0.1	0.1	0.2	0.0	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	4.7 ^a	4.2	1.5	1.7	4.8 ^a	4.2

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.72A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	6,548	6,574	108	130	6,394	6,415
AGE						
15-17	849	869	12	11	824	852
18-25	2,811	2,961	69	74	2,723	2,866
26-44	2,888	2,743	27	45	2,846	2,697
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,671	5,503	88	105	5,543	5,379
White	4,409 ^a	4,045	58	68	4,328 ^a	3,962
Black or African American	981	1,010	25	33	940	972
American Indian or Alaska Native	45	56	*	*	42	55
Native Hawaiian or Other Pacific Islander	11 ^a	33	*	*	10 ^a	32
Asian	100 ^a	182	*	*	100 ^a	182
Two or More Races	126 ^a	178	*	*	123 ^a	175
Hispanic or Latino	876 ^a	1,071	20	25	851 ^a	1,037
TRIMESTER²						
First	N/A	N/A	60	82	N/A	N/A
Second	N/A	N/A	32	31	N/A	N/A
Third	N/A	N/A	16	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.72B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	10.7	10.5	4.4	5.9	10.9	10.7
AGE						
15-17	13.8	13.9	15.7	18.3	13.7	13.8
18-25	16.8	17.2	7.4	9.0	17.3	17.6
26-44	7.5	7.0	1.9	3.4	7.7	7.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	11.2	10.9	4.4	6.0	11.5	11.1
White	11.9	11.2	3.8	5.5	12.2	11.4
Black or African American	11.2	11.8	8.4	9.6	11.2	11.8
American Indian or Alaska Native	13.4	13.7	*	*	13.1	14.1
Native Hawaiian or Other Pacific Islander	4.2 ^a	12.6	*	*	4.1 ^a	13.1
Asian	3.1	4.8	*	*	3.2	4.9
Two or More Races	15.0	15.6	*	*	15.4	16.0
Hispanic or Latino	8.1	8.8	4.6	5.6	8.2	8.9
TRIMESTER²						
First	N/A	N/A	8.1	11.6	N/A	N/A
Second	N/A	N/A	3.5	4.0	N/A	N/A
Third	N/A	N/A	2.0	2.3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.73A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	4,910	5,122	89	115	4,792	4,985
AGE						
15-17	648	709	11	10	628	694
18-25	2,300 ^a	2,485	61	62	2,226 ^a	2,406
26-44	1,963	1,928	17	43	1,938	1,884
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,321	4,372	71	97	4,223	4,261
White	3,333	3,217	47	64	3,268	3,144
Black or African American	781	804	21	30	752	770
American Indian or Alaska Native	32	39	*	*	30	38
Native Hawaiian or Other Pacific Islander	9 ^a	25	*	*	9 ^a	25
Asian	56 ^b	132	*	*	56 ^b	132
Two or More Races	109 ^a	154	*	*	107 ^a	152
Hispanic or Latino	589 ^b	750	18	18	568 ^b	724
TRIMESTER²						
First	N/A	N/A	48	76	N/A	N/A
Second	N/A	N/A	26	23	N/A	N/A
Third	N/A	N/A	14	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.73B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	8.0	8.2	3.6	5.2	8.2	8.3
AGE						
15-17	10.5	11.3	14.2	16.5	10.4	11.3
18-25	13.7	14.5	6.6	7.5	14.1	14.8
26-44	5.1	4.9	1.2	3.3	5.3	5.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.5	8.7	3.5	5.5	8.7	8.8
White	9.0	8.9	3.1	5.2	9.2	9.1
Black or African American	8.9	9.4	6.9	8.7	9.0	9.4
American Indian or Alaska Native	9.5	9.6	*	*	9.4	9.9
Native Hawaiian or Other Pacific Islander	3.6 ^a	9.7	*	*	3.5 ^a	10.0
Asian	1.7 ^a	3.4	*	*	1.8 ^a	3.6
Two or More Races	13.0	13.5	*	*	13.5	13.8
Hispanic or Latino	5.4	6.2	4.0	3.9	5.5	6.2
TRIMESTER²						
First	N/A	N/A	6.6	10.7	N/A	N/A
Second	N/A	N/A	2.9	2.8	N/A	N/A
Third	N/A	N/A	1.7	2.3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.74A Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOBACCO PRODUCTS²	17,017 ^a	16,173	407	363	16,526 ^a	15,747
Cigarettes	16,188 ^b	15,148	398	352	15,715 ^b	14,739
Smokeless Tobacco	321	279	8	11	309	267
Cigars	2,120	2,104	34	33	2,074	2,057
Pipe Tobacco	282	339	6	6	271	330
ALCOHOL	32,543	33,625	261 ^a	187	32,119 ^a	33,310
Binge Alcohol Use ³	14,600	14,913	89	59	14,444	14,798
Heavy Alcohol Use ³	3,229	3,171	23 ^a	6	3,194	3,150

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.74B Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOBACCO PRODUCTS²	27.7 ^b	25.9	16.6	16.4	28.2 ^b	26.2
Cigarettes	26.4 ^b	24.2	16.2	15.9	26.8 ^b	24.6
Smokeless Tobacco	0.5	0.4	0.3	0.5	0.5	0.4
Cigars	3.5	3.4	1.4	1.5	3.5	3.4
Pipe Tobacco	0.5	0.5	0.2	0.2	0.5	0.5
ALCOHOL	53.0	53.8	10.6	8.5	54.8	55.5
Binge Alcohol Use ³	23.8	23.9	3.6	2.7	24.6	24.7
Heavy Alcohol Use ³	5.3	5.1	0.9 ^a	0.3	5.4	5.2

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.75A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	16,188 ^b	15,148	398	352	15,715 ^b	14,739
AGE						
15-17	842 ^b	691	17	*	811 ^b	670
18-25	5,167 ^b	4,786	211	172	4,924 ^b	4,591
26-44	10,178	9,671	170	166	9,980	9,479
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	14,220 ^b	13,234	372	335	13,778 ^b	12,847
White	11,653 ^b	10,637	313	251	11,282 ^b	10,344
Black or African American	1,802	1,618	41	54	1,752	1,558
American Indian or Alaska Native	134 ^a	188	*	*	129	177
Native Hawaiian or Other Pacific Islander	34	*	*	*	30	*
Asian	318	335	*	*	318	335
Two or More Races	279 ^a	379	*	*	266 ^a	359
Hispanic or Latino	1,969	1,913	26	17	1,937	1,893
TRIMESTER²						
First	N/A	N/A	177	164	N/A	N/A
Second	N/A	N/A	124	108	N/A	N/A
Third	N/A	N/A	94	78	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.75B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	26.4 ^b	24.2	16.2	15.9	26.8 ^b	24.6
AGE						
15-17	13.7 ^b	11.0	22.7	*	13.4 ^b	10.9
18-25	30.9 ^b	27.8	22.7	20.9	31.3 ^b	28.2
26-44	26.4 ^a	24.8	11.7	12.5	27.1 ^a	25.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	28.1 ^b	26.3	18.5	19.1	28.5 ^b	26.5
White	31.4 ^b	29.4	20.7	20.3	31.8 ^b	29.8
Black or African American	20.6	18.9	13.4	15.8	20.9	19.0
American Indian or Alaska Native	40.1	46.3	*	*	40.7	45.5
Native Hawaiian or Other Pacific Islander	13.2	*	*	*	12.0	*
Asian	9.7	8.7	*	*	10.2	9.0
Two or More Races	33.2	33.2	*	*	33.5	32.7
Hispanic or Latino	18.2 ^a	15.8	5.8	3.8	18.8 ^a	16.3
TRIMESTER²						
First	N/A	N/A	24.2	23.2	N/A	N/A
Second	N/A	N/A	13.6	13.6	N/A	N/A
Third	N/A	N/A	12.0	11.1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.76A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	32,543	33,625	261 ^a	187	32,119 ^a	33,310
AGE						
15-17	1,358	1,348	10	8	1,336	1,331
18-25	9,592 ^b	9,934	85	60	9,458	9,827
26-44	21,593	22,344	166	119	21,325	22,152
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	28,115	28,413	217	162	27,763	28,136
White	22,158	22,000	165	121	21,892	21,788
Black or African American	4,068	4,065	38	31	3,996	4,018
American Indian or Alaska Native	140 ^a	197	*	*	140 ^a	196
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1,194	1,374	*	*	1,184	1,371
Two or More Races	450 ^b	672	*	*	446 ^b	658
Hispanic or Latino	4,428 ^b	5,212	44	25	4,356 ^b	5,174
TRIMESTER²						
First	N/A	N/A	145	127	N/A	N/A
Second	N/A	N/A	86 ^b	33	N/A	N/A
Third	N/A	N/A	24	26	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.76B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	53.0	53.8	10.6	8.5	54.8	55.5
AGE						
15-17	22.1	21.5	12.7	13.4	22.2	21.6
18-25	57.3	57.8	9.2	7.3	60.1	60.4
26-44	56.1	57.2	11.5	9.0	57.8	58.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	55.6	56.4	10.8	9.2	57.5	58.1
White	59.7	60.9	11.0	9.7	61.7	62.7
Black or African American	46.5	47.4	12.5	9.2	47.7	49.0
American Indian or Alaska Native	42.1	48.7	*	*	44.0	50.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	36.6	35.8	*	*	38.2	36.9
Two or More Races	53.7	58.9	*	*	56.1	60.0
Hispanic or Latino	40.8	43.0	9.9	5.5	42.2	44.5
TRIMESTER²						
First	N/A	N/A	19.8	17.9	N/A	N/A
Second	N/A	N/A	9.4 ^b	4.2	N/A	N/A
Third	N/A	N/A	3.0	3.7	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.77A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	14,600	14,913	89	59	14,444	14,798
AGE						
15-17	781	718	8	*	764	710
18-25	5,559	5,707	39	25	5,483	5,658
26-44	8,261	8,488	42	30	8,197	8,431
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12,340	12,397	70	52	12,220	12,295
White	10,043	9,817	55	35	9,955	9,737
Black or African American	1,606	1,665	14	10	1,576	1,650
American Indian or Alaska Native	88	123	*	*	88	122
Native Hawaiian or Other Pacific Islander	44	59	*	*	44	*
Asian	352	459	*	*	352	458
Two or More Races	206 ^a	274	*	*	205	268
Hispanic or Latino	2,261	2,516	19	7	2,225 ^a	2,503
TRIMESTER²						
First	N/A	N/A	73	47	N/A	N/A
Second	N/A	N/A	12	9	N/A	N/A
Third	N/A	N/A	0	3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.77B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	23.8	23.9	3.6	2.7	24.6	24.7
AGE						
15-17	12.7 ^a	11.5	10.2	*	12.7 ^a	11.5
18-25	33.2	33.2	4.2	3.0	34.9	34.8
26-44	21.4	21.7	2.9	2.3	22.2	22.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	24.4	24.6	3.5	3.0	25.3	25.4
White	27.0	27.2	3.6	2.8	28.1	28.0
Black or African American	18.4	19.4	4.5	3.1	18.8	20.1
American Indian or Alaska Native	26.3	30.3	*	*	27.6	31.3
Native Hawaiian or Other Pacific Islander	17.2	22.6	*	*	17.4	*
Asian	10.8	12.0	*	*	11.4	12.3
Two or More Races	24.6	24.0	*	*	25.8	24.5
Hispanic or Latino	20.8	20.7	4.3	1.5	21.6	21.5
TRIMESTER²						
First	N/A	N/A	10.0	6.6	N/A	N/A
Second	N/A	N/A	1.3	1.1	N/A	N/A
Third	N/A	N/A	0.1	0.4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.78A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	3,229	3,171	23 ^a	6	3,194	3,150
AGE						
15-17	171 ^b	130	4	*	164 ^a	130
18-25	1,423	1,360	14 ^a	4	1,400	1,349
26-44	1,635	1,680	5	2	1,630	1,671
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,873	2,786	17 ^a	5	2,846	2,768
White	2,467	2,304	13	4	2,446	2,287
Black or African American	290	278	4	0	285	277
American Indian or Alaska Native	23	44	*	*	23	44
Native Hawaiian or Other Pacific Islander	9	*	*	*	9	*
Asian	45	75	*	*	45	75
Two or More Races	39	59	*	*	38	59
Hispanic or Latino	356	385	6	1	348	382
TRIMESTER²						
First	N/A	N/A	19 ^a	5	N/A	N/A
Second	N/A	N/A	4	0	N/A	N/A
Third	N/A	N/A	*	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.78B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	5.3	5.1	0.9 ^a	0.3	5.4	5.2
AGE						
15-17	2.8 ^b	2.1	5.4	*	2.7 ^a	2.1
18-25	8.5	7.9	1.5 ^a	0.4	8.9	8.3
26-44	4.2	4.3	0.4	0.2	4.4	4.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5.7	5.5	0.8 ^a	0.3	5.9	5.7
White	6.6	6.4	0.8	0.3	6.9	6.6
Black or African American	3.3	3.2	1.3	0.1	3.4	3.4
American Indian or Alaska Native	7.0	10.9	*	*	7.4	11.3
Native Hawaiian or Other Pacific Islander	3.6	*	*	*	3.6	*
Asian	1.4	1.9	*	*	1.4	2.0
Two or More Races	4.6	5.2	*	*	4.8	5.3
Hispanic or Latino	3.3	3.2	1.4	0.3	3.4	3.3
TRIMESTER²						
First	N/A	N/A	2.6 ^a	0.8	N/A	N/A
Second	N/A	N/A	0.4	0.1	N/A	N/A
Third	N/A	N/A	*	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2011-2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011-2012 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.