

Table 6.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	214	233	228	270	222	257	225	217
Marijuana and Hashish	182	209	213	239	195	224	183	172
Cocaine	78	88	59	84	32	57	27	48
Crack	26	63	31	58	3	9	10	38
Heroin	20	22	17	19	16	22	32	34
Hallucinogens	33	58	35	63	27	27	22	21
LSD	10	26	11	14	6	5	16	6
PCP	3	11	9	4	*	3	*	*
Ecstasy	26	45	26	50	23	25	10	12
Inhalants	33	22	27	53	30	22	38	19
Nonmedical Use of Psychotherapeutics ^{3,4}	104	110	105	135	113	130	131	133
Pain Relievers	76	92	95	105	89	97	116	115
OxyContin [®]	26	29	31	16	19	13	27	27
Tranquilizers	64	67	59	89	58	84	74	81
Stimulants ⁴	51	45	36	62	28	54	53	39
Methamphetamine ⁴	42	25	17	53	23	28	50	33
Sedatives	23	12	13	36	21	24	35	21
ILLICIT DRUGS OTHER THAN MARIJUANA²	128	142	121	164	122	142	142	142

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	1.06	1.12	0.51	0.57	0.30	0.33	0.18	0.17
Marijuana and Hashish	0.96	1.05	0.47	0.52	0.26	0.29	0.15	0.14
Cocaine	0.47	0.51	0.14	0.19	0.04	0.08	0.02	0.04
Crack	0.17	0.37	0.07	0.14	0.00	0.01	0.01	0.03
Heroin	0.13	0.13	0.04	0.05	0.02	0.03	0.03	0.03
Hallucinogens	0.21	0.34	0.08	0.15	0.04	0.04	0.02	0.02
LSD	0.07	0.15	0.03	0.03	0.01	0.01	0.01	0.00
PCP	0.02	0.07	0.02	0.01	*	0.00	*	*
Ecstasy	0.17	0.27	0.06	0.12	0.03	0.03	0.01	0.01
Inhalants	0.21	0.13	0.06	0.12	0.04	0.03	0.03	0.02
Nonmedical Use of Psychotherapeutics ^{3,4}	0.62	0.61	0.25	0.31	0.15	0.17	0.11	0.11
Pain Relievers	0.47	0.52	0.22	0.24	0.12	0.13	0.09	0.09
OxyContin [®]	0.16	0.17	0.07	0.04	0.03	0.02	0.02	0.02
Tranquilizers	0.40	0.39	0.14	0.21	0.08	0.11	0.06	0.07
Stimulants ⁴	0.32	0.26	0.09	0.14	0.04	0.07	0.04	0.03
Methamphetamine ⁴	0.26	0.15	0.04	0.12	0.03	0.04	0.04	0.03
Sedatives	0.14	0.07	0.03	0.08	0.03	0.03	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.73	0.77	0.29	0.37	0.16	0.19	0.11	0.11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	0.8722		0.0798		0.3180		0.8033	
Marijuana and Hashish	0.9202		0.2924		0.7251		0.8341	
Cocaine	0.8997		0.4278		0.2662		0.0970	
Crack	0.2666		0.6121		0.3220		0.0241	
Heroin	0.8393		0.7175		0.9903		0.4742	
Hallucinogens	0.0519		0.4251		0.1203		0.8580	
LSD	0.5133		0.2691		0.8085		0.5955	
PCP	0.3806		0.3355		*		*	
Ecstasy	0.1005		0.6943		0.2294		0.9965	
Inhalants	0.1172		0.4718		0.2510		0.0324	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.7650		0.0267		0.2034		0.8590	
Pain Relievers	0.8934		0.0713		0.5213		0.9092	
OxyContin [®]	0.8088		0.0503		0.7129		0.9321	
Tranquilizers	0.6561		0.2089		0.4080		0.4516	
Stimulants ⁴	0.6541		0.0382		0.1384		0.5599	
Methamphetamine ⁴	0.1242		0.0716		0.6955		0.4113	
Sedatives	0.5329		0.0243		0.8828		0.3476	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.9022		0.0202		0.2583		0.9419	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	20	21	32	32	30	28	50	53
Marijuana and Hashish	21	19	30	30	29	26	41	40
Cocaine	6	3	6	6	6	2	4	3
Crack	0	0	2	1	*	*	*	*
Heroin	1	*	2	*	7	*	1	*
Hallucinogens	9	9	12	10	11	5	12	11
LSD	3	3	7	3	1	3	3	3
PCP	1	*	1	1	*	2	*	2
Ecstasy	7	6	8	7	9	4	6	6
Inhalants	4	9	9	8	6	7	16	14
Nonmedical Use of Psychotherapeutics ^{3,4}	12	10	16	17	11	12	25	31
Pain Relievers	10	8	15	15	9	11	23	26
OxyContin [®]	5	3	8	8	1	3	3	3
Tranquilizers	9	7	11	8	6	4	7	16
Stimulants ⁴	6	6	5	9	6	4	7	8
Methamphetamine ⁴	4	2	4	4	5	2	4	4
Sedatives	2	2	4	4	3	0	3	5
ILLICIT DRUGS OTHER THAN MARIJUANA²	13	14	21	20	17	14	31	35

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	3.04	3.64	1.64	1.72	1.70	1.60	0.22	0.23
Marijuana and Hashish	3.08	3.91	1.63	1.64	1.64	1.55	0.19	0.18
Cocaine	1.52	0.77	0.40	0.37	0.42	0.13	0.02	0.01
Crack	0.03	0.05	0.14	0.04	*	*	*	*
Heroin	0.22	*	0.12	*	0.49	*	0.00	*
Hallucinogens	2.14	2.65	0.77	0.63	0.76	0.32	0.05	0.05
LSD	0.86	0.95	0.45	0.19	0.05	0.20	0.01	0.01
PCP	0.37	*	0.08	0.04	*	0.16	*	0.01
Ecstasy	1.86	1.74	0.54	0.46	0.60	0.28	0.03	0.03
Inhalants	0.93	2.46	0.58	0.51	0.43	0.49	0.08	0.06
Nonmedical Use of Psychotherapeutics ^{3,4}	2.84	2.75	1.05	1.10	0.74	0.79	0.11	0.14
Pain Relievers	2.54	2.33	0.96	0.99	0.61	0.73	0.11	0.12
OxyContin [®]	1.20	0.83	0.50	0.56	0.07	0.24	0.02	0.02
Tranquilizers	2.38	2.20	0.72	0.52	0.40	0.25	0.03	0.07
Stimulants ⁴	1.61	1.78	0.34	0.62	0.41	0.30	0.03	0.04
Methamphetamine ⁴	1.03	0.68	0.26	0.28	0.33	0.14	0.02	0.02
Sedatives	0.42	0.47	0.27	0.25	0.18	0.02	0.01	0.02
ILLICIT DRUGS OTHER THAN MARIJUANA²	3.14	3.37	1.28	1.22	1.07	0.91	0.14	0.16

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	0.6027		0.2808		0.6089		0.6336	
Marijuana and Hashish	0.2395		0.3289		0.2677		0.5692	
Cocaine	0.0181		0.4601		0.0823		0.4848	
Crack	0.8493		0.3117		*		*	
Heroin	*		*		*		*	
Hallucinogens	0.5351		0.1148		0.0492		0.5956	
LSD	0.9398		0.0913		0.1579		0.6395	
PCP	*		0.4711		*		*	
Ecstasy	0.5191		0.3122		0.1635		0.9079	
Inhalants	0.0908		0.4567		0.4123		0.0305	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.3792		0.5462		0.9168		0.2144	
Pain Relievers	0.0516		0.7123		0.4115		0.4434	
OxyContin [®]	0.0588		0.5486		0.4409		0.9687	
Tranquilizers	0.3774		0.1320		0.1021		0.1587	
Stimulants ⁴	0.6095		0.0291		0.1789		0.1555	
Methamphetamine ⁴	0.2746		0.6102		0.3494		0.7134	
Sedatives	0.7164		0.8853		0.0570		0.2380	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.9204		0.0537		0.3947		0.6038	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	97	84	101	94	67	69	57	65
Marijuana and Hashish	86	81	96	90	62	66	50	58
Cocaine	33	24	24	19	8	7	7	11
Crack	3	7	5	4	1	2	1	4
Heroin	7	7	11	11	11	6	8	17
Hallucinogens	22	33	28	25	15	11	12	13
LSD	10	8	9	13	3	4	2	5
PCP	1	3	6	2	1	2	*	*
Ecstasy	17	31	21	18	11	9	8	8
Inhalants	18	14	14	12	17	5	5	11
Nonmedical Use of Psychotherapeutics ^{3,4}	41	46	40	39	30	26	27	34
Pain Relievers	32	36	36	33	25	23	22	33
OxyContin [®]	13	10	13	10	8	11	6	16
Tranquilizers	22	24	22	22	17	13	16	23
Stimulants ⁴	24	22	14	19	11	6	10	9
Methamphetamine ⁴	8	10	6	10	6	2	9	5
Sedatives	6	9	5	7	6	3	4	4
ILLICIT DRUGS OTHER THAN MARIJUANA²	55	55	52	47	37	28	31	38

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	1.33	1.28	0.81	0.77	0.81	0.84	0.39	0.44
Marijuana and Hashish	1.31	1.25	0.77	0.77	0.76	0.80	0.36	0.41
Cocaine	0.72	0.52	0.25	0.20	0.12	0.10	0.05	0.08
Crack	0.08	0.15	0.05	0.04	0.01	0.03	0.01	0.03
Heroin	0.18	0.16	0.11	0.11	0.15	0.09	0.06	0.12
Hallucinogens	0.50	0.74	0.29	0.26	0.21	0.15	0.09	0.09
LSD	0.23	0.18	0.09	0.14	0.05	0.06	0.02	0.04
PCP	0.03	0.07	0.06	0.02	0.01	0.03	*	*
Ecstasy	0.40	0.70	0.22	0.20	0.15	0.12	0.06	0.05
Inhalants	0.43	0.32	0.14	0.13	0.24	0.07	0.03	0.08
Nonmedical Use of Psychotherapeutics ^{3,4}	0.85	0.95	0.40	0.40	0.42	0.36	0.20	0.24
Pain Relievers	0.71	0.78	0.36	0.35	0.34	0.31	0.16	0.24
OxyContin [®]	0.30	0.23	0.14	0.11	0.11	0.15	0.04	0.11
Tranquilizers	0.51	0.54	0.22	0.23	0.24	0.18	0.12	0.17
Stimulants ⁴	0.55	0.48	0.14	0.20	0.15	0.09	0.08	0.06
Methamphetamine ⁴	0.19	0.23	0.06	0.10	0.09	0.03	0.07	0.04
Sedatives	0.13	0.20	0.05	0.07	0.09	0.05	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.03	1.10	0.50	0.47	0.50	0.39	0.22	0.27

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	0.4123		0.3539		0.2812		0.0091	
Marijuana and Hashish	0.6401		0.1822		0.4402		0.0764	
Cocaine	0.1681		0.3525		0.6439		0.4425	
Crack	0.1325		0.4580		0.6202		0.0331	
Heroin	0.9691		0.8147		0.2235		0.0212	
Hallucinogens	0.1077		0.8726		0.0740		0.9636	
LSD	0.4138		0.0722		0.8272		0.2632	
PCP	0.2095		0.3634		0.6355		*	
Ecstasy	0.0352		0.8751		0.1622		0.3690	
Inhalants	0.6409		0.8857		0.1631		0.2664	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.9015		0.1648		0.3448		0.1225	
Pain Relievers	0.6147		0.7648		0.6065		0.0109	
OxyContin [®]	0.6736		0.1630		0.9776		0.3523	
Tranquilizers	0.4226		0.4145		0.6239		0.8532	
Stimulants ⁴	0.7108		0.0384		0.6373		0.7368	
Methamphetamine ⁴	0.4534		0.1774		0.7211		0.3285	
Sedatives	0.7183		0.1347		0.3985		0.7917	
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.5789		0.4278		0.0961		0.1007	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS²	186	215	202	246	215	247	208	195
Marijuana and Hashish	165	192	186	216	187	213	170	153
Cocaine	69	85	54	81	31	57	25	46
Crack	26	62	30	58	3	9	10	38
Heroin	19	21	13	16	9	22	31	29
Hallucinogens	23	47	19	56	19	25	15	13
LSD	1	25	*	6	5	*	15	*
PCP	3	11	7	4	*	*	*	*
Ecstasy	19	32	14	46	18	23	*	7
Inhalants	28	15	21	51	24	18	34	7
Nonmedical Use of Psychotherapeutics ^{3,4}	92	101	99	127	109	126	125	123
Pain Relievers	67	86	88	100	85	93	111	105
OxyContin [®]	22	27	27	10	17	6	26	22
Tranquilizers	59	60	53	87	55	83	73	76
Stimulants ⁴	45	37	34	59	25	54	52	37
Methamphetamine ⁴	41	21	16	52	22	28	49	33
Sedatives	22	8	11	35	20	23	35	20
ILLCIT DRUGS OTHER THAN MARIJUANA²	110	131	109	153	116	138	132	129

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	1.35	1.47	0.61	0.71	0.32	0.36	0.23	0.22
Marijuana and Hashish	1.25	1.36	0.57	0.64	0.28	0.31	0.19	0.17
Cocaine	0.59	0.68	0.17	0.25	0.05	0.08	0.03	0.05
Crack	0.23	0.50	0.10	0.18	0.00	0.01	0.01	0.04
Heroin	0.17	0.17	0.04	0.05	0.01	0.03	0.04	0.03
Hallucinogens	0.20	0.38	0.06	0.17	0.03	0.04	0.02	0.01
LSD	0.01	0.20	*	0.02	0.01	*	0.02	*
PCP	0.03	0.09	0.02	0.01	*	*	*	*
Ecstasy	0.16	0.26	0.04	0.15	0.03	0.03	*	0.01
Inhalants	0.24	0.12	0.07	0.16	0.04	0.03	0.04	0.01
Nonmedical Use of Psychotherapeutics ^{3,4}	0.78	0.78	0.31	0.39	0.16	0.18	0.14	0.14
Pain Relievers	0.58	0.68	0.28	0.31	0.13	0.14	0.12	0.12
OxyContin [®]	0.20	0.22	0.09	0.03	0.03	0.01	0.03	0.02
Tranquilizers	0.51	0.48	0.17	0.27	0.08	0.12	0.08	0.09
Stimulants ⁴	0.39	0.29	0.11	0.18	0.04	0.08	0.06	0.04
Methamphetamine ⁴	0.35	0.17	0.05	0.16	0.03	0.04	0.06	0.04
Sedatives	0.20	0.06	0.04	0.11	0.03	0.03	0.04	0.02
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.90	0.99	0.35	0.46	0.17	0.20	0.15	0.14

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLICIT DRUGS²	0.7198		0.0120		0.1452		0.6822	
Marijuana and Hashish	0.5428		0.0538		0.4228		0.4754	
Cocaine	0.6010		0.2380		0.1675		0.1120	
Crack	0.3486		0.5442		0.3691		0.0416	
Heroin	0.8050		0.6841		0.2698		0.8603	
Hallucinogens	0.1510		0.1700		0.9474		0.4876	
LSD	0.2028		*		*		*	
PCP	0.4928		0.6849		*		*	
Ecstasy	0.5142		0.3999		0.8124		*	
Inhalants	0.0202		0.3202		0.4127		0.0895	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.9572		0.0475		0.1145		0.6274	
Pain Relievers	0.6933		0.0583		0.4575		0.3244	
OxyContin [®]	0.7947		0.0718		0.5645		0.7182	
Tranquilizers	0.9882		0.1972		0.2754		0.5560	
Stimulants ⁴	0.5976		0.2364		0.0659		0.4443	
Methamphetamine ⁴	0.0923		0.1294		0.6249		0.4922	
Sedatives	0.3988		0.0478		0.6064		0.2845	
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.6203		0.0115		0.0844		0.6825	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS¹	412	460	136	122	388	444
Marijuana or Hashish	367	406	127	114	347	389
Cocaine (Including Crack)	103	135	30	22	98	134
Heroin	35	37	14	8	31	36
Hallucinogens	56	88	31	35	45	81
Inhalants	57	62	28	22	45	57
Pain Relievers ²	148	177	40	40	142	174
Tranquilizers ²	109	141	29	24	105	139
Stimulants ^{2,3}	70	93	20	21	66	91
Methamphetamine ³	50	66	10	9	50	64
Sedatives ²	34	45	7	7	33	44
ILLCIT DRUGS OTHER THAN MARIJUANA¹	211	273	66	61	197	265

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS¹	0.29	0.32	0.90	0.83	0.29	0.33
Marijuana or Hashish	0.26	0.28	0.87	0.80	0.26	0.29
Cocaine (Including Crack)	0.08	0.10	0.29	0.23	0.08	0.11
Heroin	0.03	0.03	0.14	0.09	0.03	0.03
Hallucinogens	0.04	0.07	0.32	0.37	0.04	0.06
Inhalants	0.04	0.05	0.29	0.23	0.04	0.05
Pain Relievers ²	0.11	0.13	0.40	0.43	0.12	0.14
Tranquilizers ²	0.08	0.10	0.30	0.26	0.08	0.11
Stimulants ^{2,3}	0.05	0.07	0.20	0.22	0.05	0.07
Methamphetamine ³	0.04	0.05	0.10	0.10	0.04	0.05
Sedatives ²	0.03	0.03	0.08	0.08	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.16	0.20	0.60	0.61	0.16	0.21

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS¹	0.0670		0.1654		0.0135	
Marijuana or Hashish	0.2448		0.0607		0.0547	
Cocaine (Including Crack)	0.3142		0.0049		0.1061	
Heroin	0.7332		0.4542		0.5162	
Hallucinogens	0.1772		0.6481		0.1375	
Inhalants	0.5539		0.9290		0.5834	
Pain Relievers ²	0.1247		0.7419		0.0785	
Tranquilizers ²	0.2763		0.0266		0.0962	
Stimulants ^{2,3}	0.0497		0.5643		0.0521	
Methamphetamine ³	0.5965		0.8340		0.5959	
Sedatives ²	0.1780		0.9223		0.1587	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0214		0.2863		0.0054	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS²	261	302	82	84	243	290
Marijuana or Hashish	246	285	76	83	230	274
Cocaine (Including Crack)	48	100	14	13	46	99
Heroin	15	24	7	4	13	24
Hallucinogens	22	27	8	8	20	26
Inhalants	22	6	16	5	15	*
Pain Relievers ³	54	73	14	11	54	72
Tranquilizers ³	19	38	8	5	17	38
Stimulants ³	24	37	2	3	24	37
Methamphetamine ⁴	31	58	4	6	31	57
Sedatives ³	21	12	3	1	21	11
ILLCIT DRUGS OTHER THAN MARIJUANA²	89	132	30	21	85	130

*Low precision; no estimate reported.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS²	0.20	0.21	0.70	0.74	0.20	0.22
Marijuana or Hashish	0.18	0.20	0.67	0.73	0.19	0.21
Cocaine (Including Crack)	0.04	0.07	0.14	0.14	0.04	0.08
Heroin	0.01	0.02	0.07	0.04	0.01	0.02
Hallucinogens	0.02	0.02	0.08	0.09	0.02	0.02
Inhalants	0.02	0.00	0.16	0.06	0.01	*
Pain Relievers ³	0.04	0.05	0.14	0.12	0.04	0.06
Tranquilizers ³	0.01	0.03	0.08	0.06	0.01	0.03
Stimulants ³	0.02	0.03	0.02	0.04	0.02	0.03
Methamphetamine ⁴	0.02	0.04	0.04	0.06	0.02	0.04
Sedatives ³	0.02	0.01	0.03	0.01	0.02	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.07	0.10	0.30	0.23	0.07	0.10

*Low precision; no estimate reported.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug¹	Total (2011)	Total (2012)	Aged 12-20 (2011)	Aged 12-20 (2012)	Aged 21+ (2011)	Aged 21+ (2012)
ILLCIT DRUGS²	0.0039		0.6000		0.0020	
Marijuana or Hashish	0.0080		0.5605		0.0048	
Cocaine (Including Crack)	0.0499		0.4270		0.0330	
Heroin	0.9920		0.3148		0.7551	
Hallucinogens	0.7538		0.7068		0.8067	
Inhalants	0.3113		0.7548		*	
Pain Relievers ³	0.2863		0.3268		0.2032	
Tranquilizers ³	0.1843		0.3471		0.1106	
Stimulants ³	0.4587		0.1759		0.5322	
Methamphetamine ⁴	0.1210		0.3105		0.1423	
Sedatives ³	0.3950		0.6153		0.4319	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.0633		0.2220		0.0265	

*Low precision; no estimate reported.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.17C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS¹	412	460	50	48	179	163	365	425
Marijuana or Hashish	367	406	48	45	165	156	331	371
Cocaine (Including Crack)	103	135	11	6	42	31	93	130
Heroin	35	37	7	*	18	14	28	34
Hallucinogens	56	88	19	14	39	45	35	76
Inhalants	57	62	11	13	36	19	43	56
Pain Relievers ²	148	177	21	21	56	55	136	170
Tranquilizers ²	109	141	15	11	36	36	100	136
Stimulants ^{2,3}	70	93	10	12	32	31	61	88
Methamphetamine ³	50	66	7	5	12	14	49	63
Sedatives ²	34	45	5	4	10	12	32	43
ILLCIT DRUGS OTHER THAN MARIJUANA¹	211	273	30	28	90	82	188	253

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.17D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2011 and 2012

Drug	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS¹	0.29	0.32	1.19	1.20	0.60	0.54	0.32	0.36
Marijuana or Hashish	0.26	0.28	1.16	1.17	0.57	0.54	0.29	0.32
Cocaine (Including Crack)	0.08	0.10	0.32	0.19	0.19	0.14	0.08	0.12
Heroin	0.03	0.03	0.22	*	0.09	0.07	0.03	0.03
Hallucinogens	0.04	0.07	0.54	0.42	0.18	0.21	0.03	0.07
Inhalants	0.04	0.05	0.33	0.41	0.17	0.09	0.04	0.05
Pain Relievers ²	0.11	0.13	0.61	0.61	0.25	0.26	0.13	0.15
Tranquilizers ²	0.08	0.10	0.46	0.34	0.17	0.17	0.09	0.12
Stimulants ^{2,3}	0.05	0.07	0.30	0.37	0.15	0.14	0.06	0.08
Methamphetamine ³	0.04	0.05	0.22	0.16	0.06	0.07	0.04	0.06
Sedatives ²	0.03	0.03	0.15	0.12	0.05	0.06	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.16	0.20	0.85	0.79	0.38	0.35	0.17	0.22

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.17P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS¹	0.0670		0.1883		0.2058		0.0061	
Marijuana or Hashish	0.2448		0.0679		0.2178		0.0341	
Cocaine (Including Crack)	0.3142		0.0045		0.1010		0.0733	
Heroin	0.7332		*		0.6057		0.3339	
Hallucinogens	0.1772		0.0576		0.4759		0.0599	
Inhalants	0.5539		0.4050		0.3534		0.7572	
Pain Relievers ²	0.1247		0.4381		0.8361		0.0601	
Tranquilizers ²	0.2763		0.0213		0.9110		0.1382	
Stimulants ^{2,3}	0.0497		0.6601		0.1384		0.1183	
Methamphetamine ³	0.5965		0.4088		0.1278		0.7508	
Sedatives ²	0.1780		0.5849		0.4880		0.1971	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0214		0.0592		0.7292		0.0037	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS²	261	302	32	34	118	113	230	276
Marijuana or Hashish	246	285	31	33	111	110	217	260
Cocaine (Including Crack)	48	100	5	3	18	18	44	98
Heroin	15	24	4	*	10	5	10	24
Hallucinogens	22	27	4	3	15	13	15	24
Inhalants	22	6	1	3	16	4	15	*
Pain Relievers ³	54	73	8	5	19	23	52	69
Tranquilizers ³	19	38	4	2	9	12	16	36
Stimulants ³	24	37	1	3	7	7	23	36
Methamphetamine ⁴	31	58	4	2	3	11	31	56
Sedatives ³	21	12	3	1	*	2	21	11
ILLCIT DRUGS OTHER THAN MARIJUANA²	89	132	12	9	38	35	82	126

*Low precision; no estimate reported.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2011 and 2012

Drug¹	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLCIT DRUGS²	0.20	0.21	0.90	0.99	0.47	0.45	0.21	0.24
Marijuana or Hashish	0.18	0.20	0.88	0.97	0.46	0.45	0.20	0.23
Cocaine (Including Crack)	0.04	0.07	0.15	0.09	0.09	0.08	0.04	0.09
Heroin	0.01	0.02	0.12	*	0.05	0.02	0.01	0.02
Hallucinogens	0.02	0.02	0.11	0.11	0.07	0.06	0.01	0.02
Inhalants	0.02	0.00	0.04	0.11	0.08	0.02	0.01	*
Pain Relievers ³	0.04	0.05	0.25	0.17	0.09	0.11	0.05	0.06
Tranquilizers ³	0.01	0.03	0.13	0.08	0.04	0.06	0.01	0.03
Stimulants ³	0.02	0.03	0.04	0.09	0.03	0.04	0.02	0.03
Methamphetamine ⁴	0.02	0.04	0.13	0.06	0.01	0.05	0.03	0.05
Sedatives ³	0.02	0.01	0.09	0.04	*	0.01	0.02	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.07	0.10	0.37	0.29	0.18	0.16	0.07	0.11

*Low precision; no estimate reported.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug¹	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
ILLICIT DRUGS²	0.0039		0.5878		0.8023		0.0009	
Marijuana or Hashish	0.0080		0.6030		0.7029		0.0025	
Cocaine (Including Crack)	0.0499		0.2342		0.6346		0.0285	
Heroin	0.9920		*		0.1139		0.3732	
Hallucinogens	0.7538		0.8084		0.7424		0.4788	
Inhalants	0.3113		0.2467		0.5951		*	
Pain Relievers ³	0.2863		0.4637		0.9394		0.2323	
Tranquilizers ³	0.1843		0.3712		0.8610		0.1340	
Stimulants ³	0.4587		0.2763		0.9275		0.5076	
Methamphetamine ⁴	0.1210		0.5748		0.0188		0.2208	
Sedatives ³	0.3950		0.6125		*		0.4104	
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.0633		0.3500		0.4104		0.0189	

*Low precision; no estimate reported.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.