

Table 6.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	289	311	147	150	285	322	154	150
Marijuana and Hashish	268	282	126	126	247	281	119	110
Cocaine	82	123	26	36	60	58	5	31
Crack	41	82	10	32	*	25	*	20
Heroin	30	37	31	23	8	4	8	25
Hallucinogens	49	83	18	16	26	29	14	13
LSD	15	30	15	4	8	7	4	5
PCP	10	5	*	2	*	11	*	*
Ecstasy	40	68	7	8	21	23	7	8
Inhalants	44	26	8	7	36	55	37	18
Nonmedical Use of Psychotherapeutics ^{2,3}	129	151	83	82	131	165	104	105
Pain Relievers	103	119	73	67	109	123	89	92
OxyContin [®]	37	35	19	25	25	8	19	12
Tranquilizers	78	104	44	44	73	96	60	68
Stimulants ³	60	67	29	37	34	64	44	11
Methamphetamine ³	47	58	24	32	18	31	44	7
Sedatives	30	13	17	17	16	43	31	12
ILLCIT DRUGS OTHER THAN MARIJUANA¹	151	195	87	91	146	182	110	109

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.68	0.70	0.74	0.72	0.28	0.32	0.15	0.14
Marijuana and Hashish	0.64	0.66	0.65	0.62	0.25	0.28	0.11	0.10
Cocaine	0.21	0.32	0.14	0.18	0.06	0.06	0.01	0.03
Crack	0.11	0.22	0.05	0.16	*	0.03	*	0.02
Heroin	0.08	0.10	0.16	0.11	0.01	0.00	0.01	0.02
Hallucinogens	0.13	0.22	0.09	0.08	0.03	0.03	0.01	0.01
LSD	0.04	0.08	0.08	0.02	0.01	0.01	0.00	0.00
PCP	0.03	0.01	*	0.01	*	0.01	*	*
Ecstasy	0.11	0.18	0.04	0.04	0.02	0.02	0.01	0.01
Inhalants	0.12	0.07	0.04	0.04	0.04	0.06	0.04	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.33	0.38	0.42	0.41	0.14	0.17	0.10	0.10
Pain Relievers	0.27	0.31	0.37	0.34	0.11	0.13	0.09	0.09
OxyContin [®]	0.10	0.09	0.10	0.12	0.03	0.01	0.02	0.01
Tranquilizers	0.20	0.27	0.23	0.22	0.08	0.10	0.06	0.06
Stimulants ³	0.16	0.18	0.15	0.18	0.04	0.07	0.04	0.01
Methamphetamine ³	0.12	0.15	0.13	0.16	0.02	0.03	0.04	0.01
Sedatives	0.08	0.03	0.09	0.09	0.02	0.04	0.03	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.39	0.49	0.44	0.45	0.15	0.18	0.10	0.10

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.0965		0.9226		0.1548		0.9245	
Marijuana and Hashish	0.2339		0.8261		0.3906		0.3476	
Cocaine	0.1457		0.6194		0.7444		0.0396	
Crack	0.2828		0.0623		*		*	
Heroin	0.6070		0.7063		0.6419		0.0950	
Hallucinogens	0.1383		0.9327		0.7775		0.7839	
LSD	0.2293		0.4664		1.0000		0.7255	
PCP	0.3116		*		*		*	
Ecstasy	0.4541		0.7361		0.4522		0.7931	
Inhalants	0.1394		0.6397		0.6861		0.0371	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4334		0.1846		0.0184		0.2768	
Pain Relievers	0.2295		0.1184		0.2355		0.3594	
OxyContin [®]	0.5689		0.5317		0.0618		0.4075	
Tranquilizers	0.4327		0.7689		0.3547		0.3168	
Stimulants ³	0.9743		0.6278		0.0010		0.1609	
Methamphetamine ³	0.9867		0.5776		0.1769		0.0945	
Sedatives	0.2270		0.3520		0.0107		0.6075	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.1538		0.2860		0.0282		0.5613	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	37	36	26	22	33	33	42	45
Marijuana and Hashish	36	34	25	21	31	32	30	32
Cocaine	10	6	4	3	5	1	1	1
Crack	2	1	*	*	*	0	*	*
Heroin	7	*	1	*	0	*	*	*
Hallucinogens	16	14	5	8	10	3	11	8
LSD	7	5	1	3	3	1	3	1
PCP	2	2	*	2	0	0	*	*
Ecstasy	11	9	2	5	9	3	6	4
Inhalants	9	11	4	4	6	8	16	13
Nonmedical Use of Psychotherapeutics ^{2,3}	21	18	11	9	11	15	22	29
Pain Relievers	18	15	10	8	10	14	21	25
OxyContin [®]	8	8	3	1	3	6	2	3
Tranquilizers	15	10	4	5	5	5	6	15
Stimulants ³	10	10	4	5	3	6	5	6
Methamphetamine ³	7	4	4	2	2	3	1	3
Sedatives	3	4	2	1	4	1	2	5
ILLCIT DRUGS OTHER THAN MARIJUANA¹	26	22	12	12	17	16	29	33

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	1.83	2.13	2.43	2.47	1.33	1.31	0.19	0.20
Marijuana and Hashish	1.88	2.11	2.44	2.43	1.28	1.28	0.14	0.15
Cocaine	0.80	0.62	0.56	0.43	0.21	0.05	0.00	0.01
Crack	0.18	0.07	*	*	*	0.01	*	*
Heroin	0.63	*	0.12	*	0.01	*	*	*
Hallucinogens	1.25	1.29	0.59	1.17	0.46	0.16	0.05	0.04
LSD	0.59	0.49	0.07	0.41	0.13	0.03	0.01	0.01
PCP	0.15	0.23	*	0.27	0.02	0.01	*	*
Ecstasy	0.89	0.92	0.23	0.72	0.42	0.13	0.03	0.02
Inhalants	0.78	1.03	0.52	0.65	0.29	0.36	0.08	0.06
Nonmedical Use of Psychotherapeutics ^{2,3}	1.63	1.62	1.39	1.37	0.51	0.65	0.10	0.13
Pain Relievers	1.49	1.37	1.19	1.15	0.46	0.59	0.10	0.11
OxyContin [®]	0.71	0.74	0.38	0.17	0.13	0.26	0.01	0.02
Tranquilizers	1.21	0.96	0.49	0.77	0.22	0.23	0.03	0.07
Stimulants ³	0.81	1.00	0.56	0.81	0.14	0.26	0.02	0.03
Methamphetamine ³	0.61	0.44	0.51	0.38	0.08	0.12	0.00	0.01
Sedatives	0.26	0.38	0.23	0.22	0.19	0.05	0.01	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.86	1.89	1.48	1.79	0.75	0.72	0.13	0.15

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.0052		0.3164		0.1423		0.9467	
Marijuana and Hashish	0.0049		0.4954		0.4080		0.9918	
Cocaine	0.0534		0.5148		0.0565		0.5414	
Crack	0.3191		*		*		*	
Heroin	*		*		*		*	
Hallucinogens	0.8205		0.1720		0.0105		0.1961	
LSD	0.8254		0.1374		0.1947		0.5021	
PCP	0.8513		*		0.2832		*	
Ecstasy	0.8059		0.1019		0.0179		0.2487	
Inhalants	0.5555		0.9466		0.2945		0.0314	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.1114		0.6721		0.1165		0.0784	
Pain Relievers	0.0862		0.5665		0.0661		0.2049	
OxyContin [®]	0.7737		0.3017		0.3576		0.3507	
Tranquilizers	0.0264		0.1619		0.6951		0.2630	
Stimulants ³	0.6089		0.4380		0.3608		0.1542	
Methamphetamine ³	0.3361		0.6876		0.4320		0.1021	
Sedatives	0.3855		0.8305		0.0906		0.1965	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0540		0.8281		0.6460		0.7567	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	125	118	37	50	101	93	43	42
Marijuana and Hashish	119	115	34	44	90	89	37	36
Cocaine	38	29	7	11	17	12	3	3
Crack	6	7	1	4	2	4	*	2
Heroin	18	14	5	13	3	4	6	11
Hallucinogens	33	39	9	9	20	19	8	9
LSD	12	15	2	2	5	7	*	5
PCP	6	3	0	0	*	3	*	*
Ecstasy	26	35	7	5	16	15	4	6
Inhalants	35	18	4	4	6	6	2	11
Nonmedical Use of Psychotherapeutics ^{2,3}	58	58	20	29	33	36	19	18
Pain Relievers	48	49	16	29	27	27	16	17
OxyContin [®]	18	17	4	16	9	5	5	*
Tranquilizers	31	33	12	23	18	16	10	5
Stimulants ³	29	24	10	7	11	17	4	5
Methamphetamine ³	11	14	9	5	4	3	2	2
Sedatives	8	5	4	1	5	11	2	3
ILLCIT DRUGS OTHER THAN MARIJUANA¹	76	71	22	31	41	40	21	24

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.92	0.90	1.29	1.59	0.66	0.61	0.37	0.36
Marijuana and Hashish	0.91	0.92	1.21	1.49	0.60	0.59	0.33	0.32
Cocaine	0.41	0.33	0.25	0.41	0.14	0.09	0.03	0.02
Crack	0.06	0.08	0.03	0.15	0.01	0.03	*	0.02
Heroin	0.20	0.16	0.19	0.50	0.03	0.03	0.06	0.10
Hallucinogens	0.36	0.45	0.34	0.34	0.17	0.15	0.07	0.08
LSD	0.14	0.18	0.09	0.09	0.04	0.06	*	0.04
PCP	0.07	0.04	0.01	0.01	*	0.03	*	*
Ecstasy	0.29	0.40	0.26	0.18	0.13	0.12	0.04	0.05
Inhalants	0.39	0.22	0.17	0.14	0.05	0.05	0.01	0.10
Nonmedical Use of Psychotherapeutics ^{2,3}	0.56	0.62	0.75	1.07	0.27	0.27	0.17	0.16
Pain Relievers	0.48	0.54	0.59	1.07	0.22	0.22	0.14	0.15
OxyContin [®]	0.20	0.20	0.14	0.60	0.07	0.04	0.04	*
Tranquilizers	0.33	0.38	0.46	0.86	0.15	0.12	0.09	0.04
Stimulants ³	0.31	0.28	0.37	0.27	0.09	0.13	0.04	0.05
Methamphetamine ³	0.12	0.16	0.33	0.19	0.03	0.02	0.02	0.01
Sedatives	0.09	0.06	0.14	0.05	0.04	0.09	0.02	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.73	0.73	0.83	1.12	0.32	0.30	0.19	0.21

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.9551		0.0009		0.4858		0.4139	
Marijuana and Hashish	0.8933		0.0068		0.5462		0.9151	
Cocaine	0.1339		0.4528		0.9079		0.7438	
Crack	0.7463		0.0899		0.1827		*	
Heroin	0.6063		0.0278		0.4907		0.3327	
Hallucinogens	0.5411		0.3807		0.3340		0.3658	
LSD	0.4693		0.9577		0.3205		*	
PCP	0.4439		0.8825		*		*	
Ecstasy	0.4509		0.0775		0.1626		0.4865	
Inhalants	0.3854		0.7963		0.9329		0.1581	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3580		0.1422		0.6476		0.4034	
Pain Relievers	0.5966		0.0197		0.6131		0.1815	
OxyContin [®]	0.5487		0.1339		0.3543		*	
Tranquilizers	0.4377		0.8159		0.3127		0.3993	
Stimulants ³	0.4132		0.5401		0.0286		0.6057	
Methamphetamine ³	0.0482		0.3631		0.4089		0.7952	
Sedatives	0.3995		0.2473		0.1208		0.4261	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.7842		0.2647		0.4072		0.1623	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	249	289	140	137	261	306	140	135
Marijuana and Hashish	232	259	119	114	227	266	108	99
Cocaine	73	119	25	35	57	57	4	31
Crack	40	82	10	32	*	25	*	20
Heroin	23	34	31	18	7	*	5	23
Hallucinogens	33	73	15	11	13	20	*	7
LSD	2	26	15	*	5	*	*	*
PCP	8	4	*	*	*	11	*	*
Ecstasy	28	59	*	5	10	17	*	5
Inhalants	25	13	5	5	35	54	33	6
Nonmedical Use of Psychotherapeutics ^{2,3}	111	139	79	76	126	159	98	97
Pain Relievers	88	112	70	59	105	120	84	86
OxyContin [®]	32	30	19	19	23	4	18	11
Tranquilizers	69	97	43	38	70	94	59	66
Stimulants ³	52	61	27	36	32	63	44	8
Methamphetamine ³	45	55	23	32	17	30	44	6
Sedatives	29	12	16	17	14	41	31	10
ILLCIT DRUGS OTHER THAN MARIJUANA¹	124	179	83	83	138	174	102	99

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.82	0.88	0.85	0.79	0.31	0.36	0.19	0.18
Marijuana and Hashish	0.77	0.82	0.74	0.67	0.27	0.31	0.15	0.14
Cocaine	0.26	0.42	0.16	0.21	0.07	0.07	0.01	0.04
Crack	0.14	0.29	0.07	0.19	*	0.03	*	0.03
Heroin	0.08	0.12	0.20	0.11	0.01	*	0.01	0.03
Hallucinogens	0.12	0.26	0.10	0.07	0.02	0.02	*	0.01
LSD	0.01	0.09	0.09	*	0.01	*	*	*
PCP	0.03	0.01	*	*	*	0.01	*	*
Ecstasy	0.10	0.21	*	0.03	0.01	0.02	*	0.01
Inhalants	0.09	0.04	0.03	0.03	0.04	0.07	0.05	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.40	0.47	0.49	0.45	0.15	0.19	0.13	0.13
Pain Relievers	0.32	0.38	0.44	0.35	0.13	0.14	0.12	0.12
OxyContin [®]	0.11	0.10	0.12	0.11	0.03	0.00	0.02	0.02
Tranquilizers	0.25	0.34	0.27	0.23	0.09	0.11	0.08	0.09
Stimulants ³	0.19	0.22	0.17	0.21	0.04	0.08	0.06	0.01
Methamphetamine ³	0.16	0.20	0.14	0.19	0.02	0.04	0.06	0.01
Sedatives	0.10	0.04	0.10	0.10	0.02	0.05	0.04	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.44	0.60	0.51	0.49	0.17	0.21	0.14	0.14

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2011	2012	2011	2012	2011	2012	2011	2012
ILLCIT DRUGS¹	0.0084		0.5587		0.1374		0.6804	
Marijuana and Hashish	0.0284		0.7625		0.3560		0.2472	
Cocaine	0.0277		0.6893		0.7946		0.0448	
Crack	0.2983		0.0963		*		*	
Heroin	0.2225		0.2274		*		0.1634	
Hallucinogens	0.0626		0.9008		0.6963		*	
LSD	0.1660		*		*		*	
PCP	0.4717		*		*		*	
Ecstasy	0.4707		*		0.4751		*	
Inhalants	0.1578		0.9949		0.8210		0.0824	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3007		0.0882		0.0327		0.5905	
Pain Relievers	0.0977		0.0196		0.2630		0.7091	
OxyContin [®]	0.9243		1.0000		0.0520		0.5101	
Tranquilizers	0.2935		0.6033		0.2665		0.3509	
Stimulants ³	0.7792		0.5503		0.0071		0.0738	
Methamphetamine ³	0.7417		0.3769		0.1600		0.0766	
Sedatives	0.2662		0.4643		0.0169		0.4049	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0203		0.1737		0.0472		0.7051	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.23C Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Numbers in Thousands, 2011 and 2012

Cigarette Use/Level of Alcohol Use	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
TOTAL	460	492	68	63	142	138	420	452
Heavy Alcohol Use ¹	214	233	20	21	97	84	186	215
Binge Alcohol Use But Not Heavy Use ¹	228	270	32	32	101	94	202	246
Alcohol Use But Not Binge Use ¹	222	257	30	28	67	69	215	247
No Alcohol Use	225	217	50	53	57	65	208	195
PAST MONTH CIGARETTE USE	342	352	47	42	133	133	299	325
Heavy Alcohol Use ¹	166	179	17	18	70	68	144	166
Binge Alcohol Use But Not Heavy Use ¹	161	196	26	24	75	71	139	182
Alcohol Use But Not Binge Use ¹	142	143	20	16	46	48	130	131
No Alcohol Use	147	150	26	22	37	50	140	137
NO PAST MONTH CIGARETTE USE	330	349	52	56	113	103	300	329
Heavy Alcohol Use ¹	126	136	10	11	50	45	116	127
Binge Alcohol Use But Not Heavy Use ¹	162	172	20	22	62	55	148	161
Alcohol Use But Not Binge Use ¹	168	213	22	22	47	47	161	211
No Alcohol Use	154	150	42	45	43	42	140	135

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.23D Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Percentages, 2011 and 2012

Cigarette Use/Level of Alcohol Use	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
TOTAL	0.18	0.19	0.27	0.25	0.42	0.40	0.21	0.23
Heavy Alcohol Use ¹	1.06	1.12	3.04	3.64	1.33	1.28	1.35	1.47
Binge Alcohol Use But Not Heavy Use ¹	0.51	0.57	1.64	1.72	0.81	0.77	0.61	0.71
Alcohol Use But Not Binge Use ¹	0.30	0.33	1.70	1.60	0.81	0.84	0.32	0.36
No Alcohol Use	0.18	0.17	0.22	0.23	0.39	0.44	0.23	0.22
PAST MONTH CIGARETTE USE	0.54	0.53	1.49	1.61	0.80	0.81	0.64	0.63
Heavy Alcohol Use ¹	1.54	1.52	3.13	4.45	1.55	1.56	2.05	2.08
Binge Alcohol Use But Not Heavy Use ¹	0.95	1.05	2.37	2.91	1.28	1.26	1.23	1.38
Alcohol Use But Not Binge Use ¹	1.01	1.08	4.12	4.33	1.86	2.04	1.11	1.17
No Alcohol Use	0.74	0.72	2.43	2.47	1.29	1.59	0.85	0.79
NO PAST MONTH CIGARETTE USE	0.16	0.17	0.21	0.22	0.41	0.37	0.19	0.21
Heavy Alcohol Use ¹	1.48	1.54	5.52	5.75	2.16	2.03	1.83	1.91
Binge Alcohol Use But Not Heavy Use ¹	0.58	0.61	2.04	2.01	0.98	0.90	0.68	0.75
Alcohol Use But Not Binge Use ¹	0.27	0.33	1.70	1.66	0.81	0.81	0.29	0.37
No Alcohol Use	0.15	0.14	0.19	0.20	0.37	0.36	0.19	0.18

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.23P Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Cigarette Use/Level of Alcohol Use	Total (2011)	Total (2012)	Aged 12-17 (2011)	Aged 12-17 (2012)	Aged 18-25 (2011)	Aged 18-25 (2012)	Aged 26+ (2011)	Aged 26+ (2012)
TOTAL	0.0723		0.1309		0.8661		0.0215	
Heavy Alcohol Use ¹	0.8722		0.6027		0.4123		0.7198	
Binge Alcohol Use But Not Heavy Use ¹	0.0798		0.2808		0.3539		0.0120	
Alcohol Use But Not Binge Use ¹	0.3180		0.6089		0.2812		0.1452	
No Alcohol Use	0.8033		0.6336		0.0091		0.6822	
PAST MONTH CIGARETTE USE	0.2146		0.1642		0.2264		0.0667	
Heavy Alcohol Use ¹	0.8907		0.2192		0.3658		0.4046	
Binge Alcohol Use But Not Heavy Use ¹	0.2512		0.0654		0.9956		0.0378	
Alcohol Use But Not Binge Use ¹	0.4188		0.1173		0.6842		0.3077	
No Alcohol Use	0.9226		0.3164		0.0009		0.5587	
NO PAST MONTH CIGARETTE USE	0.1688		0.3872		0.8899		0.2070	
Heavy Alcohol Use ¹	0.9190		0.2843		0.6962		0.7420	
Binge Alcohol Use But Not Heavy Use ¹	0.2836		0.3465		0.4490		0.2580	
Alcohol Use But Not Binge Use ¹	0.2018		0.4524		0.2913		0.1316	
No Alcohol Use	0.9245		0.9467		0.4139		0.6804	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.