

Table 6.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	367	366
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	161	174	223	247
Cigars	231	268	290	284
Pipe Tobacco	95	120	110	140
ALCOHOL	716	700	1,529	1,532
Binge Alcohol Use ²	491	527	717	728
Heavy Alcohol Use ²	282	311	324	334

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	0.18	0.18
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.27	0.29	0.11	0.12
Cigars	0.39	0.42	0.14	0.14
Pipe Tobacco	0.16	0.20	0.05	0.07
ALCOHOL	0.71	0.73	0.47	0.45
Binge Alcohol Use ²	0.67	0.70	0.30	0.32
Heavy Alcohol Use ²	0.46	0.48	0.16	0.16

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A		0.3410	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.4343		0.1085	
Cigars	0.1518		0.8538	
Pipe Tobacco	0.4288		0.1462	
ALCOHOL	0.3232		0.2837	
Binge Alcohol Use ²	0.2636		0.8949	
Heavy Alcohol Use ²	0.5137		0.1034	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	31	29
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	22	23	19	20
Cigars	32	26	24	20
Pipe Tobacco	14	14	11	11
ALCOHOL	46	47	70	73
Binge Alcohol Use ²	37	40	44	48
Heavy Alcohol Use ²	19	21	15	15

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	0.13	0.12
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1.04	1.25	0.08	0.08
Cigars	1.35	1.37	0.10	0.08
Pipe Tobacco	0.71	0.83	0.05	0.05
ALCOHOL	1.53	1.62	0.28	0.28
Binge Alcohol Use ²	1.42	1.69	0.19	0.19
Heavy Alcohol Use ²	0.94	1.14	0.07	0.06

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A		0.3472	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3621		0.5898	
Cigars	0.0242		0.0474	
Pipe Tobacco	0.8946		0.9140	
ALCOHOL	0.6710		0.7267	
Binge Alcohol Use ²	0.7042		0.3392	
Heavy Alcohol Use ²	0.6379		0.8647	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	82	85
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	58	50	42	49
Cigars	94	86	66	62
Pipe Tobacco	37	31	33	27
ALCOHOL	204	190	290	276
Binge Alcohol Use ²	171	169	203	196
Heavy Alcohol Use ²	91	97	92	98

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	0.33	0.32
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.47	0.43	0.18	0.20
Cigars	0.65	0.64	0.27	0.24
Pipe Tobacco	0.31	0.27	0.15	0.11
ALCOHOL	0.69	0.74	0.67	0.62
Binge Alcohol Use ²	0.77	0.87	0.63	0.57
Heavy Alcohol Use ²	0.65	0.70	0.36	0.37

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A		0.6766	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.7119		0.5055	
Cigars	0.8771		0.4278	
Pipe Tobacco	0.9365		0.9130	
ALCOHOL	0.3690		0.7037	
Binge Alcohol Use ²	0.9974		0.6779	
Heavy Alcohol Use ²	0.3095		0.0898	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	360	350
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	146	165	218	237
Cigars	206	250	281	277
Pipe Tobacco	87	114	104	137
ALCOHOL	659	665	1,459	1,476
Binge Alcohol Use ²	445	493	671	693
Heavy Alcohol Use ²	266	296	302	318

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2011 and 2012

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A	N/A	0.23	0.22
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.33	0.36	0.14	0.15
Cigars	0.46	0.52	0.18	0.18
Pipe Tobacco	0.20	0.25	0.07	0.09
ALCOHOL	0.89	0.90	0.59	0.56
Binge Alcohol Use ²	0.81	0.85	0.38	0.40
Heavy Alcohol Use ²	0.57	0.59	0.19	0.20

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2011	2012	2011	2012
TOBACCO PRODUCTS¹	N/A		0.3529	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3558		0.1418	
Cigars	0.0225		0.8073	
Pipe Tobacco	0.3255		0.1262	
ALCOHOL	0.4483		0.2967	
Binge Alcohol Use ²	0.1457		0.9288	
Heavy Alcohol Use ²	0.5576		0.2071	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.