

130408

Table 6.28C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	319	347	431	436	477	465	550	601
Cigarettes	282	311	385	385	450	422	516	564
Smokeless Tobacco	125	126	147	152	123	126	148	182
Cigars	136	160	209	205	215	202	175	188
Pipe Tobacco	39	42	70	76	85	109	86	109

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.28D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	1.27	1.32	0.80	0.79	0.60	0.56	0.40	0.43
Cigarettes	1.32	1.33	0.77	0.74	0.58	0.53	0.38	0.41
Smokeless Tobacco	0.73	0.70	0.34	0.35	0.16	0.17	0.12	0.14
Cigars	0.77	0.87	0.47	0.46	0.28	0.26	0.14	0.15
Pipe Tobacco	0.25	0.25	0.16	0.18	0.11	0.14	0.07	0.09

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.28P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	0.4903		0.2314		0.0494		0.1556	
Cigarettes	0.4144		0.2893		0.0368		0.2683	
Smokeless Tobacco	0.4329		0.3525		0.4415		0.4843	
Cigars	0.0690		0.3819		0.4030		0.7913	
Pipe Tobacco	0.8852		0.8178		0.0810		0.2563	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.29C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	21	23	36	35	29	25	47	39
Cigarettes	19	21	32	33	25	23	40	33
Smokeless Tobacco	14	11	14	17	13	10	17	19
Cigars	14	11	23	20	17	15	25	17
Pipe Tobacco	7	5	13	9	4	10	10	9

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.29D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	2.71	3.38	1.71	1.81	1.64	1.53	0.21	0.18
Cigarettes	3.16	3.69	1.67	1.81	1.53	1.44	0.19	0.15
Smokeless Tobacco	3.15	3.08	0.96	1.11	0.85	0.70	0.08	0.09
Cigars	2.96	3.07	1.39	1.24	1.06	1.02	0.11	0.08
Pipe Tobacco	1.86	1.56	0.84	0.60	0.30	0.69	0.05	0.04

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.29P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	0.2002		0.0215		0.3813		0.0097	
Cigarettes	0.3155		0.0418		0.2635		0.0067	
Smokeless Tobacco	0.3996		0.8480		0.3598		0.1860	
Cigars	0.1353		0.0526		0.3208		0.0013	
Pipe Tobacco	0.3301		0.0450		0.0101		0.8532	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.30C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	103	110	142	129	88	78	100	103
Cigarettes	91	97	129	116	79	71	90	97
Smokeless Tobacco	39	43	47	40	18	23	28	30
Cigars	58	57	70	60	42	42	48	44
Pipe Tobacco	20	19	31	23	21	18	21	19

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.30D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	1.30	1.35	0.88	0.85	1.00	0.91	0.65	0.66
Cigarettes	1.36	1.48	0.90	0.84	0.93	0.86	0.60	0.63
Smokeless Tobacco	0.85	0.88	0.47	0.42	0.25	0.32	0.21	0.22
Cigars	1.09	1.06	0.64	0.58	0.57	0.54	0.34	0.30
Pipe Tobacco	0.48	0.41	0.31	0.24	0.29	0.25	0.16	0.14

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.30P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	0.2049		0.1086		0.3011		0.4485	
Cigarettes	0.0790		0.0896		0.0499		0.5507	
Smokeless Tobacco	0.1241		0.1159		0.2326		0.9368	
Cigars	0.7922		0.4959		0.4743		0.2043	
Pipe Tobacco	0.9691		0.4483		0.9911		0.6691	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.31C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	303	327	391	415	465	456	533	578
Cigarettes	266	296	353	370	438	415	496	540
Smokeless Tobacco	121	115	137	144	122	125	145	178
Cigars	125	150	189	196	209	199	167	181
Pipe Tobacco	33	36	60	71	82	107	83	106

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.31D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2011 and 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	1.70	1.77	1.02	1.02	0.67	0.61	0.54	0.58
Cigarettes	1.74	1.77	0.97	0.96	0.64	0.58	0.51	0.55
Smokeless Tobacco	1.00	0.89	0.43	0.45	0.18	0.18	0.16	0.20
Cigars	1.01	1.13	0.59	0.59	0.31	0.28	0.19	0.20
Pipe Tobacco	0.29	0.29	0.19	0.22	0.12	0.16	0.09	0.12

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 6.31P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2012 Versus 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2011	2012	2011	2012	2011	2012	2011	2012
TOBACCO PRODUCTS²	0.7973		0.0567		0.0759		0.0709	
Cigarettes	0.7877		0.0787		0.0824		0.1478	
Smokeless Tobacco	0.6433		0.1440		0.5036		0.5703	
Cigars	0.0377		0.3278		0.5454		0.6862	
Pipe Tobacco	0.9206		0.6706		0.1107		0.2719	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.