

**Table 6.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	105	108	16	17	26	27	101	103
From More Than One Doctor	35	44	5	5	9	9	34	43
Wrote Fake Prescription	9	6	2	4	1	4	9	3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	26	5	4	2	6	4	26
From Friend or Relative for Free	163	164	24	22	60	52	150	154
Bought from Friend or Relative	72	61	9	9	28	26	66	54
Took from Friend or Relative without Asking	40	39	12	10	15	16	35	35
Bought from Drug Dealer or Other Stranger	45	33	7	6	19	17	41	28
Bought on the Internet	19	13	0	1	1	4	19	12
Some Other Way	47	48	10	8	14	11	42	46

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.82	0.87	1.08	1.27	0.66	0.77	1.39	1.37
From More Than One Doctor	0.30	0.40	0.38	0.44	0.23	0.29	0.52	0.65
Wrote Fake Prescription	0.08	0.06	0.16	0.32	0.04	0.11	0.14	0.05
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.05	0.24	0.36	0.34	0.05	0.18	0.06	0.39
From Friend or Relative for Free	1.00	1.00	1.34	1.44	0.95	1.04	1.67	1.60
Bought from Friend or Relative	0.58	0.52	0.69	0.81	0.67	0.74	0.96	0.79
Took from Friend or Relative without Asking	0.34	0.36	0.84	0.81	0.40	0.47	0.54	0.53
Bought from Drug Dealer or Other Stranger	0.38	0.31	0.52	0.53	0.49	0.52	0.63	0.42
Bought on the Internet	0.17	0.12	0.03	0.09	0.04	0.12	0.30	0.19
Some Other Way	0.40	0.44	0.78	0.69	0.37	0.34	0.65	0.70

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.0467		0.3232		0.1539		0.1642	
From More Than One Doctor	0.6315		0.8657		0.3517		0.7444	
Wrote Fake Prescription	0.7877		0.5578		0.0323		0.4134	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0173		0.3832		0.0167		0.0297	
From Friend or Relative for Free	0.4499		0.6029		0.0589		0.6931	
Bought from Friend or Relative	0.6234		0.6467		0.3759		0.4210	
Took from Friend or Relative without Asking	0.0980		0.2321		0.8478		0.2162	
Bought from Drug Dealer or Other Stranger	0.8856		0.4419		0.5565		0.8997	
Bought on the Internet	0.3348		0.3615		0.5217		0.2312	
Some Other Way	0.7131		0.2387		0.3777		0.9032	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	142	148	16	16	46	41	134	142
From More Than One Doctor	27	28	6	5	10	11	24	23
Wrote Fake Prescription	11	11	1	*	1	2	11	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	6	3	1	3	5	2	2
From Friend or Relative for Free	40	29	7	6	15	12	37	25
Bought from Friend or Relative	23	23	5	5	13	13	19	19
Took from Friend or Relative without Asking	15	12	8	5	8	8	10	7
Bought from Drug Dealer or Other Stranger	18	14	3	4	9	8	14	11
Bought on the Internet	5	8	1	0	2	1	4	8
Some Other Way	12	14	5	4	5	4	10	13

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.00	1.03	2.18	2.18	1.29	1.36	1.47	1.37
From More Than One Doctor	0.48	0.51	1.14	1.15	0.54	0.70	0.71	0.69
Wrote Fake Prescription	0.19	0.20	0.13	*	0.06	0.13	0.31	0.31
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.08	0.11	0.69	0.32	0.16	0.33	0.05	0.06
From Friend or Relative for Free	0.69	0.53	1.31	1.29	0.79	0.75	1.07	0.76
Bought from Friend or Relative	0.42	0.44	1.08	1.12	0.72	0.83	0.56	0.57
Took from Friend or Relative without Asking	0.26	0.23	1.44	1.11	0.42	0.56	0.31	0.21
Bought from Drug Dealer or Other Stranger	0.32	0.27	0.71	1.00	0.52	0.49	0.42	0.34
Bought on the Internet	0.09	0.15	0.11	0.05	0.10	0.08	0.13	0.23
Some Other Way	0.21	0.27	0.94	0.85	0.29	0.26	0.29	0.40

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.0537		0.3049		0.3636		0.2503	
From More Than One Doctor	0.9956		0.3050		0.3551		0.6015	
Wrote Fake Prescription	0.7203		*		0.4391		0.7845	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3832		0.7196		0.1788		0.7532	
From Friend or Relative for Free	0.2914		0.6375		0.2130		0.7000	
Bought from Friend or Relative	0.5232		0.4511		0.9753		0.5162	
Took from Friend or Relative without Asking	0.0530		0.0372		0.9894		0.2820	
Bought from Drug Dealer or Other Stranger	0.0272		0.7834		0.0764		0.2031	
Bought on the Internet	0.8790		0.6313		0.5391		0.7580	
Some Other Way	0.7157		0.7421		0.4287		0.9130	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	87	74	11	10	20	21	84	70
From More Than One Doctor	36	27	5	4	9	9	33	25
Wrote Fake Prescription	10	14	2	3	1	1	9	13
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	20	4	3	3	4	6	19
From Friend or Relative for Free	109	108	16	13	38	36	102	100
Bought from Friend or Relative	70	64	8	8	25	23	63	60
Took from Friend or Relative without Asking	47	43	9	7	16	14	43	41
Bought from Drug Dealer or Other Stranger	37	37	7	5	19	18	30	31
Bought on the Internet	19	17	*	1	5	6	18	16
Some Other Way	39	25	6	5	11	11	37	22

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.50	1.46	1.87	1.94	1.22	1.54	2.41	2.22
From More Than One Doctor	0.74	0.63	0.93	0.96	0.62	0.74	1.16	0.95
Wrote Fake Prescription	0.21	0.32	0.39	0.76	0.09	0.11	0.34	0.49
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.16	0.46	0.75	0.60	0.21	0.33	0.20	0.72
From Friend or Relative for Free	1.52	1.64	2.19	2.17	1.35	1.72	2.45	2.50
Bought from Friend or Relative	1.28	1.38	1.47	1.62	1.44	1.60	2.03	2.12
Took from Friend or Relative without Asking	0.93	0.97	1.56	1.56	1.01	1.09	1.47	1.47
Bought from Drug Dealer or Other Stranger	0.77	0.88	1.28	1.17	1.16	1.37	1.09	1.18
Bought on the Internet	0.40	0.39	*	0.27	0.34	0.50	0.66	0.59
Some Other Way	0.80	0.60	1.19	1.15	0.78	0.96	1.30	0.84

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.9586		0.7105		0.3984		0.7072	
From More Than One Doctor	0.2743		0.3483		0.4938		0.3981	
Wrote Fake Prescription	0.6277		0.5410		0.9427		0.7588	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0566		0.6640		0.1153		0.0793	
From Friend or Relative for Free	0.8805		0.9432		0.2036		0.8190	
Bought from Friend or Relative	0.6933		0.6181		0.2744		0.8711	
Took from Friend or Relative without Asking	0.4485		0.1693		0.3675		0.8716	
Bought from Drug Dealer or Other Stranger	0.5281		0.2824		0.2023		0.5321	
Bought on the Internet	0.9018		*		0.5424		0.8389	
Some Other Way	0.0718		0.9731		0.8856		0.0700	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	93	96	10	9	31	28	89	91
From More Than One Doctor	31	25	5	4	11	13	28	21
Wrote Fake Prescription	13	17	1	1	4	2	12	17
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	11	1	1	5	5	2	10
From Friend or Relative for Free	37	33	6	4	15	14	33	29
Bought from Friend or Relative	31	29	6	4	15	14	27	25
Took from Friend or Relative without Asking	17	18	5	3	9	11	13	14
Bought from Drug Dealer or Other Stranger	27	23	5	4	13	12	23	19
Bought on the Internet	3	12	1	*	2	3	2	11
Some Other Way	12	15	4	3	7	6	8	13

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.48	1.64	3.33	3.27	1.77	2.12	2.28	2.26
From More Than One Doctor	1.15	1.05	2.33	2.23	1.17	1.78	1.79	1.39
Wrote Fake Prescription	0.50	0.73	0.27	0.64	0.47	0.30	0.81	1.12
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.21	0.46	0.60	0.79	0.54	0.69	0.15	0.66
From Friend or Relative for Free	1.35	1.34	2.54	2.26	1.45	1.86	2.13	1.87
Bought from Friend or Relative	1.17	1.16	2.48	2.13	1.54	1.88	1.74	1.59
Took from Friend or Relative without Asking	0.65	0.77	2.29	1.77	1.00	1.62	0.89	0.92
Bought from Drug Dealer or Other Stranger	1.01	0.95	2.06	2.27	1.40	1.67	1.50	1.24
Bought on the Internet	0.12	0.50	0.42	*	0.29	0.51	0.12	0.74
Some Other Way	0.46	0.64	1.93	1.40	0.85	0.91	0.53	0.90

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.2644		0.0969		0.4161		0.8159	
From More Than One Doctor	0.4042		0.6820		0.3725		0.1917	
Wrote Fake Prescription	0.4615		0.5845		0.3666		0.4070	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6026		0.7108		0.4506		0.2265	
From Friend or Relative for Free	0.5222		0.1781		0.2638		0.8813	
Bought from Friend or Relative	0.4471		0.1754		0.6260		0.9620	
Took from Friend or Relative without Asking	0.8797		0.2014		0.7135		0.7861	
Bought from Drug Dealer or Other Stranger	0.4092		0.4567		0.5309		0.9449	
Bought on the Internet	0.2512		*		0.6877		0.2416	
Some Other Way	0.8169		0.1765		0.4442		0.4311	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	61	61	6	6	13	12	59	59
From More Than One Doctor	19	11	2	2	5	7	19	8
Wrote Fake Prescription	1	1	1	1	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	5	2	1	5	3	*	4
From Friend or Relative for Free	119	131	13	14	37	33	114	126
Bought from Friend or Relative	42	53	7	8	23	21	35	48
Took from Friend or Relative without Asking	21	22	5	6	9	12	18	17
Bought from Drug Dealer or Other Stranger	29	29	5	6	13	15	24	23
Bought on the Internet	15	1	1	*	2	1	15	*
Some Other Way	26	24	4	3	9	11	24	21

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.12	1.08	1.46	1.41	0.78	0.77	1.79	1.63
From More Than One Doctor	0.37	0.20	0.50	0.44	0.30	0.43	0.60	0.25
Wrote Fake Prescription	0.02	0.03	0.21	0.31	0.06	0.04	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.10	0.09	0.46	0.13	0.29	0.20	*	0.11
From Friend or Relative for Free	1.46	1.43	2.38	2.40	1.40	1.48	2.23	2.04
Bought from Friend or Relative	0.80	0.95	1.66	1.97	1.17	1.21	1.10	1.36
Took from Friend or Relative without Asking	0.40	0.42	1.27	1.47	0.53	0.76	0.58	0.52
Bought from Drug Dealer or Other Stranger	0.55	0.57	1.24	1.53	0.78	0.96	0.74	0.69
Bought on the Internet	0.29	0.01	0.14	*	0.12	0.04	0.47	*
Some Other Way	0.50	0.45	1.10	0.79	0.54	0.72	0.76	0.62

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.8175		0.6420		0.6987		0.8512	
From More Than One Doctor	0.3342		0.9538		0.5583		0.2205	
Wrote Fake Prescription	0.8273		0.7592		0.5607		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9945		0.2240		0.9250		*	
From Friend or Relative for Free	0.5639		0.9487		0.2171		0.4293	
Bought from Friend or Relative	0.7261		0.2829		0.5561		0.7990	
Took from Friend or Relative without Asking	0.2541		0.8126		0.7794		0.2856	
Bought from Drug Dealer or Other Stranger	0.8898		0.9680		0.1098		0.6297	
Bought on the Internet	0.1603		*		0.2239		*	
Some Other Way	0.3779		0.1765		0.8647		0.4370	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	105	118	9	8	26	27	101	115
From More Than One Doctor	15	22	3	2	7	6	13	21
Wrote Fake Prescription	2	1	*	*	1	1	2	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	4	*	1	1	2	3	4
From Friend or Relative for Free	25	26	5	4	13	10	20	23
Bought from Friend or Relative	11	19	3	6	9	7	5	16
Took from Friend or Relative without Asking	5	7	3	2	4	4	2	6
Bought from Drug Dealer or Other Stranger	13	8	3	3	8	5	10	5
Bought on the Internet	4	1	*	*	1	*	4	*
Some Other Way	7	9	3	2	2	6	7	6

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.31	1.37	3.80	4.22	1.92	1.98	1.61	1.73
From More Than One Doctor	0.57	0.76	1.68	1.38	0.88	0.91	0.77	1.03
Wrote Fake Prescription	0.09	0.05	*	*	0.09	0.20	0.11	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.13	0.16	*	0.38	0.11	0.23	0.19	0.20
From Friend or Relative for Free	0.93	0.89	2.65	2.81	1.60	1.38	1.16	1.13
Bought from Friend or Relative	0.42	0.64	1.66	3.80	1.11	0.99	0.31	0.78
Took from Friend or Relative without Asking	0.20	0.26	1.84	1.39	0.50	0.64	0.10	0.28
Bought from Drug Dealer or Other Stranger	0.50	0.27	1.69	2.07	0.96	0.69	0.60	0.25
Bought on the Internet	0.16	0.03	*	*	0.10	*	0.24	*
Some Other Way	0.28	0.31	1.63	1.03	0.24	0.81	0.38	0.30

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.3200		0.3133		0.0336		0.6814	
From More Than One Doctor	0.9234		0.9941		0.8425		0.7643	
Wrote Fake Prescription	0.2356		*		0.7203		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6886		*		0.2211		0.9658	
From Friend or Relative for Free	0.2021		0.7176		0.0403		0.9231	
Bought from Friend or Relative	0.1383		0.1016		0.4056		0.0138	
Took from Friend or Relative without Asking	0.5120		0.4666		0.4952		0.1611	
Bought from Drug Dealer or Other Stranger	0.0175		0.4481		0.0436		0.1622	
Bought on the Internet	0.0983		*		*		*	
Some Other Way	0.9871		0.1488		0.0710		0.4746	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	33	52	4	4	10	9	31	51
From More Than One Doctor	6	5	*	1	2	4	5	2
Wrote Fake Prescription	1	1	1	1	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	13	*	*	*	3	*	*
From Friend or Relative for Free	82	81	5	8	19	19	78	77
Bought from Friend or Relative	39	36	4	5	14	13	36	34
Took from Friend or Relative without Asking	15	17	4	3	7	8	13	14
Bought from Drug Dealer or Other Stranger	18	21	4	5	9	10	14	18
Bought on the Internet	7	*	1	*	3	0	7	*
Some Other Way	16	15	2	2	4	7	15	13

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.68	2.39	3.05	3.18	1.65	1.63	2.49	3.41
From More Than One Doctor	0.31	0.27	*	0.97	0.30	0.89	0.43	0.15
Wrote Fake Prescription	0.05	0.07	0.63	0.80	0.13	0.19	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.09	0.69	*	*	*	0.55	*	*
From Friend or Relative for Free	2.41	2.63	4.05	4.33	2.35	2.67	3.34	3.66
Bought from Friend or Relative	1.91	1.80	3.53	3.67	2.12	2.20	2.72	2.51
Took from Friend or Relative without Asking	0.79	0.88	3.18	2.15	1.32	1.57	1.04	1.11
Bought from Drug Dealer or Other Stranger	0.94	1.14	2.98	3.57	1.63	1.91	1.13	1.42
Bought on the Internet	0.38	*	0.64	*	0.47	0.02	0.53	*
Some Other Way	0.82	0.80	1.87	2.00	0.70	1.33	1.21	1.04

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.4666		0.5453		0.6772		0.3812	
From More Than One Doctor	0.7457		*		0.1819		0.1413	
Wrote Fake Prescription	0.5441		0.8849		0.5382		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1599		*		*		*	
From Friend or Relative for Free	0.8277		0.0847		0.9606		0.6454	
Bought from Friend or Relative	0.1670		0.7044		0.2027		0.3584	
Took from Friend or Relative without Asking	0.3472		0.4190		0.4723		0.6203	
Bought from Drug Dealer or Other Stranger	0.6113		0.9121		0.7148		0.5756	
Bought on the Internet	*		*		0.1105		*	
Some Other Way	0.9407		0.8265		0.7651		0.7977	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	69	69	*	*	14	14	67	67
From More Than One Doctor	13	20	*	*	6	3	12	19
Wrote Fake Prescription	1	*	*	*	1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	4	*	*	1	2	*	*
From Friend or Relative for Free	24	17	*	*	8	6	22	16
Bought from Friend or Relative	12	11	*	*	6	5	9	9
Took from Friend or Relative without Asking	10	7	*	*	1	3	*	6
Bought from Drug Dealer or Other Stranger	15	5	*	*	5	3	14	3
Bought on the Internet	4	1	*	*	*	*	*	*
Some Other Way	3	8	*	*	2	*	2	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	2.87	2.72	*	*	3.24	3.63	3.77	3.56
From More Than One Doctor	1.28	1.85	*	*	2.07	1.44	1.57	2.52
Wrote Fake Prescription	0.08	*	*	*	0.32	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.16	0.41	*	*	0.49	0.79	*	*
From Friend or Relative for Free	2.19	1.72	*	*	2.74	2.63	2.81	2.17
Bought from Friend or Relative	1.20	1.11	*	*	2.35	2.08	1.28	1.18
Took from Friend or Relative without Asking	0.93	0.68	*	*	0.42	1.40	*	0.80
Bought from Drug Dealer or Other Stranger	1.46	0.51	*	*	1.83	1.47	1.90	0.43
Bought on the Internet	0.37	0.09	*	*	*	*	*	*
Some Other Way	0.33	0.80	*	*	0.82	*	0.31	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.9754		*		0.3479		0.7952	
From More Than One Doctor	0.5347		*		0.0946		0.2628	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3331		*		0.3992		*	
From Friend or Relative for Free	0.2108		*		0.0725		0.3845	
Bought from Friend or Relative	0.5526		*		0.3921		0.5514	
Took from Friend or Relative without Asking	0.9060		*		0.0655		*	
Bought from Drug Dealer or Other Stranger	0.0399		*		0.5646		0.0527	
Bought on the Internet	0.2365		*		*		*	
Some Other Way	0.3730		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	26	35	5	4	10	10	24	33
From More Than One Doctor	10	8	2	1	3	7	10	4
Wrote Fake Prescription	7	2	1	*	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	4	2	1	2	2	10	3
From Friend or Relative for Free	62	65	9	10	32	29	53	59
Bought from Friend or Relative	29	39	6	6	18	26	20	28
Took from Friend or Relative without Asking	19	15	4	4	7	5	17	14
Bought from Drug Dealer or Other Stranger	31	17	4	4	9	11	29	13
Bought on the Internet	11	5	2	*	3	1	10	5
Some Other Way	17	12	3	1	6	7	16	10

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	1.12	1.47	1.92	1.95	0.90	0.88	2.41	3.10
From More Than One Doctor	0.46	0.38	0.95	0.67	0.25	0.67	1.01	0.45
Wrote Fake Prescription	0.32	0.08	0.56	*	0.01	0.08	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.46	0.18	1.04	0.55	0.16	0.21	1.02	0.32
From Friend or Relative for Free	1.86	1.99	3.32	3.07	1.74	2.09	3.99	3.89
Bought from Friend or Relative	1.27	1.55	2.62	2.51	1.55	2.00	2.06	2.66
Took from Friend or Relative without Asking	0.82	0.67	1.65	1.65	0.67	0.45	1.75	1.45
Bought from Drug Dealer or Other Stranger	1.30	0.78	1.70	1.66	0.80	1.05	2.91	1.32
Bought on the Internet	0.49	0.24	0.75	*	0.33	0.14	1.06	0.52
Some Other Way	0.77	0.55	1.12	0.58	0.53	0.68	1.69	1.01

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.7267		0.6176		0.1581		0.4812	
From More Than One Doctor	0.7726		0.5979		0.0560		0.4133	
Wrote Fake Prescription	0.3101		*		0.3694		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2820		0.2129		0.4085		0.2932	
From Friend or Relative for Free	0.5964		0.1280		0.5248		0.5502	
Bought from Friend or Relative	0.1756		0.8048		0.4314		0.1549	
Took from Friend or Relative without Asking	0.1453		0.7282		0.0697		0.4170	
Bought from Drug Dealer or Other Stranger	0.8725		0.6437		0.1864		0.5315	
Bought on the Internet	0.0531		*		0.1129		0.1779	
Some Other Way	0.1892		0.1754		0.8300		0.1730	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	50	60	7	7	24	24	43	54
From More Than One Doctor	10	7	3	3	5	6	8	3
Wrote Fake Prescription	1	*	*	*	0	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	3	2	*	*
From Friend or Relative for Free	17	14	3	3	8	7	*	12
Bought from Friend or Relative	9	10	2	3	8	7	3	6
Took from Friend or Relative without Asking	3	4	1	3	2	2	1	*
Bought from Drug Dealer or Other Stranger	13	5	2	2	6	5	*	2
Bought on the Internet	6	2	*	1	*	0	*	*
Some Other Way	10	2	0	1	1	1	*	0

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	2.31	1.94	4.78	4.59	2.42	2.34	4.73	2.75
From More Than One Doctor	1.00	0.71	2.94	2.70	1.03	1.22	1.90	0.59
Wrote Fake Prescription	0.07	*	*	*	0.09	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.26	0.15	*	*	0.54	0.33	*	*
From Friend or Relative for Free	1.63	1.31	3.07	2.83	1.64	1.44	*	2.29
Bought from Friend or Relative	0.88	0.94	2.24	2.78	1.55	1.42	0.73	1.26
Took from Friend or Relative without Asking	0.28	0.34	1.65	2.72	0.37	0.53	0.19	*
Bought from Drug Dealer or Other Stranger	1.27	0.52	1.97	1.81	1.15	1.00	*	0.41
Bought on the Internet	0.62	0.21	*	0.68	*	0.07	*	*
Some Other Way	0.99	0.15	0.47	0.91	0.18	0.29	*	0.03

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.1046		0.6378		0.9116		0.0645	
From More Than One Doctor	0.4824		0.8887		0.4137		0.1374	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2892		*		0.3717		*	
From Friend or Relative for Free	0.8145		0.5886		0.6168		*	
Bought from Friend or Relative	0.7140		0.8521		0.9696		0.2455	
Took from Friend or Relative without Asking	0.3626		0.5857		0.2137		*	
Bought from Drug Dealer or Other Stranger	0.1005		0.9268		0.5995		*	
Bought on the Internet	0.4606		*		*		*	
Some Other Way	0.2322		0.4319		0.7405		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Stimulant Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	25	29	4	3	9	5	22	*
From More Than One Doctor	10	9	1	2	2	2	*	*
Wrote Fake Prescription	1	*	1	*	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	9	*	1	1	2	*	*
From Friend or Relative for Free	37	32	5	6	18	17	32	*
Bought from Friend or Relative	24	27	4	4	13	14	19	*
Took from Friend or Relative without Asking	18	12	2	2	6	5	16	*
Bought from Drug Dealer or Other Stranger	15	17	4	3	7	8	13	16
Bought on the Internet	8	10	1	*	3	1	*	*
Some Other Way	9	11	2	1	3	5	8	10

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Stimulant Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	2.56	3.33	3.93	3.86	2.31	1.43	4.77	*
From More Than One Doctor	1.20	1.23	1.13	2.11	0.69	0.63	*	*
Wrote Fake Prescription	0.15	*	1.32	*	0.16	0.03	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.69	1.14	*	0.97	0.17	0.54	*	*
From Friend or Relative for Free	3.06	3.34	5.24	5.26	3.04	3.13	5.52	*
Bought from Friend or Relative	2.55	3.08	4.08	4.68	2.89	3.19	4.20	*
Took from Friend or Relative without Asking	1.95	1.54	2.87	2.86	1.67	1.47	3.65	*
Bought from Drug Dealer or Other Stranger	1.68	2.19	3.98	3.96	1.83	2.33	2.97	4.03
Bought on the Internet	0.93	1.30	0.81	*	0.75	0.43	*	*
Some Other Way	1.08	1.46	2.31	1.45	0.73	1.63	2.09	2.59

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Past Month Nonmedical Use among Past Month Stimulant Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.7546		0.5857		0.0046		*	
From More Than One Doctor	0.8485		0.5648		0.8564		*	
Wrote Fake Prescription	*		*		0.4147		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7152		*		0.2659		*	
From Friend or Relative for Free	0.9729		0.1152		0.7406		*	
Bought from Friend or Relative	0.0668		0.0351		0.5118		*	
Took from Friend or Relative without Asking	0.2000		0.7052		0.8040		*	
Bought from Drug Dealer or Other Stranger	0.2258		0.5107		0.2306		0.3537	
Bought on the Internet	0.7924		*		0.3801		*	
Some Other Way	0.4262		0.6581		0.2846		0.6431	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	28	29	*	*	12	16	*	*
From More Than One Doctor	7	9	*	*	5	2	*	*
Wrote Fake Prescription	0	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0	*	*	*	*	*	*	*
From Friend or Relative for Free	16	10	*	*	6	3	*	*
Bought from Friend or Relative	8	10	*	*	6	3	*	*
Took from Friend or Relative without Asking	4	9	*	*	*	2	*	*
Bought from Drug Dealer or Other Stranger	5	11	*	*	3	3	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	2	*	*	0	1	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specific questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	4.54	2.93	*	*	4.37	2.76	*	*
From More Than One Doctor	2.02	2.69	*	*	2.93	1.16	*	*
Wrote Fake Prescription	0.03	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.09	*	*	*	*	*	*	*
From Friend or Relative for Free	4.08	2.82	*	*	3.55	2.26	*	*
Bought from Friend or Relative	2.30	2.82	*	*	3.40	1.84	*	*
Took from Friend or Relative without Asking	1.26	2.63	*	*	*	1.60	*	*
Bought from Drug Dealer or Other Stranger	1.28	3.26	*	*	1.99	1.81	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	0.49	*	*	0.25	0.46	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specific questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.0585		*		0.0428		*	
From More Than One Doctor	0.9237		*		0.1322		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.1232		*		0.1461		*	
Bought from Friend or Relative	0.7525		*		0.3734		*	
Took from Friend or Relative without Asking	0.4447		*		*		*	
Bought from Drug Dealer or Other Stranger	0.0443		*		0.8029		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		0.5567		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.59C Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

<b>Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users<sup>1</sup></b>	<b>Total (2009-2010)</b>	<b>Total (2011-2012)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 12-17 (2011-2012)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 18-25 (2011-2012)</b>	<b>Aged 26+ (2009-2010)</b>	<b>Aged 26+ (2011-2012)</b>
From Friend or Relative for Free	33	29	*	*	14	9	30	*
Bought from Friend or Relative	52	*	3	*	9	3	49	*
Took from Friend or Relative without Asking	14	8	2	*	3	*	14	*
Bought from Drug Dealer or Other Stranger	25	*	3	*	9	5	23	*
Bought on the Internet	5	2	*	*	2	2	*	*
Some Other Way	12	6	*	*	3	*	11	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.59D Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	3.42	5.63	*	*	3.97	5.85	4.57	*
Bought from Friend or Relative	4.18	*	3.95	*	3.29	2.77	5.60	*
Took from Friend or Relative without Asking	1.45	1.74	3.03	*	0.98	*	2.17	*
Bought from Drug Dealer or Other Stranger	2.47	*	3.71	*	3.16	4.26	3.43	*
Bought on the Internet	0.49	0.43	*	*	0.66	1.61	*	*
Some Other Way	1.29	1.31	*	*	1.02	*	1.79	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.59P Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	0.9059		*		0.8140		*	
Bought from Friend or Relative	*		*		0.0047		*	
Took from Friend or Relative without Asking	0.7996		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		0.7897		*	
Bought on the Internet	0.9083		*		0.3432		*	
Some Other Way	0.7104		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	22	*	*	*	*	*	*	*
Bought from Friend or Relative	17	*	*	*	6	*	*	*
Took from Friend or Relative without Asking	6	1	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	10	*	*	*	4	*	*	*
Bought on the Internet	*	1	*	*	*	*	*	*
Some Other Way	3	5	*	*	3	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	4.84	*	*	*	*	*	*	*
Bought from Friend or Relative	4.45	*	*	*	4.99	*	*	*
Took from Friend or Relative without Asking	1.78	0.57	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3.02	*	*	*	3.87	*	*	*
Bought on the Internet	*	0.98	*	*	*	*	*	*
Some Other Way	1.00	3.57	*	*	2.65	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.1696		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.0233		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users<sup>1</sup></b>	<b>Total (2009-2010)</b>	<b>Total (2011-2012)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 12-17 (2011-2012)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 18-25 (2011-2012)</b>	<b>Aged 26+ (2009-2010)</b>	<b>Aged 26+ (2011-2012)</b>
From Friend or Relative for Free	23	*	*	*	*	*	*	*
Bought from Friend or Relative	26	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	21	*	*	*	*	*	*	*
Bought on the Internet	3	*	*	*	*	*	*	*
Some Other Way	9	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users<sup>1</sup></b>	<b>Total (2009-2010)</b>	<b>Total (2011-2012)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 12-17 (2011-2012)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 18-25 (2011-2012)</b>	<b>Aged 26+ (2009-2010)</b>	<b>Aged 26+ (2011-2012)</b>
From Friend or Relative for Free	5.03	*	*	*	*	*	*	*
Bought from Friend or Relative	5.19	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4.63	*	*	*	*	*	*	*
Bought on the Internet	0.77	*	*	*	*	*	*	*
Some Other Way	2.51	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users<sup>1</sup></b>	<b>Total (2009-2010)</b>	<b>Total (2011-2012)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 12-17 (2011-2012)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 18-25 (2011-2012)</b>	<b>Aged 26+ (2009-2010)</b>	<b>Aged 26+ (2011-2012)</b>
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	21	27	4	3	6	4	20	*
From More Than One Doctor	11	4	1	*	2	2	*	*
Wrote Fake Prescription	1	1	0	*	0	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	2	*	2	*	1	*	*
From Friend or Relative for Free	53	31	7	5	10	9	52	*
Bought from Friend or Relative	17	10	4	2	5	4	16	*
Took from Friend or Relative without Asking	11	5	3	2	6	*	9	*
Bought from Drug Dealer or Other Stranger	10	4	1	1	2	2	10	*
Bought on the Internet	*	0	*	0	*	0	*	*
Some Other Way	13	10	2	0	2	2	13	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	2.50	4.36	4.07	4.71	3.43	3.62	3.37	*
From More Than One Doctor	1.39	0.81	1.05	*	1.57	1.38	*	*
Wrote Fake Prescription	0.07	0.19	0.44	*	0.28	0.23	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.47	0.45	*	2.66	*	1.00	*	*
From Friend or Relative for Free	3.82	4.21	5.36	5.71	5.01	5.30	5.13	*
Bought from Friend or Relative	2.03	2.01	3.69	3.36	3.15	3.59	2.73	*
Took from Friend or Relative without Asking	1.42	0.92	3.22	2.84	3.89	*	1.54	*
Bought from Drug Dealer or Other Stranger	1.27	0.72	1.14	1.56	1.17	2.05	1.78	*
Bought on the Internet	*	0.03	*	0.06	*	0.15	*	*
Some Other Way	1.59	1.90	1.98	0.59	1.21	2.07	2.23	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.0429		0.3049		0.2156		*	
From More Than One Doctor	0.6266		*		0.9207		*	
Wrote Fake Prescription	0.4433		*		0.8923		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5338		*		*		*	
From Friend or Relative for Free	0.4427		0.4985		0.2696		*	
Bought from Friend or Relative	0.7563		0.2440		0.7424		*	
Took from Friend or Relative without Asking	0.2964		0.6594		*		*	
Bought from Drug Dealer or Other Stranger	0.3232		0.6943		0.1291		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.8374		0.1722		0.7197		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	51	29	*	*	*	*	*	*
From More Than One Doctor	3	3	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	*	*	*	*	*	*	*
From Friend or Relative for Free	11	4	*	*	*	*	*	*
Bought from Friend or Relative	2	1	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4	3	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	2	1	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	3.29	2.72	*	*	*	*	*	*
From More Than One Doctor	0.64	1.27	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.54	*	*	*	*	*	*	*
From Friend or Relative for Free	2.42	1.71	*	*	*	*	*	*
Bought from Friend or Relative	0.56	0.56	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.66	1.12	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1.02	1.05	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.43	0.31	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	0.8719		*		*		*	
From More Than One Doctor	0.1988		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.7597		*		*		*	
Bought from Friend or Relative	0.8463		*		*		*	
Took from Friend or Relative without Asking	0.3979		*		*		*	
Bought from Drug Dealer or Other Stranger	0.9483		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.7813		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Sedative Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	19	*	*	*	*	*	*	*
From More Than One Doctor	5	*	*	*	*	*	*	*
Wrote Fake Prescription	0	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	1	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	6	4	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	0	0	*	*	*	*	*	*
Some Other Way	9	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source for Past Month Nonmedical Use among Past Month Sedative Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	5.11	*	*	*	*	*	*	*
From More Than One Doctor	1.31	*	*	*	*	*	*	*
Wrote Fake Prescription	0.13	0.43	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.09	0.61	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.72	1.96	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	0.03	0.02	*	*	*	*	*	*
Some Other Way	2.46	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source for Past Month Nonmedical Use among Past Month Sedative Users <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	0.4191		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6596		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.7854		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	0.7797		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

**Table 6.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2009-2010)	Total (2011-2012)	Aged 12-17 (2009-2010)	Aged 12-17 (2011-2012)	Aged 18-25 (2009-2010)	Aged 18-25 (2011-2012)	Aged 26+ (2009-2010)	Aged 26+ (2011-2012)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.