

Table 6.71C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
ILLCIT DRUGS²	139	151	11	19	139	148
Marijuana and Hashish	117	122	10	19	116	119
Cocaine	37	30	2	2	37	30
Crack	23	12	1	*	23	12
Heroin	14	16	2	3	13	16
Hallucinogens	34	26	4	1	33	26
LSD	8	6	4	*	7	6
PCP	7	*	4	0	6	*
Ecstasy	27	23	4	1	27	23
Inhalants	19	16	4	1	19	16
Nonmedical Use of Psychotherapeutics ^{3,4}	83	82	6	10	82	82
Pain Relievers	69	64	6	6	68	64
OxyContin [®]	20	13	2	1	20	14
Tranquilizers	43	50	6	3	42	50
Stimulants ⁴	36	30	4	7	35	29
Methamphetamine ⁴	22	22	4	6	22	21
Sedatives	12	13	4	0	11	13
ILLCIT DRUGS OTHER THAN MARIJUANA²	94	90	7	10	93	89

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.71D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
ILLCIT DRUGS²	0.21	0.22	0.47	0.86	0.22	0.22
Marijuana and Hashish	0.18	0.18	0.42	0.82	0.19	0.18
Cocaine	0.06	0.05	0.08	0.11	0.06	0.05
Crack	0.04	0.02	0.04	*	0.04	0.02
Heroin	0.02	0.03	0.09	0.14	0.02	0.03
Hallucinogens	0.05	0.04	0.17	0.05	0.06	0.04
LSD	0.01	0.01	0.16	*	0.01	0.01
PCP	0.01	*	0.16	0.01	0.01	*
Ecstasy	0.04	0.04	0.17	0.03	0.05	0.04
Inhalants	0.03	0.02	0.16	0.03	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.13	0.13	0.26	0.43	0.14	0.13
Pain Relievers	0.11	0.10	0.24	0.28	0.11	0.10
OxyContin [®]	0.03	0.02	0.06	0.07	0.03	0.02
Tranquilizers	0.07	0.08	0.23	0.15	0.07	0.08
Stimulants ⁴	0.06	0.05	0.17	0.30	0.06	0.05
Methamphetamine ⁴	0.04	0.04	0.16	0.29	0.04	0.04
Sedatives	0.02	0.02	0.16	0.01	0.02	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.15	0.14	0.29	0.44	0.15	0.14

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.71P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
ILLCIT DRUGS²	0.6507		0.1216		0.4930	
Marijuana and Hashish	0.4195		0.0839		0.6117	
Cocaine	0.2799		0.8102		0.2764	
Crack	0.0361		*		0.0410	
Heroin	0.5198		0.8713		0.4640	
Hallucinogens	0.0098		0.2260		0.0124	
LSD	0.4301		*		0.7506	
PCP	*		0.3399		*	
Ecstasy	0.0422		0.1399		0.0716	
Inhalants	0.0719		0.2257		0.1183	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.2040		0.4403		0.1773	
Pain Relievers	0.0228		0.9049		0.0252	
OxyContin [®]	0.0007		0.9615		0.0007	
Tranquilizers	0.8205		0.3985		0.7419	
Stimulants ⁴	0.0733		0.6820		0.0770	
Methamphetamine ⁴	0.9477		0.8002		0.9338	
Sedatives	0.8161		0.3565		0.9636	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.0166		0.7124		0.0150	

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.72C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	139	151	11	19	139	148
AGE						
15-17	30	31	3	3	29	30
18-25	57	65	8	12	66	81
26-44	118	115	8	15	118	112
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	130	137	10	17	130	135
White	112	119	8	14	112	117
Black or African American	56	57	6	10	55	54
American Indian or Alaska Native	6	10	*	*	6	10
Native Hawaiian or Other Pacific Islander	3	9	*	*	3	9
Asian	17	29	*	*	17	29
Two or More Races	14	17	*	*	14	17
Hispanic or Latino	52	61	6	9	52	60
TRIMESTER²						
First	N/A	N/A	9	16	N/A	N/A
Second	N/A	N/A	6	9	N/A	N/A
Third	N/A	N/A	4	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.72D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.21	0.22	0.47	0.86	0.22	0.22
AGE						
15-17	0.43	0.44	3.40	4.44	0.44	0.44
18-25	0.34	0.38	0.85	1.42	0.35	0.40
26-44	0.29	0.28	0.53	1.10	0.30	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.24	0.24	0.51	0.97	0.25	0.25
White	0.29	0.29	0.55	1.13	0.30	0.30
Black or African American	0.60	0.57	1.82	2.74	0.61	0.57
American Indian or Alaska Native	1.88	2.21	*	*	1.94	2.29
Native Hawaiian or Other Pacific Islander	1.23	3.49	*	*	1.24	3.68
Asian	0.53	0.73	*	*	0.56	0.76
Two or More Races	1.62	1.46	*	*	1.70	1.51
Hispanic or Latino	0.45	0.48	1.24	1.89	0.47	0.49
TRIMESTER²						
First	N/A	N/A	1.27	2.19	N/A	N/A
Second	N/A	N/A	0.64	1.15	N/A	N/A
Third	N/A	N/A	0.47	0.63	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.72P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.6507		0.1216		0.4930	
AGE						
15-17	0.8681		0.6359		0.7954	
18-25	0.3539		0.3285		0.5573	
26-44	0.2444		0.2094		0.1919	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4114		0.1375		0.3140	
White	0.0967		0.1749		0.0580	
Black or African American	0.5084		0.7157		0.4474	
American Indian or Alaska Native	0.9195		*		0.7478	
Native Hawaiian or Other Pacific Islander	0.0222		*		0.0211	
Asian	0.0660		*		0.0817	
Two or More Races	0.7783		*		0.8141	
Hispanic or Latino	0.2563		0.6549		0.3269	
TRIMESTER²						
First	N/A		0.1633		N/A	
Second	N/A		0.7169		N/A	
Third	N/A		0.7154		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.73C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	117	122	10	19	116	119
AGE						
15-17	26	27	3	3	25	27
18-25	53	60	8	11	60	73
26-44	98	92	6	15	98	90
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	111	113	9	17	110	110
White	95	97	8	14	95	96
Black or African American	49	51	5	10	48	48
American Indian or Alaska Native	5	6	*	*	5	6
Native Hawaiian or Other Pacific Islander	3	7	*	*	3	7
Asian	12	22	*	*	12	22
Two or More Races	13	16	*	*	13	16
Hispanic or Latino	39	46	5	8	39	45
TRIMESTER²						
First	N/A	N/A	8	16	N/A	N/A
Second	N/A	N/A	5	8	N/A	N/A
Third	N/A	N/A	3	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.73D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.18	0.18	0.42	0.82	0.19	0.18
AGE						
15-17	0.38	0.39	3.28	4.28	0.39	0.39
18-25	0.31	0.35	0.82	1.31	0.33	0.36
26-44	0.25	0.23	0.42	1.09	0.26	0.23
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.21	0.20	0.44	0.96	0.22	0.21
White	0.25	0.25	0.50	1.12	0.26	0.25
Black or African American	0.52	0.52	1.51	2.71	0.54	0.51
American Indian or Alaska Native	1.54	1.38	*	*	1.60	1.44
Native Hawaiian or Other Pacific Islander	1.16	2.88	*	*	1.17	3.04
Asian	0.37	0.57	*	*	0.39	0.59
Two or More Races	1.56	1.39	*	*	1.63	1.45
Hispanic or Latino	0.36	0.36	1.20	1.64	0.37	0.37
TRIMESTER²						
First	N/A	N/A	1.14	2.17	N/A	N/A
Second	N/A	N/A	0.57	1.00	N/A	N/A
Third	N/A	N/A	0.44	0.63	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.73P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.4195		0.0839		0.6117	
AGE						
15-17	0.1388		0.6660		0.1239	
18-25	0.1086		0.5293		0.1761	
26-44	0.6372		0.0785		0.4784	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6271		0.0575		0.8303	
White	0.8362		0.0995		0.6393	
Black or African American	0.5500		0.5575		0.5769	
American Indian or Alaska Native	0.9554		*		0.8399	
Native Hawaiian or Other Pacific Islander	0.0492		*		0.0467	
Asian	0.0133		*		0.0159	
Two or More Races	0.8095		*		0.8746	
Hispanic or Latino	0.1428		0.9703		0.1723	
TRIMESTER²						
First	N/A		0.0910		N/A	
Second	N/A		0.9943		N/A	
Third	N/A		0.4709		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.74C Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOBACCO PRODUCTS²	249	271	32	32	246	265
Cigarettes	245	259	31	31	242	254
Smokeless Tobacco	31	28	4	6	31	28
Cigars	78	86	6	8	78	86
Pipe Tobacco	22	28	4	2	21	27
ALCOHOL	407	459	27	21	401	459
Binge Alcohol Use ³	238	259	14	10	237	258
Heavy Alcohol Use ³	104	118	6	3	103	118

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.74D Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOBACCO PRODUCTS²	0.35	0.37	1.19	1.36	0.36	0.37
Cigarettes	0.35	0.36	1.18	1.34	0.36	0.36
Smokeless Tobacco	0.05	0.05	0.17	0.28	0.05	0.05
Cigars	0.13	0.13	0.24	0.38	0.13	0.14
Pipe Tobacco	0.04	0.04	0.16	0.10	0.04	0.05
ALCOHOL	0.39	0.41	1.02	0.93	0.40	0.42
Binge Alcohol Use ³	0.32	0.32	0.55	0.46	0.34	0.34
Heavy Alcohol Use ³	0.16	0.18	0.25	0.13	0.17	0.19

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.74P Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOBACCO PRODUCTS²	0.0003		0.9389		0.0001	
Cigarettes	0.0000		0.8667		0.0000	
Smokeless Tobacco	0.2707		0.5697		0.2452	
Cigars	0.6384		0.8520		0.5501	
Pipe Tobacco	0.1311		0.9662		0.1239	
ALCOHOL	0.1313		0.1210		0.2134	
Binge Alcohol Use ³	0.8300		0.1880		0.9704	
Heavy Alcohol Use ³	0.4474		0.0179		0.4262	

*Low precision; no estimate reported.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.75C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	245	259	31	31	242	254
AGE						
15-17	29	25	4	*	28	24
18-25	69	72	16	16	88	92
26-44	217	227	27	25	216	223
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	229	236	31	31	225	230
White	210	208	29	26	206	204
Black or African American	88	80	8	11	87	78
American Indian or Alaska Native	16	21	*	*	16	20
Native Hawaiian or Other Pacific Islander	7	*	*	*	6	*
Asian	39	39	*	*	39	39
Two or More Races	23	36	*	*	23	35
Hispanic or Latino	94	94	6	5	94	94
TRIMESTER²						
First	N/A	N/A	21	23	N/A	N/A
Second	N/A	N/A	17	16	N/A	N/A
Third	N/A	N/A	15	12	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.75D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.35	0.36	1.18	1.34	0.36	0.36
AGE						
15-17	0.42	0.37	4.29	*	0.42	0.36
18-25	0.41	0.42	1.48	1.69	0.43	0.43
26-44	0.51	0.51	1.74	1.82	0.52	0.52
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.40	0.39	1.39	1.63	0.41	0.40
White	0.48	0.47	1.68	1.96	0.49	0.48
Black or African American	0.86	0.77	2.56	3.14	0.88	0.78
American Indian or Alaska Native	3.74	3.58	*	*	3.92	3.59
Native Hawaiian or Other Pacific Islander	2.72	*	*	*	2.46	*
Asian	1.15	0.97	*	*	1.20	1.00
Two or More Races	2.37	2.45	*	*	2.47	2.50
Hispanic or Latino	0.77	0.72	1.35	1.10	0.81	0.75
TRIMESTER²						
First	N/A	N/A	2.47	2.85	N/A	N/A
Second	N/A	N/A	1.82	1.97	N/A	N/A
Third	N/A	N/A	1.89	1.72	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.75P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.0000		0.8667		0.0000	
AGE						
15-17	0.0000		*		0.0000	
18-25	0.0000		0.4228		0.0000	
26-44	0.0234		0.7505		0.0125	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0008		0.7865		0.0004	
White	0.0036		0.8556		0.0024	
Black or African American	0.1150		0.5202		0.0939	
American Indian or Alaska Native	0.2404		*		0.3701	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4901		*		0.4145	
Two or More Races	0.9930		*		0.8411	
Hispanic or Latino	0.0226		0.2354		0.0212	
TRIMESTER²						
First	N/A		0.7975		N/A	
Second	N/A		0.9838		N/A	
Third	N/A		0.7480		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.76C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	407	459	27	21	401	459
AGE						
15-17	38	36	2	3	38	36
18-25	81	88	11	7	156	185
26-44	352	393	25	20	347	394
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	375	410	23	19	369	410
White	330	358	21	17	325	358
Black or African American	137	165	8	8	136	164
American Indian or Alaska Native	16	21	*	*	16	21
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	90	95	*	*	89	95
Two or More Races	32	47	*	*	32	46
Hispanic or Latino	145	182	14	7	143	181
TRIMESTER²						
First	N/A	N/A	19	17	N/A	N/A
Second	N/A	N/A	17	8	N/A	N/A
Third	N/A	N/A	8	9	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.76D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.39	0.41	1.02	0.93	0.40	0.42
AGE						
15-17	0.51	0.50	2.96	4.11	0.52	0.51
18-25	0.48	0.51	1.13	0.89	0.49	0.53
26-44	0.56	0.59	1.60	1.46	0.58	0.61
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.40	0.44	1.09	1.08	0.41	0.45
White	0.46	0.48	1.29	1.34	0.47	0.48
Black or African American	0.97	1.10	2.39	2.18	1.00	1.15
American Indian or Alaska Native	3.61	3.45	*	*	3.81	3.43
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.92	1.96	*	*	2.04	1.99
Two or More Races	2.59	2.52	*	*	2.64	2.58
Hispanic or Latino	0.98	1.06	2.90	1.48	1.02	1.10
TRIMESTER²						
First	N/A	N/A	2.30	2.30	N/A	N/A
Second	N/A	N/A	1.76	0.97	N/A	N/A
Third	N/A	N/A	0.98	1.19	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.76P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.1313		0.1210		0.2134	
AGE						
15-17	0.4560		0.8957		0.4375	
18-25	0.4505		0.1949		0.7374	
26-44	0.1410		0.2498		0.1865	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1568		0.3131		0.2696	
White	0.0646		0.5119		0.1435	
Black or African American	0.5520		0.3033		0.3901	
American Indian or Alaska Native	0.1920		*		0.2138	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7717		*		0.6521	
Two or More Races	0.1551		*		0.3128	
Hispanic or Latino	0.1227		0.1839		0.1184	
TRIMESTER²						
First	N/A		0.5643		N/A	
Second	N/A		0.0093		N/A	
Third	N/A		0.6484		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.77C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	238	259	14	10	237	258
AGE						
15-17	29	24	2	*	29	24
18-25	76	83	6	4	105	129
26-44	203	214	12	9	202	214
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	211	229	10	10	210	228
White	191	204	10	8	190	203
Black or African American	78	88	4	3	78	87
American Indian or Alaska Native	12	17	*	*	12	17
Native Hawaiian or Other Pacific Islander	10	15	*	*	10	*
Asian	41	48	*	*	41	49
Two or More Races	19	27	*	*	19	27
Hispanic or Latino	103	109	9	3	102	109
TRIMESTER²						
First	N/A	N/A	13	9	N/A	N/A
Second	N/A	N/A	4	3	N/A	N/A
Third	N/A	N/A	0	2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.77D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.32	0.32	0.55	0.46	0.34	0.34
AGE						
15-17	0.43	0.36	2.71	*	0.44	0.37
18-25	0.45	0.48	0.67	0.55	0.47	0.50
26-44	0.46	0.46	0.84	0.67	0.48	0.47
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.34	0.36	0.51	0.56	0.36	0.37
White	0.42	0.42	0.63	0.67	0.44	0.43
Black or African American	0.78	0.84	1.22	0.86	0.81	0.88
American Indian or Alaska Native	3.17	3.22	*	*	3.33	3.28
Native Hawaiian or Other Pacific Islander	3.81	5.02	*	*	3.89	*
Asian	1.23	1.21	*	*	1.27	1.24
Two or More Races	2.14	2.16	*	*	2.24	2.23
Hispanic or Latino	0.80	0.76	1.98	0.71	0.85	0.79
TRIMESTER²						
First	N/A	N/A	1.67	1.32	N/A	N/A
Second	N/A	N/A	0.45	0.44	N/A	N/A
Third	N/A	N/A	0.05	0.22	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.77P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.8300		0.1880		0.9704	
AGE						
15-17	0.0288		*		0.0421	
18-25	0.9928		0.1654		0.8795	
26-44	0.6307		0.5624		0.7592	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6487		0.5091		0.8252	
White	0.8198		0.3815		0.9460	
Black or African American	0.3506		0.3554		0.2584	
American Indian or Alaska Native	0.3877		*		0.4307	
Native Hawaiian or Other Pacific Islander	0.3873		*		*	
Asian	0.4804		*		0.5751	
Two or More Races	0.8469		*		0.6691	
Hispanic or Latino	0.9171		0.1831		0.9729	
TRIMESTER²						
First	N/A		0.1051		N/A	
Second	N/A		0.8270		N/A	
Third	N/A		0.1706		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.78C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	104	118	6	3	103	118
AGE						
15-17	12	11	1	*	12	11
18-25	42	52	5	2	46	59
26-44	91	99	4	2	91	99
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	97	105	5	3	97	105
White	88	94	4	3	87	93
Black or African American	33	37	2	0	33	37
American Indian or Alaska Native	6	12	*	*	6	12
Native Hawaiian or Other Pacific Islander	3	*	*	*	3	*
Asian	14	17	*	*	14	17
Two or More Races	6	9	*	*	6	9
Hispanic or Latino	33	47	4	1	33	47
TRIMESTER²						
First	N/A	N/A	6	3	N/A	N/A
Second	N/A	N/A	3	0	N/A	N/A
Third	N/A	N/A	*	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.78D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2009-2010 and 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.16	0.18	0.25	0.13	0.17	0.19
AGE						
15-17	0.19	0.17	1.87	*	0.19	0.17
18-25	0.25	0.30	0.48	0.21	0.26	0.31
26-44	0.23	0.25	0.29	0.18	0.24	0.25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.18	0.20	0.24	0.15	0.19	0.20
White	0.22	0.24	0.29	0.21	0.23	0.25
Black or African American	0.38	0.42	0.66	0.15	0.39	0.44
American Indian or Alaska Native	1.69	2.72	*	*	1.77	2.82
Native Hawaiian or Other Pacific Islander	1.12	*	*	*	1.14	*
Asian	0.41	0.43	*	*	0.43	0.44
Two or More Races	0.78	0.80	*	*	0.83	0.84
Hispanic or Latino	0.31	0.38	0.91	0.22	0.32	0.40
TRIMESTER²						
First	N/A	N/A	0.76	0.41	N/A	N/A
Second	N/A	N/A	0.32	0.04	N/A	N/A
Third	N/A	N/A	*	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.

Table 6.78P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011-2012 Versus 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2009-2010	2011-2012	2009-2010	2011-2012	2009-2010	2011-2012
TOTAL	0.4474		0.0179		0.4262	
AGE						
15-17	0.0047		*		0.0126	
18-25	0.1228		0.0450		0.1248	
26-44	0.8599		0.5561		0.9396	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5794		0.0415		0.5344	
White	0.3933		0.1382		0.3292	
Black or African American	0.8919		0.0846		0.9827	
American Indian or Alaska Native	0.2364		*		0.2479	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3402		*		0.3672	
Two or More Races	0.6165		*		0.6451	
Hispanic or Latino	0.8176		0.2266		0.8752	
TRIMESTER²						
First	N/A		0.0282		N/A	
Second	N/A		0.2605		N/A	
Third	N/A		*		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2009-2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009-2012.