

Table 7.1A Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|---------|
| ILLCIT DRUGS¹ | 108,255 ^b | 110,205 ^b | 110,057 ^b | 112,085 ^b | 111,747 ^b | 114,509 ^b | 117,815 ^b | 119,259 ^b | 119,933 ^b | 121,078 ^b | 124,808 |
| Marijuana and Hashish | 94,946 ^b | 96,611 ^b | 96,772 ^b | 97,545 ^b | 97,792 ^b | 100,737 ^b | 102,857 ^b | 104,950 ^b | 106,613 ^b | 107,842 ^a | 111,239 |
| Cocaine | 33,910 ^b | 34,891 ^b | 34,153 ^b | 33,673 ^b | 35,303 ^a | 35,947 | 36,916 | 36,742 | 37,361 | 36,921 | 37,688 |
| Crack | 8,402 | 7,949 ^a | 7,840 ^b | 7,928 ^a | 8,559 | 8,615 | 8,454 | 8,390 | 9,208 | 8,214 | 9,015 |
| Heroin | 3,668 ^b | 3,744 ^a | 3,145 ^b | 3,534 ^b | 3,788 ^a | 3,806 ^a | 3,795 ^a | 3,680 ^b | 4,144 | 4,162 | 4,565 |
| Hallucinogens | 34,314 ^b | 34,363 ^b | 34,333 ^b | 33,728 ^b | 35,290 ^b | 34,278 ^b | 36,064 | 37,404 | 37,544 | 36,362 | 37,908 |
| LSD | 24,516 | 24,424 | 23,398 | 22,433 | 23,365 | 22,720 | 23,604 | 23,714 | 23,375 | 23,000 | 23,735 |
| PCP | 7,418 ^a | 7,107 | 6,762 | 6,603 | 6,633 | 6,179 | 6,657 | 6,271 | 6,255 | 6,103 | 6,552 |
| Ecstasy | 10,150 ^b | 10,904 ^b | 11,130 ^b | 11,495 ^b | 12,282 ^b | 12,418 ^b | 12,964 ^b | 14,280 ^b | 15,929 | 14,570 ^b | 16,162 |
| Inhalants | 22,870 ^b | 22,995 ^b | 22,798 ^a | 22,745 ^a | 22,856 ^b | 22,558 ^a | 22,430 | 22,552 ^a | 21,778 | 20,523 | 21,095 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 47,958 ^b | 49,001 ^b | 49,157 ^b | 49,571 ^b | 50,983 ^b | 50,541 ^b | 52,154 ^a | 51,991 ^a | 51,832 ^a | 51,243 ^b | 54,389 |
| Pain Relievers | 29,611 ^b | 31,207 ^b | 31,768 ^b | 32,692 ^b | 33,478 ^b | 33,132 ^b | 34,992 ^a | 35,197 ^a | 34,908 ^a | 34,247 ^b | 37,045 |
| OxyContin [®] | 1,924 ^b | 2,832 ^b | 3,072 ^b | 3,481 ^b | 4,116 ^b | 4,379 ^b | 4,857 ^b | 5,863 ^a | 6,160 | 5,917 ^a | 6,579 |
| Tranquilizers | 19,267 ^b | 20,220 ^b | 19,852 ^b | 21,041 ^b | 21,318 ^b | 20,253 ^b | 21,551 ^b | 21,848 ^a | 22,187 | 21,655 ^b | 23,639 |
| Stimulants ³ | 23,496 ^b | 23,004 ^a | 22,297 | 20,983 | 22,495 | 21,669 | 21,285 | 22,033 | 21,739 | 20,379 | 21,498 |
| Methamphetamine ³ | 15,365 ^b | 15,139 ^b | 14,512 ^b | 12,663 | 14,226 ^b | 13,081 | 12,634 | 12,908 | 13,060 | 11,928 | 12,259 |
| Sedatives | 9,960 ^b | 9,510 ^b | 9,891 ^b | 8,982 ^a | 8,842 | 8,401 | 8,920 | 8,633 | 7,653 | 7,515 | 7,939 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 70,300 ^b | 71,128 ^b | 70,657 ^b | 71,822 ^b | 72,934 ^b | 73,654 ^b | 75,907 | 76,080 | 76,472 | 75,447 ^a | 78,034 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.1B Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|-------------------|------|
| ILLCIT DRUGS¹ | 46.0 ^b | 46.4 ^b | 45.8 ^b | 46.1 ^b | 45.4 ^b | 46.2 ^b | 47.2 | 47.4 | 47.3 | 47.0 | 48.0 |
| Marijuana and Hashish | 40.4 ^b | 40.6 ^b | 40.2 ^b | 40.1 ^b | 39.7 ^b | 40.6 ^b | 41.2 ^b | 41.7 ^a | 42.0 | 41.9 | 42.8 |
| Cocaine | 14.4 | 14.7 | 14.2 | 13.8 | 14.3 | 14.5 | 14.8 | 14.6 | 14.7 | 14.3 | 14.5 |
| Crack | 3.6 | 3.3 | 3.3 | 3.3 | 3.5 | 3.5 | 3.4 | 3.3 | 3.6 | 3.2 | 3.5 |
| Heroin | 1.6 | 1.6 | 1.3 ^b | 1.5 ^a | 1.5 | 1.5 | 1.5 | 1.5 ^a | 1.6 | 1.6 | 1.8 |
| Hallucinogens | 14.6 | 14.5 | 14.3 | 13.9 ^a | 14.3 | 13.8 ^a | 14.4 | 14.9 | 14.8 | 14.1 | 14.6 |
| LSD | 10.4 ^b | 10.3 ^b | 9.7 ^a | 9.2 | 9.5 | 9.2 | 9.4 | 9.4 | 9.2 | 8.9 | 9.1 |
| PCP | 3.2 ^b | 3.0 ^b | 2.8 | 2.7 | 2.7 | 2.5 | 2.7 | 2.5 | 2.5 | 2.4 | 2.5 |
| Ecstasy | 4.3 ^b | 4.6 ^b | 4.6 ^b | 4.7 ^b | 5.0 ^b | 5.0 ^b | 5.2 ^b | 5.7 ^a | 6.3 | 5.7 ^b | 6.2 |
| Inhalants | 9.7 ^b | 9.7 ^b | 9.5 ^b | 9.4 ^b | 9.3 ^b | 9.1 ^b | 9.0 ^b | 9.0 ^b | 8.6 | 8.0 | 8.1 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 20.4 | 20.6 | 20.4 | 20.4 | 20.7 | 20.4 | 20.9 | 20.6 | 20.4 | 19.9 ^a | 20.9 |
| Pain Relievers | 12.6 ^b | 13.1 ^b | 13.2 ^b | 13.4 ^a | 13.6 ^a | 13.4 ^b | 14.0 | 14.0 | 13.8 | 13.3 ^b | 14.2 |
| OxyContin [®] | 0.8 ^b | 1.2 ^b | 1.3 ^b | 1.4 ^b | 1.7 ^b | 1.8 ^b | 1.9 ^b | 2.3 | 2.4 | 2.3 | 2.5 |
| Tranquilizers | 8.2 ^b | 8.5 ^a | 8.3 ^b | 8.7 | 8.7 | 8.2 ^b | 8.6 | 8.7 | 8.7 | 8.4 ^a | 9.1 |
| Stimulants ³ | 10.0 ^b | 9.7 ^b | 9.3 ^b | 8.6 | 9.1 ^b | 8.7 | 8.5 | 8.7 | 8.6 | 7.9 | 8.3 |
| Methamphetamine ³ | 6.5 ^b | 6.4 ^b | 6.0 ^b | 5.2 ^a | 5.8 ^b | 5.3 ^a | 5.1 | 5.1 | 5.1 | 4.6 | 4.7 |
| Sedatives | 4.2 ^b | 4.0 ^b | 4.1 ^b | 3.7 ^b | 3.6 ^b | 3.4 | 3.6 ^a | 3.4 | 3.0 | 2.9 | 3.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 29.9 | 29.9 | 29.4 | 29.5 | 29.6 | 29.7 | 30.4 | 30.2 | 30.2 | 29.3 | 30.0 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.2A Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 35,132 ^b | 34,993 ^b | 34,807 ^b | 35,041 ^b | 35,810 ^b | 35,789 ^b | 35,643 ^b | 38,123 ^b | 38,908 ^b | 38,287 ^b | 41,479 |
| Marijuana and Hashish | 25,755 ^b | 25,231 ^b | 25,451 ^b | 25,375 ^b | 25,432 ^b | 25,136 ^b | 25,891 ^b | 28,688 ^b | 29,301 ^b | 29,739 ^a | 31,513 |
| Cocaine | 5,902 ^b | 5,908 ^b | 5,658 ^b | 5,523 ^b | 6,075 ^b | 5,748 ^b | 5,283 ^a | 4,806 | 4,533 | 3,857 ^b | 4,671 |
| Crack | 1,554 ^b | 1,406 ^b | 1,304 ^a | 1,381 ^b | 1,478 ^b | 1,452 ^b | 1,113 | 1,008 | 885 | 625 ^a | 921 |
| Heroin | 404 ^b | 314 ^b | 398 ^b | 379 ^b | 560 | 373 ^b | 455 ^a | 582 | 621 | 620 | 669 |
| Hallucinogens | 4,749 | 3,936 | 3,878 | 3,809 ^a | 3,970 | 3,759 ^a | 3,670 ^b | 4,536 | 4,543 | 4,069 | 4,306 |
| LSD | 999 | 558 ^b | 592 ^b | 563 ^b | 667 ^b | 621 ^b | 808 ^a | 790 ^b | 881 | 880 | 1,057 |
| PCP | 235 | 219 | 210 | 164 | 187 | 139 | 99 ^a | 125 | 96 | 119 | 172 |
| Ecstasy | 3,167 ^b | 2,119 ^b | 1,915 ^b | 1,960 ^b | 2,138 ^b | 2,122 ^b | 2,116 ^b | 2,804 | 2,653 | 2,422 | 2,610 |
| Inhalants | 2,084 ^a | 2,075 ^b | 2,255 ^b | 2,187 ^b | 2,219 ^b | 2,085 ^a | 2,065 ^a | 2,109 ^a | 2,048 ^a | 1,861 | 1,693 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 14,795 ^b | 15,163 ^b | 14,849 ^b | 15,346 ^a | 16,482 | 16,334 | 15,190 ^b | 16,064 | 16,051 | 14,657 ^b | 16,666 |
| Pain Relievers | 10,992 ^b | 11,671 | 11,256 ^b | 11,815 | 12,650 | 12,517 | 11,916 | 12,450 | 12,242 | 11,143 ^b | 12,489 |
| OxyContin [®] | -- | -- | 1,213 ^a | 1,226 | 1,324 | 1,427 | 1,462 | 1,685 | 1,879 ^b | 1,623 | 1,477 |
| Tranquilizers | 4,849 ^b | 5,051 ^b | 5,068 ^b | 5,249 ^a | 5,066 ^b | 5,293 ^a | 5,099 ^b | 5,493 | 5,583 | 5,109 ^b | 6,073 |
| Stimulants ³ | 3,380 | 3,031 | 3,254 | 3,088 | 3,789 | 2,994 | 2,648 ^b | 3,093 | 2,902 | 2,700 ^b | 3,317 |
| Methamphetamine ³ | 1,755 ^b | 1,602 ^b | 1,808 ^b | 1,603 ^b | 1,885 ^b | 1,345 | 854 ^a | 1,167 | 959 | 1,033 | 1,155 |
| Sedatives | 981 ^b | 831 ^a | 737 | 750 | 927 ^b | 864 ^a | 626 | 816 ^a | 906 ^a | 526 | 590 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 20,423 | 20,305 | 19,658 ^b | 20,109 | 21,262 | 21,205 | 20,018 ^a | 21,077 | 20,625 | 18,959 ^b | 21,267 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.2B Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|-------------------|------|
| ILLCIT DRUGS¹ | 14.9 ^b | 14.7 ^b | 14.5 ^b | 14.4 ^b | 14.6 ^b | 14.4 ^b | 14.3 ^b | 15.1 ^a | 15.3 | 14.9 ^b | 16.0 |
| Marijuana and Hashish | 11.0 ^b | 10.6 ^b | 10.6 ^b | 10.4 ^b | 10.3 ^b | 10.1 ^b | 10.4 ^b | 11.4 ^a | 11.6 | 11.5 ^a | 12.1 |
| Cocaine | 2.5 ^b | 2.5 ^b | 2.4 ^b | 2.3 ^b | 2.5 ^b | 2.3 ^b | 2.1 ^b | 1.9 | 1.8 | 1.5 ^b | 1.8 |
| Crack | 0.7 ^b | 0.6 ^b | 0.5 ^b | 0.6 ^b | 0.6 ^b | 0.6 ^b | 0.4 | 0.4 | 0.3 | 0.2 ^a | 0.4 |
| Heroin | 0.2 ^a | 0.1 ^b | 0.2 ^a | 0.2 ^b | 0.2 | 0.2 ^b | 0.2 ^a | 0.2 | 0.2 | 0.2 | 0.3 |
| Hallucinogens | 2.0 ^b | 1.7 | 1.6 | 1.6 | 1.6 | 1.5 | 1.5 ^a | 1.8 | 1.8 | 1.6 | 1.7 |
| LSD | 0.4 | 0.2 ^b | 0.2 ^b | 0.2 ^b | 0.3 ^b | 0.3 ^b | 0.3 ^a | 0.3 ^a | 0.3 | 0.3 | 0.4 |
| PCP | 0.1 ^a | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Ecstasy | 1.3 ^b | 0.9 | 0.8 ^b | 0.8 ^b | 0.9 | 0.9 ^a | 0.8 ^a | 1.1 | 1.0 | 0.9 | 1.0 |
| Inhalants | 0.9 ^b | 0.8 ^b | 0.8 ^b | 0.8 ^b | 0.8 ^a | 0.7 | 0.7 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 6.3 | 6.4 | 6.2 | 6.3 | 6.7 | 6.6 | 6.1 | 6.4 | 6.3 | 5.7 ^b | 6.4 |
| Pain Relievers | 4.7 | 4.9 | 4.7 | 4.9 | 5.1 | 5.1 | 4.8 | 4.9 | 4.8 | 4.3 ^b | 4.8 |
| OxyContin [®] | -- | -- | 0.5 | 0.5 | 0.5 | 0.6 | 0.6 | 0.7 | 0.7 ^b | 0.6 | 0.6 |
| Tranquilizers | 2.1 ^a | 2.1 | 2.1 | 2.2 | 2.1 ^a | 2.1 | 2.0 ^a | 2.2 | 2.2 | 2.0 ^b | 2.3 |
| Stimulants ³ | 1.4 | 1.3 | 1.4 | 1.3 | 1.5 ^b | 1.2 | 1.1 ^b | 1.2 | 1.1 | 1.0 ^b | 1.3 |
| Methamphetamine ³ | 0.7 ^b | 0.7 ^b | 0.8 ^b | 0.7 ^b | 0.8 ^b | 0.5 | 0.3 | 0.5 | 0.4 | 0.4 | 0.4 |
| Sedatives | 0.4 ^b | 0.3 ^b | 0.3 ^a | 0.3 ^a | 0.4 ^b | 0.3 ^b | 0.3 | 0.3 ^a | 0.4 ^b | 0.2 | 0.2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 8.7 ^a | 8.5 | 8.2 | 8.3 | 8.6 | 8.6 | 8.0 | 8.4 | 8.1 | 7.4 ^b | 8.2 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.3A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 19,522 ^b | 19,470 ^b | 19,071 ^b | 19,720 ^b | 20,387 ^b | 19,892 ^b | 20,123 ^b | 21,930 ^b | 22,648 | 22,454 ^a | 23,863 |
| Marijuana and Hashish | 14,584 ^b | 14,638 ^b | 14,576 ^b | 14,626 ^b | 14,849 ^b | 14,470 ^b | 15,269 ^b | 16,826 ^b | 17,409 ^a | 18,071 | 18,855 |
| Cocaine | 2,020 | 2,281 ^b | 2,021 ^a | 2,397 ^b | 2,426 ^b | 2,077 ^a | 1,865 | 1,642 | 1,472 | 1,369 | 1,650 |
| Crack | 567 | 604 | 467 | 682 | 704 | 611 | 363 | 497 | 378 | 228 ^a | 443 |
| Heroin | 166 ^a | 119 ^b | 166 ^b | 136 ^b | 339 | 161 ^b | 213 | 193 ^a | 239 | 281 | 335 |
| Hallucinogens | 1,196 | 1,042 | 929 | 1,088 | 1,008 | 998 | 1,059 | 1,263 | 1,207 | 972 | 1,134 |
| LSD | 112 ^a | 133 | 141 | 104 ^a | 130 | 144 | 157 | 159 | 155 | 152 | 184 |
| PCP | 58 | 56 | 49 | 48 | 31 | 43 | 25 | 54 | 36 | 26 | 32 |
| Ecstasy | 676 | 470 | 450 ^a | 502 | 529 | 503 | 546 | 763 | 691 | 544 | 628 |
| Inhalants | 635 | 570 | 638 | 611 | 762 ^b | 617 | 644 | 562 | 690 | 624 | 493 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 6,287 | 6,451 | 6,110 ^a | 6,491 | 7,098 | 6,907 | 6,210 | 6,980 | 6,957 | 6,119 ^a | 6,831 |
| Pain Relievers | 4,377 | 4,693 | 4,404 | 4,658 | 5,226 | 5,188 | 4,732 | 5,282 | 5,093 | 4,471 | 4,862 |
| OxyContin [®] | -- | -- | 325 | 334 | 275 | 371 | 436 | 510 ^a | 566 ^a | 434 | 358 |
| Tranquilizers | 1,804 | 1,830 | 1,616 ^b | 1,817 | 1,761 | 1,838 | 1,796 | 2,030 | 2,159 | 1,840 | 2,139 |
| Stimulants ³ | 1,303 | 1,310 | 1,312 | 1,188 | 1,385 | 1,051 | 908 ^a | 1,299 | 1,082 | 970 | 1,171 |
| Methamphetamine ³ | 683 ^a | 726 ^b | 706 ^a | 628 | 731 ^b | 530 | 314 | 502 | 353 | 439 | 440 |
| Sedatives | 436 ^a | 294 | 265 | 272 | 385 | 345 | 237 | 367 | 375 | 231 | 270 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 8,777 | 8,849 | 8,247 | 8,963 | 9,620 | 9,295 | 8,556 | 9,195 | 9,016 | 8,020 ^a | 8,883 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.3B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------|
| ILLCIT DRUGS¹ | 8.3 ^b | 8.2 ^b | 7.9 ^b | 8.1 ^b | 8.3 ^b | 8.0 ^b | 8.1 ^b | 8.7 | 8.9 | 8.7 | 9.2 |
| Marijuana and Hashish | 6.2 ^b | 6.2 ^b | 6.1 ^b | 6.0 ^b | 6.0 ^b | 5.8 ^b | 6.1 ^b | 6.7 ^a | 6.9 | 7.0 | 7.3 |
| Cocaine | 0.9 ^b | 1.0 ^b | 0.8 ^b | 1.0 ^b | 1.0 ^b | 0.8 ^a | 0.7 | 0.7 | 0.6 | 0.5 | 0.6 |
| Crack | 0.2 | 0.3 | 0.2 | 0.3 ^a | 0.3 ^a | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 ^a | 0.2 |
| Heroin | 0.1 ^a | 0.1 ^b | 0.1 ^a | 0.1 ^b | 0.1 | 0.1 ^a | 0.1 | 0.1 ^a | 0.1 | 0.1 | 0.1 |
| Hallucinogens | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | 0.4 | 0.4 |
| LSD | 0.0 | 0.1 | 0.1 | 0.0 ^a | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| PCP | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ecstasy | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 |
| Inhalants | 0.3 ^a | 0.2 | 0.3 ^a | 0.3 | 0.3 ^b | 0.2 | 0.3 | 0.2 | 0.3 ^a | 0.2 | 0.2 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 2.7 | 2.7 | 2.5 | 2.7 | 2.9 | 2.8 | 2.5 | 2.8 | 2.7 | 2.4 | 2.6 |
| Pain Relievers | 1.9 | 2.0 | 1.8 | 1.9 | 2.1 ^a | 2.1 | 1.9 | 2.1 | 2.0 | 1.7 | 1.9 |
| OxyContin [®] | -- | -- | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 ^a | 0.2 ^b | 0.2 | 0.1 |
| Tranquilizers | 0.8 | 0.8 | 0.7 ^a | 0.7 | 0.7 | 0.7 | 0.7 | 0.8 | 0.9 | 0.7 | 0.8 |
| Stimulants ³ | 0.6 | 0.6 | 0.5 | 0.5 | 0.6 | 0.4 | 0.4 | 0.5 | 0.4 | 0.4 | 0.5 |
| Methamphetamine ³ | 0.3 ^b | 0.3 ^b | 0.3 ^b | 0.3 ^a | 0.3 ^b | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 |
| Sedatives | 0.2 ^a | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 3.7 | 3.7 | 3.4 | 3.7 | 3.9 ^b | 3.8 | 3.4 | 3.7 | 3.6 | 3.1 ^a | 3.4 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.4A Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|
| ILLCIT DRUGS¹ | 7,642 ^b | 7,631 ^b | 7,566 ^b | 7,022 ^b | 7,014 ^b | 6,638 ^b | 6,565 ^b | 6,629 ^b | 6,278 | 6,373 ^a | 6,044 |
| Marijuana and Hashish | 5,104 ^b | 4,888 ^b | 4,788 ^b | 4,402 | 4,400 | 4,096 | 4,138 | 4,204 | 4,157 | 4,378 | 4,232 |
| Cocaine | 670 ^b | 648 ^b | 611 ^b | 582 ^b | 571 ^b | 544 ^b | 473 ^b | 405 ^b | 363 ^b | 337 ^a | 265 |
| Crack | 163 ^b | 157 ^b | 125 ^b | 108 ^b | 129 ^b | 107 ^b | 80 ^b | 58 | 60 | 68 | 34 |
| Heroin | 94 | 80 | 78 | 60 | 60 | 55 | 67 | 56 | 46 | 70 | 59 |
| Hallucinogens | 1,419 ^b | 1,250 ^b | 1,160 ^b | 998 ^b | 990 ^b | 898 | 972 ^b | 1,015 ^b | 1,000 ^b | 930 | 823 |
| LSD | 670 ^b | 397 ^b | 313 ^a | 288 | 219 | 202 | 271 | 249 | 231 | 229 | 249 |
| PCP | 233 ^b | 204 ^b | 175 ^b | 182 ^b | 128 | 105 | 102 | 99 | 76 | 76 | 96 |
| Ecstasy | 820 ^b | 603 ^b | 523 | 399 ^a | 476 | 446 | 524 | 572 | 618 ^b | 605 ^a | 488 |
| Inhalants | 2,605 ^b | 2,670 ^b | 2,762 ^b | 2,658 ^b | 2,559 ^b | 2,428 ^b | 2,326 ^b | 2,283 ^b | 2,015 ^b | 1,877 ^b | 1,613 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 3,402 ^b | 3,372 ^b | 3,403 ^b | 3,019 ^b | 3,175 ^b | 2,954 ^b | 2,783 ^b | 2,810 ^b | 2,591 | 2,522 | 2,482 |
| Pain Relievers | 2,775 ^b | 2,798 ^b | 2,879 ^b | 2,517 ^b | 2,650 ^b | 2,457 ^b | 2,353 ^b | 2,393 ^b | 2,245 ^a | 2,150 | 2,069 |
| OxyContin [®] | 224 ^b | 254 ^a | 305 | 276 | 343 | 360 | 344 | 381 | 369 | 369 | 322 |
| Tranquilizers | 834 ^a | 874 ^b | 816 ^a | 766 | 742 | 748 | 721 | 696 | 723 | 699 | 703 |
| Stimulants ³ | 1,112 ^b | 1,046 ^b | 906 ^b | 870 ^b | 930 ^b | 730 ^b | 673 ^b | 583 ^a | 531 | 508 | 495 |
| Methamphetamine ³ | 416 ^b | 382 ^b | 346 ^b | 319 ^b | 341 ^b | 221 ^b | 195 | 187 | 157 | 184 | 145 |
| Sedatives | 238 ^a | 249 ^b | 251 ^b | 232 ^a | 224 | 213 | 205 | 224 | 178 | 144 | 175 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 5,286 ^b | 5,319 ^b | 5,355 ^b | 4,944 ^b | 4,951 ^b | 4,649 ^b | 4,513 ^b | 4,501 ^b | 4,166 ^b | 4,007 ^b | 3,691 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.4B Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 30.9 ^b | 30.5 ^b | 30.0 ^b | 27.7 ^b | 27.6 ^b | 26.3 ^b | 26.4 ^b | 26.9 ^b | 25.8 ^b | 25.5 ^a | 24.2 |
| Marijuana and Hashish | 20.6 ^b | 19.6 ^b | 19.0 ^b | 17.4 | 17.3 | 16.2 | 16.6 | 17.1 | 17.1 | 17.5 | 17.0 |
| Cocaine | 2.7 ^b | 2.6 ^b | 2.4 ^b | 2.3 ^b | 2.2 ^b | 2.2 ^b | 1.9 ^b | 1.6 ^b | 1.5 ^b | 1.3 ^a | 1.1 |
| Crack | 0.7 ^b | 0.6 ^b | 0.5 ^b | 0.4 ^b | 0.5 ^b | 0.4 ^b | 0.3 ^b | 0.2 | 0.2 | 0.3 | 0.1 |
| Heroin | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 |
| Hallucinogens | 5.7 ^b | 5.0 ^b | 4.6 ^b | 3.9 ^b | 3.9 ^b | 3.6 | 3.9 ^b | 4.1 ^b | 4.1 ^b | 3.7 | 3.3 |
| LSD | 2.7 ^b | 1.6 ^b | 1.2 | 1.1 | 0.9 | 0.8 | 1.1 | 1.0 | 0.9 | 0.9 | 1.0 |
| PCP | 0.9 ^b | 0.8 ^b | 0.7 ^b | 0.7 ^b | 0.5 | 0.4 | 0.4 | 0.4 | 0.3 | 0.3 | 0.4 |
| Ecstasy | 3.3 ^b | 2.4 ^b | 2.1 | 1.6 ^a | 1.9 | 1.8 | 2.1 | 2.3 ^a | 2.5 ^b | 2.4 ^a | 2.0 |
| Inhalants | 10.5 ^b | 10.7 ^b | 11.0 ^b | 10.5 ^b | 10.1 ^b | 9.6 ^b | 9.3 ^b | 9.3 ^b | 8.3 ^b | 7.5 ^b | 6.5 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 13.7 ^b | 13.5 ^b | 13.5 ^b | 11.9 ^b | 12.5 ^b | 11.7 ^b | 11.2 ^b | 11.4 ^b | 10.6 | 10.1 | 10.0 |
| Pain Relievers | 11.2 ^b | 11.2 ^b | 11.4 ^b | 9.9 ^b | 10.4 ^b | 9.7 ^b | 9.5 ^b | 9.7 ^b | 9.2 ^b | 8.6 | 8.3 |
| OxyContin [®] | 0.9 ^b | 1.0 ^a | 1.2 | 1.1 | 1.4 | 1.4 | 1.4 | 1.5 | 1.5 | 1.5 | 1.3 |
| Tranquilizers | 3.4 ^a | 3.5 ^b | 3.2 ^a | 3.0 | 2.9 | 3.0 | 2.9 | 2.8 | 3.0 | 2.8 | 2.8 |
| Stimulants ³ | 4.5 ^b | 4.2 ^b | 3.6 ^b | 3.4 ^b | 3.7 ^b | 2.9 ^b | 2.7 ^b | 2.4 ^a | 2.2 | 2.0 | 2.0 |
| Methamphetamine ³ | 1.7 ^b | 1.5 ^b | 1.4 ^b | 1.3 ^b | 1.3 ^b | 0.9 ^b | 0.8 | 0.8 | 0.6 | 0.7 | 0.6 |
| Sedatives | 1.0 ^a | 1.0 ^b | 1.0 ^b | 0.9 | 0.9 | 0.8 | 0.8 | 0.9 | 0.7 | 0.6 | 0.7 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 21.4 ^b | 21.3 ^b | 21.2 ^b | 19.5 ^b | 19.5 ^b | 18.4 ^b | 18.1 ^b | 18.3 ^b | 17.1 ^b | 16.0 ^b | 14.8 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.5A Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|
| ILLCIT DRUGS¹ | 5,494 ^b | 5,455 ^b | 5,300 ^b | 5,041 ^b | 4,978 ^b | 4,737 ^a | 4,751 ^a | 4,838 ^b | 4,745 ^a | 4,735 ^a | 4,468 |
| Marijuana and Hashish | 3,905 ^b | 3,745 ^b | 3,656 ^b | 3,383 | 3,355 | 3,158 | 3,258 | 3,363 | 3,414 | 3,553 | 3,363 |
| Cocaine | 508 ^b | 438 ^b | 411 ^b | 423 ^b | 411 ^b | 385 ^b | 306 ^b | 242 ^a | 239 ^a | 229 | 180 |
| Crack | 100 ^b | 95 ^b | 63 ^b | 54 ^b | 74 ^b | 65 ^b | 34 | 29 | 18 | 29 | 19 |
| Heroin | 51 | 36 | 48 | 37 | 37 | 24 | 41 | 34 | 28 | 54 | 31 |
| Hallucinogens | 945 ^b | 778 ^b | 753 ^b | 665 ^a | 653 ^a | 629 | 689 ^b | 737 ^b | 747 ^b | 648 ^a | 553 |
| LSD | 314 ^b | 149 | 157 | 150 | 106 ^a | 126 | 185 | 143 | 136 | 159 | 152 |
| PCP | 110 ^b | 89 ^a | 72 | 85 ^a | 59 | 56 | 61 | 48 | 35 | 39 | 48 |
| Ecstasy | 547 ^b | 329 | 310 | 256 | 308 | 328 | 346 | 416 ^b | 465 ^b | 419 ^b | 288 |
| Inhalants | 1,078 ^b | 1,119 ^b | 1,168 ^b | 1,144 ^b | 1,109 ^b | 995 ^b | 986 ^b | 960 ^b | 885 ^b | 820 ^b | 647 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 2,285 ^b | 2,295 ^b | 2,228 ^b | 2,101 ^b | 2,184 ^b | 2,037 ^b | 1,919 ^b | 1,895 ^b | 1,812 ^a | 1,749 | 1,642 |
| Pain Relievers | 1,870 ^b | 1,917 ^b | 1,865 ^b | 1,741 ^b | 1,821 ^b | 1,684 ^b | 1,613 ^b | 1,628 ^b | 1,522 ^a | 1,484 ^a | 1,333 |
| OxyContin [®] | -- | -- | 194 | 187 | 200 | 232 | 226 | 225 | 235 | 219 | 187 |
| Tranquilizers | 568 ^a | 586 ^b | 532 | 497 | 484 | 505 | 483 | 462 | 466 | 475 | 452 |
| Stimulants ³ | 665 ^b | 578 ^b | 507 ^b | 506 ^b | 548 ^b | 415 ^a | 373 | 334 | 325 | 299 | 324 |
| Methamphetamine ³ | 245 ^b | 185 ^b | 177 ^b | 178 ^b | 185 ^b | 116 | 94 | 87 | 98 | 106 | 82 |
| Sedatives | 138 ^b | 131 ^a | 133 ^b | 118 | 110 | 111 | 113 | 126 ^a | 111 | 84 | 81 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 3,338 ^b | 3,351 ^b | 3,269 ^b | 3,129 ^b | 3,147 ^b | 2,921 ^b | 2,860 ^b | 2,849 ^b | 2,758 ^b | 2,565 ^b | 2,297 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.5B Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 22.2 ^b | 21.8 ^b | 21.0 ^b | 19.9 ^b | 19.6 ^b | 18.8 | 19.1 ^a | 19.7 ^b | 19.5 ^b | 19.0 ^a | 17.9 |
| Marijuana and Hashish | 15.8 ^b | 15.0 ^b | 14.5 ^a | 13.3 | 13.2 | 12.5 ^a | 13.1 | 13.7 | 14.0 | 14.2 | 13.5 |
| Cocaine | 2.1 ^b | 1.8 ^b | 1.6 ^b | 1.7 ^b | 1.6 ^b | 1.5 ^b | 1.2 ^b | 1.0 ^a | 1.0 ^a | 0.9 | 0.7 |
| Crack | 0.4 ^b | 0.4 ^b | 0.3 ^b | 0.2 ^b | 0.3 ^b | 0.3 ^b | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Heroin | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 |
| Hallucinogens | 3.8 ^b | 3.1 ^b | 3.0 ^b | 2.6 ^a | 2.6 | 2.5 | 2.8 ^b | 3.0 ^b | 3.1 ^b | 2.6 | 2.2 |
| LSD | 1.3 ^b | 0.6 | 0.6 | 0.6 | 0.4 ^a | 0.5 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 |
| PCP | 0.4 ^b | 0.4 ^a | 0.3 | 0.3 ^a | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 |
| Ecstasy | 2.2 ^b | 1.3 | 1.2 | 1.0 | 1.2 | 1.3 | 1.4 | 1.7 ^b | 1.9 ^b | 1.7 ^b | 1.2 |
| Inhalants | 4.4 ^b | 4.5 ^b | 4.6 ^b | 4.5 ^b | 4.4 ^b | 3.9 ^b | 4.0 ^b | 3.9 ^b | 3.6 ^b | 3.3 ^b | 2.6 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 9.2 ^b | 9.2 ^b | 8.8 ^b | 8.3 ^b | 8.6 ^b | 8.1 ^b | 7.7 ^b | 7.7 ^b | 7.4 ^b | 7.0 | 6.6 |
| Pain Relievers | 7.6 ^b | 7.7 ^b | 7.4 ^b | 6.9 ^b | 7.2 ^b | 6.7 ^b | 6.5 ^b | 6.6 ^b | 6.3 ^b | 5.9 ^a | 5.3 |
| OxyContin [®] | -- | -- | 0.8 | 0.7 | 0.8 | 0.9 | 0.9 | 0.9 | 1.0 | 0.9 | 0.7 |
| Tranquilizers | 2.3 ^a | 2.3 ^b | 2.1 | 2.0 | 1.9 | 2.0 | 1.9 | 1.9 | 1.9 | 1.9 | 1.8 |
| Stimulants ³ | 2.7 ^b | 2.3 ^b | 2.0 ^b | 2.0 ^b | 2.2 ^b | 1.6 ^a | 1.5 | 1.4 | 1.3 | 1.2 | 1.3 |
| Methamphetamine ³ | 1.0 ^b | 0.7 ^b | 0.7 ^b | 0.7 ^b | 0.7 ^b | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 |
| Sedatives | 0.6 ^b | 0.5 ^a | 0.5 ^b | 0.5 | 0.4 | 0.4 | 0.5 | 0.5 ^a | 0.5 | 0.3 | 0.3 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 13.5 ^b | 13.4 ^b | 13.0 ^b | 12.3 ^b | 12.4 ^b | 11.6 ^b | 11.5 ^b | 11.6 ^b | 11.3 ^b | 10.3 ^b | 9.2 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.6A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|
| ILLCIT DRUGS¹ | 2,878 ^b | 2,811 ^b | 2,674 ^b | 2,511 | 2,488 | 2,415 | 2,312 | 2,484 | 2,468 | 2,522 | 2,377 |
| Marijuana and Hashish | 2,023 ^b | 1,971 ^a | 1,909 | 1,730 | 1,695 | 1,693 | 1,668 | 1,813 | 1,802 | 1,967 ^a | 1,800 |
| Cocaine | 146 ^b | 153 ^b | 126 ^b | 142 ^b | 110 ^b | 104 ^b | 98 ^b | 70 ^b | 60 ^a | 67 ^b | 30 |
| Crack | 26 | 32 | 16 | 25 | 12 | 16 | 7 | 8 | 4 | 3 | * |
| Heroin | 11 | 18 | 15 | 15 | 16 | 3 | 14 | 13 | 8 | 15 | * |
| Hallucinogens | 255 ^b | 260 ^b | 211 ^a | 205 | 182 | 174 | 245 ^b | 217 ^a | 215 ^a | 215 | 160 |
| LSD | 61 ^a | 51 ^a | 38 | 32 | 31 | 30 | 52 ^a | 35 | 46 | 34 | 26 |
| PCP | 23 ^b | 20 ^a | 11 | 17 | 12 | 9 | 15 ^a | 12 | 6 | 4 | 5 |
| Ecstasy | 124 ^b | 91 | 77 | 74 | 83 | 65 | 105 | 115 ^a | 123 ^b | 108 | 73 |
| Inhalants | 301 ^b | 320 ^b | 312 ^b | 313 ^b | 340 ^b | 294 ^b | 270 ^b | 244 | 262 ^a | 223 | 188 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 986 ^b | 997 ^b | 914 ^b | 838 ^b | 846 ^b | 830 ^a | 711 | 774 | 721 | 688 | 693 |
| Pain Relievers | 794 ^b | 801 ^b | 751 ^b | 686 ^b | 692 ^b | 671 ^a | 566 | 658 ^a | 607 | 563 | 558 |
| OxyContin [®] | -- | -- | 66 | 34 | 29 | 47 | 49 | 70 | 47 | 51 | 49 |
| Tranquilizers | 188 | 226 ^b | 161 | 161 | 125 | 181 | 151 | 144 | 134 | 162 | 140 |
| Stimulants ³ | 208 ^b | 222 ^b | 180 ^b | 172 ^a | 171 ^a | 126 | 112 | 127 | 107 | 98 | 118 |
| Methamphetamine ³ | 71 ^b | 70 ^b | 58 ^a | 72 ^b | 53 | 34 | 29 | 36 | 25 | 37 | 30 |
| Sedatives | 47 | 47 | 26 | 27 | 45 | 34 | 35 | 40 | 35 | 24 | 26 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 1,421 ^b | 1,417 ^b | 1,328 ^b | 1,253 ^b | 1,252 ^b | 1,182 ^b | 1,092 ^b | 1,122 ^b | 1,092 ^b | 1,026 | 918 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.6B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------|
| ILLCIT DRUGS¹ | 11.6 ^b | 11.2 ^b | 10.6 ^b | 9.9 | 9.8 | 9.6 | 9.3 | 10.1 | 10.1 | 10.1 | 9.5 |
| Marijuana and Hashish | 8.2 ^b | 7.9 ^a | 7.6 | 6.8 | 6.7 | 6.7 | 6.7 | 7.4 | 7.4 | 7.9 ^a | 7.2 |
| Cocaine | 0.6 ^b | 0.6 ^b | 0.5 ^b | 0.6 ^b | 0.4 ^b | 0.4 ^b | 0.4 ^b | 0.3 ^b | 0.2 ^a | 0.3 ^b | 0.1 |
| Crack | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | * |
| Heroin | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | * |
| Hallucinogens | 1.0 ^b | 1.0 ^b | 0.8 | 0.8 | 0.7 | 0.7 | 1.0 ^b | 0.9 ^a | 0.9 ^a | 0.9 | 0.6 |
| LSD | 0.2 ^a | 0.2 ^a | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 ^a | 0.1 | 0.2 | 0.1 | 0.1 |
| PCP | 0.1 ^b | 0.1 ^a | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 ^a | 0.1 | 0.0 | 0.0 | 0.0 |
| Ecstasy | 0.5 ^b | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 ^a | 0.5 ^b | 0.4 | 0.3 |
| Inhalants | 1.2 ^b | 1.3 ^b | 1.2 ^b | 1.2 ^b | 1.3 ^b | 1.2 ^b | 1.1 ^b | 1.0 | 1.1 ^b | 0.9 | 0.8 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 4.0 ^b | 4.0 ^b | 3.6 ^b | 3.3 ^a | 3.3 ^a | 3.3 ^a | 2.9 | 3.1 | 3.0 | 2.8 | 2.8 |
| Pain Relievers | 3.2 ^b | 3.2 ^b | 3.0 ^b | 2.7 ^a | 2.7 ^a | 2.7 ^a | 2.3 | 2.7 ^a | 2.5 | 2.3 | 2.2 |
| OxyContin [®] | -- | -- | 0.3 | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 |
| Tranquilizers | 0.8 | 0.9 ^b | 0.6 | 0.6 | 0.5 | 0.7 | 0.6 | 0.6 | 0.5 | 0.6 | 0.6 |
| Stimulants ³ | 0.8 ^b | 0.9 ^b | 0.7 ^a | 0.7 ^a | 0.7 ^a | 0.5 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 |
| Methamphetamine ³ | 0.3 ^b | 0.3 ^b | 0.2 ^a | 0.3 ^b | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Sedatives | 0.2 ^a | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 5.7 ^b | 5.7 ^b | 5.3 ^b | 4.9 ^b | 4.9 ^b | 4.7 ^b | 4.4 ^b | 4.6 ^b | 4.5 ^b | 4.1 | 3.7 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.7A Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------------------|---------------------|--------|--------|
| ILLCIT DRUGS¹ | 18,562 ^b | 19,195 ^b | 19,043 ^b | 19,222 ^b | 19,328 ^b | 18,831 ^b | 18,772 ^b | 19,658 | 19,538 | 19,534 | 19,977 |
| Marijuana and Hashish | 16,680 ^b | 17,103 ^b | 16,983 ^b | 17,032 ^b | 17,180 ^b | 16,652 ^b | 16,718 ^b | 17,677 | 17,529 ^a | 17,796 | 18,052 |
| Cocaine | 4,786 ^b | 4,755 ^b | 4,893 ^b | 4,905 ^b | 5,139 ^b | 4,925 ^b | 4,772 ^b | 4,997 ^b | 4,568 | 4,253 | 4,267 |
| Crack | 1,182 ^b | 1,193 ^b | 1,135 ^b | 1,145 ^b | 1,176 ^b | 985 ^b | 916 ^b | 965 ^b | 886 ^b | 714 | 654 |
| Heroin | 487 ^b | 516 ^b | 529 ^a | 496 ^b | 514 ^b | 498 ^b | 467 ^b | 587 | 614 | 594 | 664 |
| Hallucinogens | 7,517 ^b | 7,377 ^b | 6,855 ^b | 6,820 ^b | 6,625 ^b | 6,291 | 5,876 | 6,418 | 6,295 | 6,029 | 6,103 |
| LSD | 4,928 ^b | 4,450 ^b | 3,900 ^b | 3,417 ^b | 2,933 ^b | 2,381 ^b | 2,167 | 2,326 ^a | 2,194 | 2,054 | 2,056 |
| PCP | 834 ^b | 948 ^b | 737 ^b | 716 ^b | 641 ^b | 540 ^b | 523 ^b | 480 ^b | 447 ^a | 351 | 345 |
| Ecstasy | 4,674 | 4,683 | 4,439 | 4,464 | 4,392 | 4,186 | 4,008 ^b | 4,194 | 4,230 | 4,222 | 4,452 |
| Inhalants | 4,867 ^b | 4,729 ^b | 4,516 ^b | 4,333 ^b | 4,097 ^b | 3,703 ^b | 3,445 ^b | 3,620 ^b | 3,396 ^b | 3,106 | 2,918 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 8,778 ^b | 9,360 | 9,564 | 9,990 | 10,067 | 9,807 | 9,694 | 9,896 | 9,832 | 9,373 | 9,709 |
| Pain Relievers | 6,854 ^b | 7,513 | 7,813 | 8,279 ^b | 8,353 ^b | 8,143 ^a | 8,096 | 8,221 ^a | 8,143 ^a | 7,600 | 7,759 |
| OxyContin [®] | 820 ^b | 1,145 ^b | 1,376 ^b | 1,616 ^b | 1,684 ^b | 1,716 ^b | 1,765 ^a | 2,110 | 2,149 | 2,079 | 2,011 |
| Tranquilizers | 3,473 ^b | 3,898 ^b | 3,919 ^b | 4,356 | 4,286 | 4,155 | 4,069 ^a | 4,323 | 4,503 | 4,232 | 4,420 |
| Stimulants ³ | 3,796 ^b | 3,873 ^b | 3,892 ^b | 4,008 ^b | 3,926 ^b | 3,569 | 3,485 | 3,587 | 3,534 | 3,172 | 3,374 |
| Methamphetamine ³ | 2,301 ^b | 2,192 ^b | 2,247 ^b | 2,190 ^b | 2,103 ^b | 1,847 ^b | 1,548 ^b | 1,518 ^b | 1,361 ^b | 1,150 | 1,089 |
| Sedatives | 648 ^b | 565 ^b | 627 ^b | 650 ^b | 612 ^b | 583 ^b | 511 | 528 ^a | 469 | 472 | 420 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 12,436 | 12,770 ^a | 12,630 | 12,800 ^b | 12,959 ^b | 12,573 | 12,146 | 12,592 | 12,410 | 11,860 | 12,216 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.7B Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|------|
| ILLCIT DRUGS¹ | 59.8 ^b | 60.5 ^b | 59.2 | 59.2 ^a | 59.0 | 57.5 | 57.0 | 58.5 | 57.3 | 56.9 | 57.8 |
| Marijuana and Hashish | 53.8 ^a | 53.9 ^a | 52.8 | 52.4 | 52.5 | 50.9 | 50.8 ^a | 52.6 | 51.4 | 51.9 | 52.2 |
| Cocaine | 15.4 ^b | 15.0 ^b | 15.2 ^b | 15.1 ^b | 15.7 ^b | 15.0 ^b | 14.5 ^b | 14.9 ^b | 13.4 ^a | 12.4 | 12.3 |
| Crack | 3.8 ^b | 3.8 ^b | 3.5 ^b | 3.5 ^b | 3.6 ^b | 3.0 ^b | 2.8 ^b | 2.9 ^b | 2.6 ^b | 2.1 | 1.9 |
| Heroin | 1.6 ^a | 1.6 | 1.6 | 1.5 ^a | 1.6 ^a | 1.5 ^a | 1.4 ^b | 1.7 | 1.8 | 1.7 | 1.9 |
| Hallucinogens | 24.2 ^b | 23.3 ^b | 21.3 ^b | 21.0 ^b | 20.2 ^b | 19.2 ^b | 17.8 | 19.1 ^b | 18.5 | 17.6 | 17.6 |
| LSD | 15.9 ^b | 14.0 ^b | 12.1 ^b | 10.5 ^b | 9.0 ^b | 7.3 ^b | 6.6 | 6.9 ^b | 6.4 | 6.0 | 5.9 |
| PCP | 2.7 ^b | 3.0 ^b | 2.3 ^b | 2.2 ^b | 2.0 ^b | 1.6 ^b | 1.6 ^b | 1.4 ^b | 1.3 ^a | 1.0 | 1.0 |
| Ecstasy | 15.1 ^b | 14.8 ^b | 13.8 | 13.7 | 13.4 | 12.8 | 12.2 | 12.5 | 12.4 | 12.3 | 12.9 |
| Inhalants | 15.7 ^b | 14.9 ^b | 14.0 ^b | 13.3 ^b | 12.5 ^b | 11.3 ^b | 10.5 ^b | 10.8 ^b | 10.0 ^b | 9.1 | 8.4 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 28.3 | 29.5 ^a | 29.7 ^b | 30.8 ^b | 30.7 ^b | 30.0 ^b | 29.4 ^a | 29.5 ^a | 28.9 | 27.3 | 28.1 |
| Pain Relievers | 22.1 | 23.7 ^a | 24.3 ^b | 25.5 ^b | 25.5 ^b | 24.9 ^b | 24.6 ^b | 24.5 ^b | 23.9 ^b | 22.2 | 22.4 |
| OxyContin [®] | 2.6 ^b | 3.6 ^b | 4.3 ^b | 5.0 ^b | 5.1 ^a | 5.2 | 5.4 | 6.3 | 6.3 | 6.1 | 5.8 |
| Tranquilizers | 11.2 ^b | 12.3 | 12.2 | 13.4 | 13.1 | 12.7 | 12.4 | 12.9 | 13.2 | 12.3 | 12.8 |
| Stimulants ³ | 12.2 ^b | 12.2 ^b | 12.1 ^b | 12.3 ^b | 12.0 ^b | 10.9 ^b | 10.6 ^a | 10.7 ^a | 10.4 | 9.2 | 9.8 |
| Methamphetamine ³ | 7.4 ^b | 6.9 ^b | 7.0 ^b | 6.7 ^b | 6.4 ^b | 5.6 ^b | 4.7 ^b | 4.5 ^b | 4.0 ^b | 3.4 | 3.1 |
| Sedatives | 2.1 ^b | 1.8 ^b | 1.9 ^b | 2.0 ^b | 1.9 ^b | 1.8 ^b | 1.6 ^a | 1.6 ^a | 1.4 | 1.4 | 1.2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 40.1 ^b | 40.2 ^b | 39.2 ^b | 39.4 ^b | 39.6 ^b | 38.4 ^b | 36.9 ^a | 37.5 ^b | 36.4 | 34.6 | 35.3 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.8A Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 11,000 ^b | 10,976 ^b | 10,913 ^b | 11,117 ^b | 11,267 ^b | 10,889 ^b | 11,090 ^b | 12,186 | 11,990 ^a | 12,067 ^a | 12,551 |
| Marijuana and Hashish | 9,238 ^b | 9,044 ^b | 8,963 ^b | 9,111 ^b | 9,188 ^b | 9,008 ^b | 9,154 ^b | 10,343 ^a | 10,215 ^b | 10,554 | 10,887 |
| Cocaine | 2,087 ^b | 2,098 ^b | 2,134 ^b | 2,247 ^b | 2,263 ^b | 2,087 ^b | 1,844 ^a | 1,790 | 1,593 | 1,563 | 1,600 |
| Crack | 266 ^b | 275 ^b | 263 ^b | 328 ^b | 293 ^b | 249 ^b | 217 ^b | 172 | 180 | 107 | 140 |
| Heroin | 122 ^b | 96 ^b | 118 ^b | 159 ^b | 147 ^b | 142 ^b | 149 ^b | 183 ^b | 211 | 230 | 272 |
| Hallucinogens | 2,610 ^b | 2,112 | 1,944 ^a | 2,051 | 2,151 | 2,042 | 2,101 | 2,430 | 2,439 | 2,323 | 2,236 |
| LSD | 553 | 336 ^b | 317 ^b | 322 ^b | 380 ^b | 375 ^b | 507 | 523 | 549 | 586 | 622 |
| PCP | 95 ^a | 117 ^b | 100 ^a | 57 | 76 | 57 | 30 | 35 | 37 | 63 | 57 |
| Ecstasy | 1,784 ^b | 1,164 ^b | 999 ^b | 998 ^b | 1,235 ^a | 1,146 ^b | 1,270 | 1,456 | 1,500 | 1,413 | 1,412 |
| Inhalants | 685 ^b | 662 ^b | 674 ^b | 685 ^b | 578 ^a | 528 | 538 | 635 ^b | 597 ^a | 500 | 473 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 4,452 | 4,637 | 4,807 | 4,915 | 5,130 ^a | 4,920 | 4,891 | 5,061 ^a | 4,876 | 4,366 ^a | 4,729 |
| Pain Relievers | 3,540 | 3,803 ^a | 3,834 ^a | 4,030 ^b | 4,077 ^b | 3,981 ^b | 3,956 ^b | 4,027 ^b | 3,779 ^a | 3,353 | 3,507 |
| OxyContin [®] | -- | -- | 562 | 559 | 577 | 568 | 602 | 773 ^b | 752 ^b | 628 | 568 |
| Tranquilizers | 1,509 ^a | 1,682 | 1,658 | 1,796 | 1,793 | 1,702 | 1,762 | 1,802 | 1,803 | 1,555 | 1,718 |
| Stimulants ³ | 1,233 | 1,197 | 1,280 | 1,242 | 1,376 | 1,132 ^a | 1,123 ^b | 1,238 | 1,242 | 1,065 ^b | 1,361 |
| Methamphetamine ³ | 619 ^b | 594 ^b | 619 ^b | 576 ^b | 553 ^b | 406 | 270 | 303 | 268 | 225 ^a | 341 |
| Sedatives | 162 | 156 | 163 | 191 ^a | 185 ^a | 171 | 178 ^a | 160 | 166 | 128 | 119 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 6,277 | 6,266 | 6,201 | 6,467 | 6,638 ^a | 6,374 | 6,253 | 6,701 ^b | 6,395 | 5,972 | 6,217 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.8B Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 35.5 | 34.6 ^a | 33.9 ^b | 34.2 ^b | 34.4 ^b | 33.3 ^b | 33.7 ^b | 36.3 | 35.2 | 35.2 | 36.3 |
| Marijuana and Hashish | 29.8 ^a | 28.5 ^b | 27.8 ^b | 28.0 ^b | 28.1 ^b | 27.5 ^b | 27.8 ^b | 30.8 | 30.0 ^a | 30.8 | 31.5 |
| Cocaine | 6.7 ^b | 6.6 ^b | 6.6 ^b | 6.9 ^b | 6.9 ^b | 6.4 ^b | 5.6 ^b | 5.3 ^a | 4.7 | 4.6 | 4.6 |
| Crack | 0.9 ^b | 0.9 ^b | 0.8 ^b | 1.0 ^b | 0.9 ^b | 0.8 ^b | 0.7 ^b | 0.5 | 0.5 | 0.3 | 0.4 |
| Heroin | 0.4 ^b | 0.3 ^b | 0.4 ^b | 0.5 ^b | 0.4 ^b | 0.4 ^b | 0.5 ^b | 0.5 ^a | 0.6 | 0.7 | 0.8 |
| Hallucinogens | 8.4 ^b | 6.7 | 6.0 | 6.3 | 6.6 | 6.2 | 6.4 | 7.2 ^a | 7.2 | 6.8 | 6.5 |
| LSD | 1.8 | 1.1 ^b | 1.0 ^b | 1.0 ^b | 1.2 ^b | 1.1 ^b | 1.5 | 1.6 | 1.6 | 1.7 | 1.8 |
| PCP | 0.3 ^a | 0.4 ^b | 0.3 ^a | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 |
| Ecstasy | 5.8 ^b | 3.7 | 3.1 ^b | 3.1 ^b | 3.8 | 3.5 ^a | 3.9 | 4.3 | 4.4 | 4.1 | 4.1 |
| Inhalants | 2.2 ^b | 2.1 ^b | 2.1 ^b | 2.1 ^b | 1.8 ^a | 1.6 | 1.6 | 1.9 ^b | 1.8 ^a | 1.5 | 1.4 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 14.4 | 14.6 ^a | 14.9 ^b | 15.1 ^b | 15.7 ^b | 15.0 ^b | 14.9 ^a | 15.1 ^b | 14.3 | 12.7 ^a | 13.7 |
| Pain Relievers | 11.4 ^b | 12.0 ^b | 11.9 ^b | 12.4 ^b | 12.5 ^b | 12.2 ^b | 12.0 ^b | 12.0 ^b | 11.1 ^a | 9.8 | 10.1 |
| OxyContin [®] | -- | -- | 1.7 | 1.7 | 1.8 | 1.7 | 1.8 | 2.3 ^b | 2.2 ^b | 1.8 | 1.6 |
| Tranquilizers | 4.9 | 5.3 | 5.2 | 5.5 | 5.5 | 5.2 | 5.4 | 5.4 | 5.3 | 4.5 | 5.0 |
| Stimulants ³ | 4.0 | 3.8 | 4.0 | 3.8 | 4.2 | 3.5 | 3.4 | 3.7 | 3.6 | 3.1 ^b | 3.9 |
| Methamphetamine ³ | 2.0 ^b | 1.9 ^b | 1.9 ^b | 1.8 ^b | 1.7 ^b | 1.2 | 0.8 | 0.9 | 0.8 | 0.7 ^a | 1.0 |
| Sedatives | 0.5 ^a | 0.5 | 0.5 ^a | 0.6 ^b | 0.6 ^b | 0.5 ^a | 0.5 ^a | 0.5 | 0.5 | 0.4 | 0.3 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 20.2 ^b | 19.7 ^b | 19.3 ^a | 19.9 ^b | 20.3 ^b | 19.5 ^b | 19.0 | 20.0 ^b | 18.8 | 17.4 | 18.0 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.9A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|-------|
| ILLCIT DRUGS¹ | 6,264 ^b | 6,435 ^b | 6,239 ^b | 6,543 ^b | 6,498 ^b | 6,475 ^b | 6,490 ^b | 7,176 | 7,345 | 7,352 | 7,381 |
| Marijuana and Hashish | 5,376 ^b | 5,400 ^b | 5,196 ^b | 5,388 ^b | 5,337 ^b | 5,392 ^b | 5,459 ^b | 6,128 | 6,313 | 6,526 | 6,485 |
| Cocaine | 631 ^b | 696 ^b | 692 ^b | 832 ^b | 730 ^b | 572 ^b | 515 ^a | 466 | 503 ^a | 464 | 398 |
| Crack | 52 | 55 | 80 ^b | 104 ^b | 72 ^a | 56 | 54 | 40 | 72 | 20 | 38 |
| Heroin | 39 ^b | 25 ^b | 43 ^b | 59 ^b | 55 ^b | 49 ^b | 76 ^a | 67 ^b | 90 | 107 | 138 |
| Hallucinogens | 589 | 545 | 477 ^a | 503 | 564 | 475 ^a | 545 | 620 | 672 | 557 | 600 |
| LSD | 36 ^b | 67 ^a | 83 | 60 ^b | 58 ^b | 69 ^a | 98 | 97 | 105 | 92 | 118 |
| PCP | 15 | 30 | 17 | 11 | 14 | 15 | 4 | 5 | 6 | 12 | 12 |
| Ecstasy | 355 | 230 ^b | 231 ^b | 267 ^a | 326 | 230 ^b | 295 | 378 | 402 | 315 | 361 |
| Inhalants | 150 | 134 | 144 | 146 | 130 | 131 | 104 | 148 | 135 | 144 | 123 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 1,698 | 1,922 | 1,978 | 2,046 ^a | 2,122 ^b | 1,947 | 1,936 | 2,133 ^b | 1,997 | 1,711 | 1,832 |
| Pain Relievers | 1,262 | 1,486 | 1,504 ^a | 1,535 ^a | 1,627 ^b | 1,518 ^a | 1,489 ^a | 1,610 ^b | 1,510 ^a | 1,220 | 1,311 |
| OxyContin [®] | -- | -- | 142 | 146 | 140 | 157 | 130 | 179 | 239 ^b | 158 | 145 |
| Tranquilizers | 487 | 525 | 566 | 606 | 651 | 573 | 570 | 622 | 570 | 546 | 537 |
| Stimulants ³ | 409 | 412 | 479 | 468 | 457 | 350 | 351 | 424 | 414 | 344 | 400 |
| Methamphetamine ³ | 183 ^b | 197 ^b | 220 ^b | 225 ^b | 183 ^b | 116 | 74 | 84 | 83 | 79 | 97 |
| Sedatives | 49 | 58 | 49 | 74 | 54 | 71 | 67 | 84 | 59 | 42 | 52 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 2,459 | 2,665 | 2,619 | 2,850 ^b | 2,912 ^b | 2,642 | 2,579 | 2,811 ^b | 2,709 ^a | 2,385 | 2,435 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.9B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|------------------|------|------|
| ILLCIT DRUGS¹ | 20.2 ^a | 20.3 | 19.4 ^b | 20.1 ^a | 19.8 ^b | 19.8 ^b | 19.7 ^b | 21.4 | 21.6 | 21.4 | 21.3 |
| Marijuana and Hashish | 17.3 ^b | 17.0 ^b | 16.1 ^b | 16.6 ^b | 16.3 ^b | 16.5 ^b | 16.6 ^b | 18.2 | 18.5 | 19.0 | 18.7 |
| Cocaine | 2.0 ^b | 2.2 ^b | 2.1 ^b | 2.6 ^b | 2.2 ^b | 1.7 ^b | 1.6 ^b | 1.4 | 1.5 ^a | 1.4 | 1.1 |
| Crack | 0.2 | 0.2 | 0.3 ^b | 0.3 ^b | 0.2 ^a | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 |
| Heroin | 0.1 ^b | 0.1 ^b | 0.1 ^b | 0.2 ^b | 0.2 ^b | 0.1 ^b | 0.2 ^a | 0.2 ^b | 0.3 | 0.3 | 0.4 |
| Hallucinogens | 1.9 | 1.7 | 1.5 | 1.5 | 1.7 | 1.5 | 1.7 | 1.8 | 2.0 | 1.6 | 1.7 |
| LSD | 0.1 ^b | 0.2 ^a | 0.3 | 0.2 ^a | 0.2 ^b | 0.2 ^a | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| PCP | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ecstasy | 1.1 | 0.7 ^a | 0.7 ^a | 0.8 | 1.0 | 0.7 ^b | 0.9 | 1.1 | 1.2 | 0.9 | 1.0 |
| Inhalants | 0.5 | 0.4 | 0.4 | 0.5 | 0.4 | 0.4 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 5.5 | 6.1 ^a | 6.1 ^b | 6.3 ^b | 6.5 ^b | 5.9 ^a | 5.9 | 6.4 ^b | 5.9 | 5.0 | 5.3 |
| Pain Relievers | 4.1 | 4.7 ^b | 4.7 ^b | 4.7 ^b | 5.0 ^b | 4.6 ^b | 4.5 ^b | 4.8 ^b | 4.4 ^a | 3.6 | 3.8 |
| OxyContin [®] | -- | -- | 0.4 | 0.4 | 0.4 | 0.5 | 0.4 | 0.5 | 0.7 ^b | 0.5 | 0.4 |
| Tranquilizers | 1.6 | 1.7 | 1.8 | 1.9 | 2.0 ^a | 1.7 | 1.7 | 1.9 | 1.7 | 1.6 | 1.6 |
| Stimulants ³ | 1.3 | 1.3 | 1.5 ^a | 1.4 ^a | 1.4 | 1.1 | 1.1 | 1.3 | 1.2 | 1.0 | 1.2 |
| Methamphetamine ³ | 0.6 ^b | 0.6 ^b | 0.7 ^b | 0.7 ^b | 0.6 ^b | 0.4 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 |
| Sedatives | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 7.9 ^b | 8.4 ^b | 8.1 ^b | 8.8 ^b | 8.9 ^b | 8.1 ^b | 7.8 ^a | 8.4 ^b | 8.0 ^b | 7.0 | 7.0 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.10A Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 82,051 ^b | 83,380 ^b | 83,448 ^b | 85,841 ^b | 85,405 ^b | 89,040 ^b | 92,478 ^b | 92,973 ^b | 94,117 ^b | 95,171 ^b | 98,787 |
| Marijuana and Hashish | 73,162 ^b | 74,620 ^b | 75,001 ^b | 76,111 ^b | 76,213 ^b | 79,988 ^b | 82,001 ^b | 83,070 ^b | 84,927 ^b | 85,668 ^a | 88,955 |
| Cocaine | 28,453 ^b | 29,488 ^b | 28,649 ^b | 28,185 ^b | 29,593 ^b | 30,477 ^b | 31,671 | 31,339 | 32,430 | 32,332 | 33,156 |
| Crack | 7,057 ^b | 6,599 ^b | 6,580 ^b | 6,675 ^b | 7,254 ^a | 7,524 | 7,458 | 7,368 ^a | 8,262 | 7,433 ^a | 8,327 |
| Heroin | 3,087 ^a | 3,148 ^a | 2,538 ^b | 2,979 ^b | 3,214 | 3,253 | 3,262 | 3,036 ^a | 3,483 | 3,499 | 3,841 |
| Hallucinogens | 25,378 ^b | 25,736 ^b | 26,318 ^b | 25,910 ^b | 27,675 ^b | 27,088 ^b | 29,217 | 29,970 | 30,248 | 29,403 | 30,982 |
| LSD | 18,917 ^b | 19,577 ^b | 19,185 ^b | 18,728 ^b | 20,213 | 20,137 | 21,166 | 21,138 | 20,950 | 20,716 | 21,430 |
| PCP | 6,352 | 5,955 | 5,851 | 5,705 | 5,865 | 5,533 | 6,032 | 5,692 | 5,731 | 5,676 | 6,112 |
| Ecstasy | 4,656 ^b | 5,619 ^b | 6,168 ^b | 6,632 ^b | 7,414 ^b | 7,786 ^b | 8,431 ^b | 9,513 ^b | 11,082 | 9,743 ^b | 11,222 |
| Inhalants | 15,398 | 15,596 | 15,521 | 15,754 | 16,200 | 16,426 | 16,659 | 16,649 | 16,367 | 15,540 | 16,563 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 35,779 ^b | 36,269 ^b | 36,190 ^b | 36,562 ^b | 37,741 ^b | 37,781 ^b | 39,677 ^a | 39,285 ^b | 39,409 ^b | 39,348 ^b | 42,198 |
| Pain Relievers | 19,983 ^b | 20,896 ^b | 21,076 ^b | 21,896 ^b | 22,475 ^b | 22,533 ^b | 24,543 ^b | 24,583 ^b | 24,521 ^b | 24,497 ^b | 27,218 |
| OxyContin [®] | 881 ^b | 1,433 ^b | 1,392 ^b | 1,589 ^b | 2,089 ^b | 2,303 ^b | 2,748 ^b | 3,373 ^b | 3,642 | 3,469 ^a | 4,247 |
| Tranquilizers | 14,959 ^b | 15,447 ^b | 15,117 ^b | 15,919 ^b | 16,290 ^b | 15,350 ^b | 16,762 ^a | 16,830 ^a | 16,960 ^a | 16,725 ^a | 18,515 |
| Stimulants ³ | 18,588 | 18,085 | 17,499 | 16,105 ^a | 17,639 | 17,369 | 17,126 | 17,863 | 17,674 | 16,699 | 17,629 |
| Methamphetamine ³ | 12,648 ^b | 12,565 ^b | 11,920 | 10,154 | 11,782 | 11,012 | 10,891 | 11,203 | 11,542 | 10,594 | 11,025 |
| Sedatives | 9,075 ^b | 8,696 ^b | 9,012 ^b | 8,101 | 8,007 | 7,604 | 8,204 | 7,882 | 7,006 | 6,899 | 7,344 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 52,578 ^b | 53,040 ^b | 52,672 ^b | 54,078 ^b | 55,024 ^b | 56,431 ^b | 59,248 ^a | 58,987 ^b | 59,895 | 59,580 ^a | 62,127 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.10B Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 45.7 ^b | 46.1 ^b | 45.6 ^b | 46.3 ^b | 45.5 ^b | 46.9 ^b | 48.2 | 48.0 | 48.2 | 48.0 | 49.3 |
| Marijuana and Hashish | 40.8 ^b | 41.2 ^b | 41.0 ^b | 41.1 ^b | 40.6 ^b | 42.1 ^b | 42.7 ^a | 42.9 ^a | 43.5 | 43.2 | 44.4 |
| Cocaine | 15.9 | 16.3 | 15.6 | 15.2 ^b | 15.7 | 16.1 | 16.5 | 16.2 | 16.6 | 16.3 | 16.5 |
| Crack | 3.9 | 3.6 ^a | 3.6 ^a | 3.6 ^a | 3.9 | 4.0 | 3.9 | 3.8 | 4.2 | 3.7 | 4.2 |
| Heroin | 1.7 | 1.7 | 1.4 ^b | 1.6 | 1.7 | 1.7 | 1.7 | 1.6 ^a | 1.8 | 1.8 | 1.9 |
| Hallucinogens | 14.1 ^b | 14.2 ^b | 14.4 ^a | 14.0 ^b | 14.7 | 14.3 ^a | 15.2 | 15.5 | 15.5 | 14.8 | 15.4 |
| LSD | 10.5 | 10.8 | 10.5 | 10.1 | 10.8 | 10.6 | 11.0 | 10.9 | 10.7 | 10.4 | 10.7 |
| PCP | 3.5 ^a | 3.3 | 3.2 | 3.1 | 3.1 | 2.9 | 3.1 | 2.9 | 2.9 | 2.9 | 3.0 |
| Ecstasy | 2.6 ^b | 3.1 ^b | 3.4 ^b | 3.6 ^b | 3.9 ^b | 4.1 ^b | 4.4 ^b | 4.9 ^b | 5.7 | 4.9 ^b | 5.6 |
| Inhalants | 8.6 | 8.6 | 8.5 | 8.5 | 8.6 | 8.7 | 8.7 | 8.6 | 8.4 | 7.8 | 8.3 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 19.9 ^a | 20.0 ^a | 19.8 ^a | 19.7 ^a | 20.1 | 19.9 ^a | 20.7 | 20.3 | 20.2 | 19.8 ^a | 21.0 |
| Pain Relievers | 11.1 ^b | 11.5 ^b | 11.5 ^b | 11.8 ^b | 12.0 ^b | 11.9 ^b | 12.8 | 12.7 ^a | 12.6 ^a | 12.4 ^b | 13.6 |
| OxyContin [®] | 0.5 ^b | 0.8 ^b | 0.8 ^b | 0.9 ^b | 1.1 ^b | 1.2 ^b | 1.4 ^b | 1.7 ^a | 1.9 | 1.7 ^a | 2.1 |
| Tranquilizers | 8.3 ^a | 8.5 ^a | 8.3 ^b | 8.6 | 8.7 | 8.1 ^b | 8.7 | 8.7 | 8.7 | 8.4 ^a | 9.2 |
| Stimulants ³ | 10.4 ^b | 10.0 ^b | 9.6 ^a | 8.7 | 9.4 | 9.1 | 8.9 | 9.2 | 9.1 | 8.4 | 8.8 |
| Methamphetamine ³ | 7.1 ^b | 6.9 ^b | 6.5 ^b | 5.5 | 6.3 ^b | 5.8 | 5.7 | 5.8 | 5.9 | 5.3 | 5.5 |
| Sedatives | 5.1 ^b | 4.8 ^b | 4.9 ^b | 4.4 ^b | 4.3 ^a | 4.0 | 4.3 ^a | 4.1 | 3.6 | 3.5 | 3.7 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 29.3 ^b | 29.3 ^b | 28.8 ^b | 29.2 ^b | 29.3 ^b | 29.7 ^a | 30.9 | 30.5 | 30.7 | 30.0 | 31.0 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.11A Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 18,639 ^b | 18,562 ^b | 18,594 ^b | 18,883 ^b | 19,565 ^b | 20,163 ^b | 19,801 ^b | 21,099 ^b | 22,174 ^b | 21,485 ^b | 24,460 |
| Marijuana and Hashish | 12,612 ^b | 12,441 ^b | 12,833 ^b | 12,880 ^b | 12,889 ^b | 12,970 ^b | 13,479 ^b | 14,983 ^b | 15,672 ^a | 15,632 ^a | 17,263 |
| Cocaine | 3,307 | 3,372 | 3,114 | 2,852 | 3,400 | 3,275 | 3,133 | 2,773 | 2,700 | 2,065 ^b | 2,890 |
| Crack | 1,188 ^b | 1,036 | 978 | 1,000 | 1,111 ^a | 1,138 ^a | 862 | 807 | 686 | 490 ^a | 762 |
| Heroin | 231 | 182 ^a | 232 | 184 ^a | 376 | 207 | 265 | 364 | 382 | 336 | 366 |
| Hallucinogens | 1,194 | 1,046 ^b | 1,182 | 1,093 ^a | 1,166 ^a | 1,089 ^a | 881 ^b | 1,369 | 1,356 | 1,098 ^a | 1,516 |
| LSD | 131 ^a | 73 ^b | 117 ^a | 91 ^b | 181 | 120 ^a | 115 ^a | 124 ^a | 197 | 135 ^a | 283 |
| PCP | 31 | 13 | 38 | 22 | 52 | 26 | * | 42 | 24 | 17 | 67 |
| Ecstasy | 836 | 626 ^a | 606 ^a | 705 | 595 ^a | 647 | 499 ^b | 932 | 687 | 590 ^a | 910 |
| Inhalants | 320 ^a | 295 ^a | 413 | 358 | 532 | 562 | 540 | 514 | 566 | 541 | 573 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 8,057 ^b | 8,231 ^b | 7,814 ^b | 8,331 ^b | 9,168 ^a | 9,377 | 8,380 ^b | 9,108 ^a | 9,362 | 8,542 ^b | 10,296 |
| Pain Relievers | 5,582 ^b | 5,950 ^b | 5,556 ^b | 6,044 ^b | 6,752 ^a | 6,851 | 6,347 ^b | 6,795 | 6,941 | 6,306 ^b | 7,649 |
| OxyContin [®] | -- | -- | 457 ^a | 480 ^a | 547 | 626 | 634 | 687 | 893 | 776 | 723 |
| Tranquilizers | 2,772 ^b | 2,783 ^b | 2,878 ^b | 2,956 ^b | 2,788 ^b | 3,086 ^b | 2,853 ^b | 3,228 ^a | 3,314 | 3,080 ^b | 3,903 |
| Stimulants ³ | 1,482 | 1,256 ^a | 1,467 | 1,340 | 1,864 | 1,447 | 1,152 ^b | 1,521 | 1,335 | 1,336 | 1,632 |
| Methamphetamine ³ | 891 | 823 | 1,012 | 849 | 1,146 ^a | 823 | 490 | 777 | 593 | 703 | 732 |
| Sedatives | 681 ^a | 544 | 441 | 441 | 632 ^a | 582 | 335 | 530 | 630 | 314 | 390 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 10,808 ^b | 10,689 ^b | 10,189 ^b | 10,513 ^b | 11,477 ^a | 11,910 | 10,906 ^b | 11,526 ^a | 11,472 ^a | 10,422 ^b | 12,753 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.11B Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 10.4 ^b | 10.3 ^b | 10.2 ^b | 10.2 ^b | 10.4 ^b | 10.6 ^b | 10.3 ^b | 10.9 ^b | 11.4 ^a | 10.8 ^b | 12.2 |
| Marijuana and Hashish | 7.0 ^b | 6.9 ^b | 7.0 ^b | 6.9 ^b | 6.9 ^b | 6.8 ^b | 7.0 ^b | 7.7 ^a | 8.0 | 7.9 ^a | 8.6 |
| Cocaine | 1.8 ^b | 1.9 ^b | 1.7 | 1.5 | 1.8 ^a | 1.7 | 1.6 | 1.4 | 1.4 | 1.0 ^b | 1.4 |
| Crack | 0.7 ^b | 0.6 ^a | 0.5 | 0.5 ^a | 0.6 ^a | 0.6 ^a | 0.4 | 0.4 | 0.4 | 0.2 ^a | 0.4 |
| Heroin | 0.1 | 0.1 | 0.1 | 0.1 ^a | 0.2 | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Hallucinogens | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 ^a | 0.5 ^b | 0.7 | 0.7 | 0.6 ^a | 0.8 |
| LSD | 0.1 | 0.0 ^b | 0.1 ^a | 0.0 ^a | 0.1 | 0.1 ^a | 0.1 ^a | 0.1 ^a | 0.1 | 0.1 ^a | 0.1 |
| PCP | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | * | 0.0 | 0.0 | 0.0 | 0.0 |
| Ecstasy | 0.5 | 0.3 | 0.3 | 0.4 | 0.3 ^a | 0.3 | 0.3 ^b | 0.5 | 0.4 | 0.3 ^a | 0.5 |
| Inhalants | 0.2 | 0.2 ^a | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 4.5 ^a | 4.5 ^a | 4.3 ^b | 4.5 ^a | 4.9 | 4.9 | 4.4 ^b | 4.7 | 4.8 | 4.3 ^b | 5.1 |
| Pain Relievers | 3.1 ^b | 3.3 ^a | 3.0 ^b | 3.3 ^a | 3.6 | 3.6 | 3.3 ^a | 3.5 | 3.6 | 3.2 ^b | 3.8 |
| OxyContin [®] | -- | -- | 0.2 ^a | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.4 | 0.4 |
| Tranquilizers | 1.5 ^a | 1.5 ^b | 1.6 ^a | 1.6 ^a | 1.5 ^b | 1.6 ^a | 1.5 ^b | 1.7 | 1.7 | 1.6 ^b | 1.9 |
| Stimulants ³ | 0.8 | 0.7 | 0.8 | 0.7 | 1.0 | 0.8 | 0.6 ^a | 0.8 | 0.7 | 0.7 | 0.8 |
| Methamphetamine ³ | 0.5 | 0.5 | 0.6 ^a | 0.5 | 0.6 ^b | 0.4 | 0.3 | 0.4 | 0.3 | 0.4 | 0.4 |
| Sedatives | 0.4 ^b | 0.3 | 0.2 | 0.2 | 0.3 ^a | 0.3 ^a | 0.2 | 0.3 | 0.3 ^a | 0.2 | 0.2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 6.0 | 5.9 | 5.6 ^b | 5.7 ^a | 6.1 | 6.3 | 5.7 ^a | 6.0 | 5.9 | 5.3 ^b | 6.4 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.12A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 10,380 ^b | 10,224 ^b | 10,159 ^b | 10,666 ^b | 11,400 ^b | 11,001 ^b | 11,321 ^b | 12,270 ^b | 12,835 ^a | 12,580 ^a | 14,105 |
| Marijuana and Hashish | 7,185 ^b | 7,267 ^b | 7,471 ^b | 7,508 ^b | 7,818 ^b | 7,385 ^b | 8,142 ^b | 8,885 ^b | 9,294 ^a | 9,578 | 10,571 |
| Cocaine | 1,243 | 1,431 | 1,204 | 1,423 | 1,586 | 1,400 | 1,252 | 1,106 | 909 | 838 ^a | 1,222 |
| Crack | 488 | 517 | 370 | 553 | 619 | 538 | 302 | 448 | 303 | 205 | 405 |
| Heroin | 116 | 76 ^a | 108 | 62 ^b | 268 | 109 | 124 | 112 | 141 | 159 | 197 |
| Hallucinogens | 352 | 238 | 241 | 380 | 262 | 349 | 269 | 427 | 320 | 200 ^a | 374 |
| LSD | 15 | 15 | 20 | 12 | 41 | 45 | * | 27 | * | 26 | 39 |
| PCP | 20 | * | 21 | 20 | * | 19 | * | 36 | 24 | * | 16 |
| Ecstasy | 197 | 149 | 142 | 161 | 120 | 209 | 146 | 269 | 165 | 120 | 193 |
| Inhalants | 183 | 117 | 181 | 152 | 292 | 191 | 270 | 170 | 293 | 257 | 182 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 3,602 ^a | 3,532 ^a | 3,218 ^b | 3,607 ^a | 4,130 | 4,130 | 3,563 ^a | 4,072 | 4,239 | 3,720 | 4,306 |
| Pain Relievers | 2,321 ^a | 2,406 ^a | 2,149 ^b | 2,437 ^a | 2,908 | 2,998 | 2,676 | 3,014 | 2,975 | 2,688 | 2,993 |
| OxyContin [®] | -- | -- | 118 | 153 | 106 | 168 | 256 | 262 | 280 | 225 | 165 |
| Tranquilizers | 1,129 | 1,078 ^a | 889 ^b | 1,050 ^a | 984 ^b | 1,085 | 1,075 ^a | 1,265 | 1,456 | 1,132 | 1,462 |
| Stimulants ³ | 685 | 676 | 653 | 548 | 757 | 574 | 445 | 748 | 561 | 528 | 654 |
| Methamphetamine ³ | 428 | 459 | 428 | 331 | 495 | 380 | 211 | 383 | 245 | 323 | 312 |
| Sedatives | 340 | 189 | 190 | 171 | 285 | 240 | 136 | 243 | 280 | 164 | 192 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 4,897 | 4,767 | 4,301 ^b | 4,860 | 5,456 | 5,471 | 4,884 | 5,262 | 5,214 | 4,610 ^a | 5,530 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.12B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------|------------------|------|
| ILLCIT DRUGS¹ | 5.8 ^b | 5.6 ^b | 5.5 ^b | 5.8 ^b | 6.1 ^b | 5.8 ^b | 5.9 ^b | 6.3 ^a | 6.6 | 6.3 ^a | 7.0 |
| Marijuana and Hashish | 4.0 ^b | 4.0 ^b | 4.1 ^b | 4.1 ^b | 4.2 ^b | 3.9 ^b | 4.2 ^b | 4.6 ^b | 4.8 | 4.8 | 5.3 |
| Cocaine | 0.7 | 0.8 | 0.7 | 0.8 | 0.8 ^a | 0.7 | 0.7 | 0.6 | 0.5 | 0.4 ^a | 0.6 |
| Crack | 0.3 | 0.3 | 0.2 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 |
| Heroin | 0.1 | 0.0 ^a | 0.1 | 0.0 ^b | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Hallucinogens | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 ^a | 0.2 |
| LSD | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | * | 0.0 | * | 0.0 | 0.0 |
| PCP | 0.0 | * | 0.0 | 0.0 | * | 0.0 | * | 0.0 | 0.0 | * | 0.0 |
| Ecstasy | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Inhalants | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 2.0 | 2.0 | 1.8 ^a | 1.9 | 2.2 | 2.2 | 1.9 | 2.1 | 2.2 | 1.9 | 2.1 |
| Pain Relievers | 1.3 | 1.3 | 1.2 ^a | 1.3 | 1.5 | 1.6 | 1.4 | 1.6 | 1.5 | 1.4 | 1.5 |
| OxyContin [®] | -- | -- | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Tranquilizers | 0.6 | 0.6 | 0.5 ^b | 0.6 | 0.5 ^a | 0.6 | 0.6 | 0.7 | 0.7 | 0.6 | 0.7 |
| Stimulants ³ | 0.4 | 0.4 | 0.4 | 0.3 | 0.4 | 0.3 | 0.2 | 0.4 | 0.3 | 0.3 | 0.3 |
| Methamphetamine ³ | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 ^a | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 |
| Sedatives | 0.2 ^a | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 2.7 | 2.6 | 2.3 ^a | 2.6 | 2.9 | 2.9 | 2.5 | 2.7 | 2.7 | 2.3 ^a | 2.8 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.13A Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|---------|
| ILLCIT DRUGS¹ | 100,612 ^b | 102,574 ^b | 102,491 ^b | 105,063 ^b | 104,733 ^b | 107,871 ^b | 111,250 ^b | 112,631 ^b | 113,655 ^b | 114,705 ^b | 118,764 |
| Marijuana and Hashish | 89,842 ^b | 91,723 ^b | 91,984 ^b | 93,143 ^b | 93,393 ^b | 96,641 ^b | 98,720 ^b | 100,747 ^b | 102,456 ^b | 103,464 ^a | 107,007 |
| Cocaine | 33,239 ^b | 34,243 ^b | 33,542 ^b | 33,090 ^b | 34,732 ^b | 35,402 ^a | 36,443 | 36,337 | 36,998 | 36,584 | 37,423 |
| Crack | 8,239 | 7,792 ^b | 7,715 ^b | 7,820 ^a | 8,430 | 8,509 | 8,374 | 8,332 | 9,148 | 8,146 | 8,981 |
| Heroin | 3,574 ^b | 3,664 ^b | 3,067 ^b | 3,475 ^b | 3,728 ^a | 3,751 ^a | 3,729 ^a | 3,623 ^b | 4,097 | 4,093 | 4,505 |
| Hallucinogens | 32,895 ^b | 33,113 ^b | 33,173 ^b | 32,730 ^b | 34,300 ^b | 33,380 ^b | 35,093 ^a | 36,388 | 36,543 | 35,432 | 37,085 |
| LSD | 23,845 | 24,027 | 23,085 | 22,145 | 23,146 | 22,518 | 23,333 | 23,464 | 23,143 | 22,771 | 23,487 |
| PCP | 7,185 | 6,903 | 6,587 | 6,421 | 6,505 | 6,073 | 6,554 | 6,173 | 6,179 | 6,027 | 6,456 |
| Ecstasy | 9,330 ^b | 10,301 ^b | 10,607 ^b | 11,096 ^b | 11,806 ^b | 11,971 ^b | 12,439 ^b | 13,707 ^b | 15,312 | 13,964 ^b | 15,673 |
| Inhalants | 20,265 | 20,325 | 20,036 | 20,087 | 20,298 | 20,130 | 20,104 | 20,269 | 19,763 | 18,646 | 19,482 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 44,556 ^b | 45,629 ^b | 45,754 ^b | 46,552 ^b | 47,809 ^b | 47,588 ^b | 49,371 ^a | 49,181 ^a | 49,241 ^a | 48,721 ^b | 51,908 |
| Pain Relievers | 26,836 ^b | 28,409 ^b | 28,890 ^b | 30,175 ^b | 30,828 ^b | 30,676 ^b | 32,639 ^b | 32,804 ^a | 32,664 ^b | 32,097 ^b | 34,977 |
| OxyContin [®] | 1,700 ^b | 2,579 ^b | 2,767 ^b | 3,205 ^b | 3,773 ^b | 4,019 ^b | 4,513 ^b | 5,483 ^a | 5,791 | 5,547 ^a | 6,257 |
| Tranquilizers | 18,433 ^b | 19,346 ^b | 19,036 ^b | 20,275 ^b | 20,576 ^b | 19,505 ^b | 20,831 ^b | 21,152 ^a | 21,463 ^a | 20,956 ^b | 22,936 |
| Stimulants ³ | 22,384 | 21,958 | 21,391 | 20,113 | 21,565 | 20,939 | 20,612 | 21,450 | 21,209 | 19,871 | 21,003 |
| Methamphetamine ³ | 14,949 ^b | 14,756 ^b | 14,166 ^b | 12,344 | 13,885 ^b | 12,859 | 12,438 | 12,721 | 12,903 | 11,743 | 12,114 |
| Sedatives | 9,723 ^b | 9,261 ^b | 9,640 ^b | 8,751 | 8,619 | 8,188 | 8,715 | 8,409 | 7,475 | 7,371 | 7,764 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 65,014 ^b | 65,809 ^b | 65,302 ^b | 66,879 ^b | 67,984 ^b | 69,005 ^b | 71,394 ^a | 71,579 ^a | 72,306 | 71,440 ^a | 74,343 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.13B Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|------|-------------------|------|
| ILLCIT DRUGS¹ | 47.8 ^b | 48.2 ^b | 47.6 ^b | 48.2 ^b | 47.5 ^b | 48.5 ^b | 49.5 | 49.6 | 49.6 | 49.3 ^a | 50.5 |
| Marijuana and Hashish | 42.7 ^b | 43.1 ^b | 42.7 ^b | 42.8 ^b | 42.3 ^b | 43.4 ^b | 43.9 ^b | 44.3 | 44.7 | 44.5 | 45.5 |
| Cocaine | 15.8 | 16.1 | 15.6 | 15.2 | 15.7 | 15.9 | 16.2 | 16.0 | 16.1 | 15.7 | 15.9 |
| Crack | 3.9 | 3.7 | 3.6 | 3.6 | 3.8 | 3.8 | 3.7 | 3.7 | 4.0 | 3.5 | 3.8 |
| Heroin | 1.7 | 1.7 | 1.4 ^b | 1.6 ^a | 1.7 | 1.7 | 1.7 | 1.6 ^a | 1.8 | 1.8 | 1.9 |
| Hallucinogens | 15.6 | 15.6 | 15.4 | 15.0 | 15.5 | 15.0 | 15.6 | 16.0 | 15.9 | 15.2 | 15.8 |
| LSD | 11.3 ^b | 11.3 ^b | 10.7 ^a | 10.2 | 10.5 | 10.1 | 10.4 | 10.3 | 10.1 | 9.8 | 10.0 |
| PCP | 3.4 ^b | 3.2 ^b | 3.1 | 2.9 | 2.9 | 2.7 | 2.9 | 2.7 | 2.7 | 2.6 | 2.7 |
| Ecstasy | 4.4 ^b | 4.8 ^b | 4.9 ^b | 5.1 ^b | 5.4 ^b | 5.4 ^b | 5.5 ^b | 6.0 ^b | 6.7 | 6.0 ^b | 6.7 |
| Inhalants | 9.6 ^b | 9.6 ^b | 9.3 ^b | 9.2 ^b | 9.2 ^b | 9.0 ^a | 8.9 ^a | 8.9 ^a | 8.6 | 8.0 | 8.3 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 21.2 | 21.5 | 21.3 | 21.4 | 21.7 | 21.4 | 22.0 | 21.6 | 21.5 | 20.9 ^a | 22.1 |
| Pain Relievers | 12.8 ^b | 13.4 ^b | 13.4 ^b | 13.9 ^b | 14.0 ^b | 13.8 ^b | 14.5 | 14.4 | 14.2 | 13.8 ^b | 14.9 |
| OxyContin [®] | 0.8 ^b | 1.2 ^b | 1.3 ^b | 1.5 ^b | 1.7 ^b | 1.8 ^b | 2.0 ^b | 2.4 | 2.5 | 2.4 ^a | 2.7 |
| Tranquilizers | 8.8 ^b | 9.1 ^a | 8.8 ^b | 9.3 | 9.3 | 8.8 ^b | 9.3 | 9.3 | 9.4 | 9.0 ^a | 9.8 |
| Stimulants ³ | 10.6 ^b | 10.3 ^b | 9.9 ^b | 9.2 | 9.8 ^a | 9.4 | 9.2 | 9.4 | 9.3 | 8.5 | 8.9 |
| Methamphetamine ³ | 7.1 ^b | 6.9 ^b | 6.6 ^b | 5.7 ^a | 6.3 ^b | 5.8 ^a | 5.5 | 5.6 | 5.6 | 5.0 | 5.2 |
| Sedatives | 4.6 ^b | 4.4 ^b | 4.5 ^b | 4.0 ^b | 3.9 ^b | 3.7 | 3.9 ^a | 3.7 | 3.3 | 3.2 | 3.3 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 30.9 | 30.9 | 30.3 ^a | 30.7 | 30.8 | 31.0 | 31.7 | 31.5 | 31.5 | 30.7 | 31.6 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.14A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 29,638 ^b | 29,538 ^b | 29,506 ^b | 29,999 ^b | 30,831 ^b | 31,052 ^b | 30,892 ^b | 33,285 ^b | 34,163 ^b | 33,552 ^b | 37,011 |
| Marijuana and Hashish | 21,850 ^b | 21,486 ^b | 21,796 ^b | 21,991 ^b | 22,077 ^b | 21,978 ^b | 22,633 ^b | 25,325 ^b | 25,887 ^b | 26,186 ^b | 28,150 |
| Cocaine | 5,394 ^b | 5,470 ^b | 5,247 ^a | 5,099 ^a | 5,663 ^b | 5,362 ^b | 4,977 | 4,564 | 4,294 | 3,628 ^b | 4,491 |
| Crack | 1,454 ^b | 1,310 ^a | 1,241 ^a | 1,327 ^b | 1,404 ^b | 1,387 ^b | 1,079 | 979 | 866 | 597 ^a | 902 |
| Heroin | 353 ^b | 278 ^b | 350 ^b | 343 ^b | 524 | 349 ^b | 415 ^a | 547 | 593 | 566 | 638 |
| Hallucinogens | 3,804 | 3,158 ^b | 3,125 ^b | 3,144 ^b | 3,317 | 3,130 ^b | 2,982 ^b | 3,799 | 3,795 | 3,421 | 3,753 |
| LSD | 684 ^a | 409 ^b | 434 ^b | 413 ^b | 561 ^b | 495 ^b | 623 ^b | 647 ^b | 746 | 721 | 905 |
| PCP | 126 | 130 | 138 | 79 | 129 | 83 | 37 ^b | 77 | 61 | 80 | 124 |
| Ecstasy | 2,621 | 1,790 ^b | 1,605 ^b | 1,703 ^b | 1,830 ^b | 1,794 ^b | 1,770 ^b | 2,388 | 2,188 | 2,003 | 2,322 |
| Inhalants | 1,005 | 956 | 1,087 | 1,043 | 1,110 | 1,090 | 1,079 | 1,149 | 1,163 | 1,041 | 1,046 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 12,509 ^b | 12,868 ^b | 12,621 ^b | 13,246 ^b | 14,298 | 14,297 | 13,271 ^b | 14,169 | 14,238 | 12,908 ^b | 15,025 |
| Pain Relievers | 9,122 ^b | 9,753 ^b | 9,391 ^b | 10,074 ^a | 10,829 | 10,833 | 10,304 | 10,822 | 10,720 | 9,659 ^b | 11,157 |
| OxyContin [®] | -- | -- | 1,019 ^a | 1,039 ^a | 1,124 | 1,195 | 1,236 | 1,460 | 1,645 ^a | 1,404 | 1,290 |
| Tranquilizers | 4,281 ^b | 4,465 ^b | 4,536 ^b | 4,752 ^b | 4,581 ^b | 4,788 ^a | 4,616 ^b | 5,031 | 5,117 | 4,635 ^b | 5,621 |
| Stimulants ³ | 2,716 | 2,453 ^b | 2,747 | 2,581 ^a | 3,241 | 2,579 | 2,275 ^b | 2,759 | 2,577 | 2,401 ^b | 2,992 |
| Methamphetamine ³ | 1,510 ^b | 1,417 ^a | 1,631 ^b | 1,425 ^a | 1,700 ^b | 1,229 | 760 ^a | 1,080 | 861 | 928 | 1,073 |
| Sedatives | 843 ^a | 700 | 604 | 632 | 818 ^b | 754 ^a | 513 | 690 | 796 ^a | 442 | 509 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 17,086 ^b | 16,954 ^b | 16,389 ^b | 16,980 ^b | 18,114 | 18,283 | 17,159 ^b | 18,228 | 17,866 | 16,395 ^b | 18,970 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.14B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 14.1 ^b | 13.9 ^b | 13.7 ^b | 13.8 ^b | 14.0 ^b | 13.9 ^b | 13.7 ^b | 14.6 ^b | 14.9 ^a | 14.4 ^b | 15.7 |
| Marijuana and Hashish | 10.4 ^b | 10.1 ^b | 10.1 ^b | 10.1 ^b | 10.0 ^b | 9.9 ^b | 10.1 ^b | 11.1 ^a | 11.3 ^a | 11.3 ^a | 12.0 |
| Cocaine | 2.6 ^b | 2.6 ^b | 2.4 ^b | 2.3 ^b | 2.6 ^b | 2.4 ^b | 2.2 ^a | 2.0 | 1.9 | 1.6 ^b | 1.9 |
| Crack | 0.7 ^b | 0.6 ^b | 0.5 | 0.4 | 0.4 | 0.3 ^a | 0.4 |
| Heroin | 0.2 ^a | 0.1 ^b | 0.2 ^b | 0.2 ^b | 0.2 | 0.2 ^b | 0.2 ^a | 0.2 | 0.3 | 0.2 | 0.3 |
| Hallucinogens | 1.8 ^a | 1.5 | 1.5 | 1.4 | 1.5 | 1.4 ^a | 1.3 ^b | 1.7 | 1.7 | 1.5 | 1.6 |
| LSD | 0.3 | 0.2 ^b | 0.2 ^b | 0.2 ^b | 0.3 ^b | 0.2 ^b | 0.3 ^b | 0.3 ^a | 0.3 | 0.3 | 0.4 |
| PCP | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 ^a | 0.0 | 0.0 | 0.0 | 0.1 |
| Ecstasy | 1.2 ^b | 0.8 ^a | 0.7 ^b | 0.8 ^b | 0.8 ^a | 0.8 ^a | 0.8 ^b | 1.1 | 1.0 | 0.9 | 1.0 |
| Inhalants | 0.5 | 0.4 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 | 0.4 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 5.9 | 6.1 | 5.9 ^a | 6.1 | 6.5 | 6.4 | 5.9 ^a | 6.2 | 6.2 | 5.5 ^b | 6.4 |
| Pain Relievers | 4.3 ^a | 4.6 | 4.4 | 4.6 | 4.9 | 4.9 | 4.6 | 4.8 | 4.7 | 4.2 ^b | 4.7 |
| OxyContin [®] | -- | -- | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.7 ^a | 0.6 | 0.5 |
| Tranquilizers | 2.0 ^a | 2.1 ^a | 2.1 ^a | 2.2 | 2.1 ^a | 2.2 | 2.1 ^a | 2.2 | 2.2 | 2.0 ^b | 2.4 |
| Stimulants ³ | 1.3 | 1.2 | 1.3 | 1.2 | 1.5 | 1.2 | 1.0 ^b | 1.2 | 1.1 | 1.0 ^a | 1.3 |
| Methamphetamine ³ | 0.7 ^b | 0.7 ^b | 0.8 ^b | 0.7 ^b | 0.8 ^b | 0.6 | 0.3 ^a | 0.5 | 0.4 | 0.4 | 0.5 |
| Sedatives | 0.4 ^b | 0.3 ^a | 0.3 | 0.3 | 0.4 ^b | 0.3 ^a | 0.2 | 0.3 ^a | 0.3 ^a | 0.2 | 0.2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 8.1 | 8.0 | 7.6 | 7.8 | 8.2 | 8.2 | 7.6 | 8.0 | 7.8 | 7.0 ^b | 8.1 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.15A Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------|
| ILLCIT DRUGS¹ | 16,644 ^b | 16,659 ^b | 16,397 ^b | 17,209 ^b | 17,898 ^b | 17,477 ^b | 17,811 ^b | 19,446 ^b | 20,180 | 19,932 ^a | 21,486 |
| Marijuana and Hashish | 12,561 ^b | 12,667 ^b | 12,667 ^b | 12,896 ^b | 13,154 ^b | 12,777 ^b | 13,601 ^b | 15,013 ^b | 15,607 ^a | 16,103 | 17,055 |
| Cocaine | 1,874 | 2,128 ^a | 1,895 | 2,255 ^b | 2,316 ^b | 1,972 | 1,767 | 1,572 | 1,412 | 1,302 | 1,620 |
| Crack | 541 | 572 | 451 | 657 | 691 | 595 | 356 | 489 | 375 | 225 ^a | 443 |
| Heroin | 155 ^b | 101 ^b | 151 ^b | 121 ^b | 323 | 157 ^b | 200 ^a | 180 ^a | 231 | 267 | 335 |
| Hallucinogens | 941 | 783 | 719 ^a | 883 | 826 | 824 | 814 | 1,047 | 992 | 757 ^a | 974 |
| LSD | 51 ^b | 82 ^a | 103 | 72 ^a | 99 | 114 | 105 | 124 | 109 | 118 | 158 |
| PCP | 35 | 35 | 37 | 31 | 19 | 34 | * | 42 | 30 | 22 | 27 |
| Ecstasy | 553 | 379 ^a | 373 ^a | 428 | 446 | 438 | 441 | 647 | 567 | 435 | 554 |
| Inhalants | 334 | 250 | 325 | 298 | 422 | 323 | 374 | 318 | 428 | 402 | 305 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 5,300 ^a | 5,453 | 5,196 ^b | 5,653 | 6,252 | 6,077 | 5,499 | 6,205 | 6,236 | 5,431 ^a | 6,138 |
| Pain Relievers | 3,583 ^b | 3,892 | 3,653 ^a | 3,972 | 4,534 | 4,517 | 4,165 | 4,624 | 4,485 | 3,908 | 4,304 |
| OxyContin [®] | -- | -- | 260 | 299 | 246 | 324 | 387 | 440 | 519 ^b | 383 | 310 |
| Tranquilizers | 1,617 | 1,604 ^a | 1,455 ^b | 1,657 | 1,636 | 1,657 | 1,646 | 1,887 | 2,026 | 1,678 | 1,999 |
| Stimulants ³ | 1,094 | 1,088 | 1,132 | 1,016 | 1,214 | 924 | 796 ^a | 1,172 | 975 | 872 | 1,053 |
| Methamphetamine ³ | 612 | 655 ^a | 648 ^a | 556 | 678 ^a | 496 | 285 | 466 | 328 | 402 | 409 |
| Sedatives | 389 | 247 | 239 | 245 | 340 | 311 | 203 | 327 | 340 | 206 | 244 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 7,356 | 7,432 | 6,920 ^b | 7,710 | 8,368 | 8,113 | 7,463 | 8,073 | 7,924 | 6,994 ^a | 7,965 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.15B Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------|
| ILLCIT DRUGS¹ | 7.9 ^b | 7.8 ^b | 7.6 ^b | 7.9 ^b | 8.1 ^b | 7.9 ^b | 7.9 ^b | 8.6 ^a | 8.8 | 8.6 ^a | 9.1 |
| Marijuana and Hashish | 6.0 ^b | 6.0 ^b | 5.9 ^b | 5.9 ^b | 6.0 ^b | 5.7 ^b | 6.0 ^b | 6.6 ^b | 6.8 | 6.9 | 7.3 |
| Cocaine | 0.9 ^a | 1.0 ^b | 0.9 ^a | 1.0 ^b | 1.0 ^b | 0.9 ^a | 0.8 | 0.7 | 0.6 | 0.6 | 0.7 |
| Crack | 0.3 | 0.3 | 0.2 | 0.3 ^a | 0.3 ^a | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 ^a | 0.2 |
| Heroin | 0.1 ^a | 0.0 ^b | 0.1 ^b | 0.1 ^b | 0.1 | 0.1 ^b | 0.1 ^a | 0.1 ^a | 0.1 | 0.1 | 0.1 |
| Hallucinogens | 0.4 | 0.4 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 0.4 | 0.3 ^a | 0.4 |
| LSD | 0.0 ^b | 0.0 | 0.0 | 0.0 ^a | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 0.1 | 0.1 |
| PCP | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | * | 0.0 | 0.0 | 0.0 | 0.0 |
| Ecstasy | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 |
| Inhalants | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 2.5 | 2.6 | 2.4 | 2.6 | 2.8 | 2.7 | 2.4 | 2.7 | 2.7 | 2.3 | 2.6 |
| Pain Relievers | 1.7 | 1.8 | 1.7 | 1.8 | 2.1 | 2.0 | 1.9 | 2.0 | 2.0 | 1.7 | 1.8 |
| OxyContin [®] | -- | -- | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 ^b | 0.2 | 0.1 |
| Tranquilizers | 0.8 | 0.8 | 0.7 ^a | 0.8 | 0.7 | 0.7 | 0.7 | 0.8 | 0.9 | 0.7 | 0.9 |
| Stimulants ³ | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.4 | 0.4 | 0.5 | 0.4 | 0.4 | 0.4 |
| Methamphetamine ³ | 0.3 ^a | 0.3 ^b | 0.3 ^b | 0.3 | 0.3 ^b | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 |
| Sedatives | 0.2 ^a | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 3.5 | 3.5 | 3.2 | 3.5 | 3.8 ^a | 3.6 | 3.3 | 3.6 | 3.5 | 3.0 ^a | 3.4 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.16A Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|
| ILLCIT DRUGS¹ | 4,035 ^b | 3,968 ^b | 3,956 ^b | 3,679 ^b | 3,658 ^b | 3,501 ^b | 3,335 ^a | 3,524 ^b | 3,256 | 3,336 ^a | 3,123 |
| Marijuana and Hashish | 2,714 ^b | 2,569 ^b | 2,552 ^b | 2,348 | 2,315 | 2,233 | 2,203 | 2,352 | 2,269 | 2,387 | 2,266 |
| Cocaine | 337 ^b | 319 ^b | 339 ^b | 283 ^b | 257 ^b | 258 ^b | 218 ^b | 220 ^b | 190 ^a | 175 | 129 |
| Crack | 61 ^b | 89 ^b | 60 ^b | 48 ^b | 51 ^b | 43 ^b | 32 | 28 | 34 | 36 | 15 |
| Heroin | 34 | 34 | 43 | 32 | 25 | 28 | 37 | 25 | 23 | 33 | 27 |
| Hallucinogens | 740 ^b | 618 ^b | 616 ^b | 518 ^b | 512 ^b | 480 | 501 ^a | 537 ^b | 516 ^b | 496 ^a | 405 |
| LSD | 364 ^b | 206 ^b | 174 ^a | 177 ^a | 125 | 111 | 149 | 140 | 120 | 137 | 128 |
| PCP | 106 ^b | 100 ^b | 97 ^b | 97 ^b | 53 | 53 | 57 | 44 | 27 | 40 | 45 |
| Ecstasy | 376 ^b | 257 | 243 | 181 | 202 | 215 | 224 | 277 | 277 ^a | 298 ^b | 214 |
| Inhalants | 1,426 ^b | 1,390 ^b | 1,429 ^b | 1,360 ^b | 1,302 ^b | 1,225 ^b | 1,081 ^b | 1,109 ^b | 985 ^b | 909 ^b | 749 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 1,643 ^b | 1,639 ^b | 1,616 ^b | 1,444 ^b | 1,561 ^b | 1,441 ^b | 1,263 | 1,352 ^a | 1,198 | 1,215 | 1,209 |
| Pain Relievers | 1,362 ^b | 1,391 ^b | 1,378 ^b | 1,211 ^b | 1,315 ^b | 1,200 ^b | 1,068 | 1,178 ^a | 1,053 | 1,027 | 1,029 |
| OxyContin [®] | 112 ^a | 146 | 162 | 136 | 160 | 173 | 159 | 189 | 184 | 165 | 157 |
| Tranquilizers | 385 ^a | 398 ^a | 353 | 342 | 316 | 323 | 317 | 303 | 325 | 315 | 309 |
| Stimulants ³ | 469 ^b | 472 ^b | 387 ^b | 364 ^b | 399 ^b | 302 ^b | 248 | 234 | 230 | 232 | 203 |
| Methamphetamine ³ | 150 ^b | 160 ^b | 140 ^b | 133 ^b | 126 ^b | 81 | 65 | 62 | 76 | 81 | 59 |
| Sedatives | 84 | 126 ^b | 128 ^b | 110 | 97 | 89 | 90 | 103 | 77 | 66 | 73 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 2,726 ^b | 2,711 ^b | 2,721 ^b | 2,488 ^b | 2,527 ^b | 2,378 ^b | 2,179 ^b | 2,252 ^b | 2,030 ^b | 2,008 ^a | 1,811 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.16B Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|
| ILLCIT DRUGS¹ | 31.9 ^b | 31.1 ^b | 30.7 ^b | 28.4 ^b | 28.2 ^b | 27.2 ^b | 26.2 ^a | 28.0 ^b | 26.1 ^a | 26.1 ^a | 24.5 |
| Marijuana and Hashish | 21.5 ^b | 20.1 ^b | 19.8 ^b | 18.1 | 17.9 | 17.3 | 17.3 | 18.7 | 18.2 | 18.7 | 17.8 |
| Cocaine | 2.7 ^b | 2.5 ^b | 2.6 ^b | 2.2 ^b | 2.0 ^b | 2.0 ^b | 1.7 ^b | 1.8 ^b | 1.5 ^a | 1.4 | 1.0 |
| Crack | 0.5 ^b | 0.7 ^b | 0.5 ^b | 0.4 ^b | 0.4 ^b | 0.3 ^a | 0.2 | 0.2 | 0.3 | 0.3 | 0.1 |
| Heroin | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 |
| Hallucinogens | 5.8 ^b | 4.8 ^b | 4.8 ^b | 4.0 ^a | 4.0 ^a | 3.7 | 3.9 ^a | 4.3 ^b | 4.1 ^b | 3.9 ^a | 3.2 |
| LSD | 2.9 ^b | 1.6 ^b | 1.4 | 1.4 | 1.0 | 0.9 | 1.2 | 1.1 | 1.0 | 1.1 | 1.0 |
| PCP | 0.8 ^b | 0.8 ^b | 0.8 ^b | 0.7 ^b | 0.4 | 0.4 | 0.5 | 0.3 | 0.2 | 0.3 | 0.4 |
| Ecstasy | 3.0 ^b | 2.0 | 1.9 | 1.4 | 1.6 | 1.7 | 1.8 | 2.2 ^a | 2.2 ^a | 2.3 ^a | 1.7 |
| Inhalants | 11.3 ^b | 10.9 ^b | 11.1 ^b | 10.5 ^b | 10.0 ^b | 9.5 ^b | 8.5 ^b | 8.8 ^b | 7.9 ^b | 7.1 ^b | 5.9 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 13.0 ^b | 12.8 ^b | 12.6 ^b | 11.1 ^b | 12.0 ^b | 11.2 ^b | 9.9 | 10.8 ^a | 9.6 | 9.5 | 9.5 |
| Pain Relievers | 10.8 ^b | 10.9 ^b | 10.7 ^b | 9.4 ^b | 10.1 ^b | 9.3 ^a | 8.4 | 9.4 ^b | 8.5 | 8.0 | 8.1 |
| OxyContin [®] | 0.9 ^a | 1.1 | 1.3 | 1.0 | 1.2 | 1.3 | 1.3 | 1.5 | 1.5 | 1.3 | 1.2 |
| Tranquilizers | 3.0 ^a | 3.1 ^a | 2.7 | 2.6 | 2.4 | 2.5 | 2.5 | 2.4 | 2.6 | 2.5 | 2.4 |
| Stimulants ³ | 3.7 ^b | 3.7 ^b | 3.0 ^b | 2.8 ^b | 3.1 ^b | 2.3 ^b | 2.0 | 1.9 | 1.8 | 1.8 | 1.6 |
| Methamphetamine ³ | 1.2 ^b | 1.3 ^b | 1.1 ^b | 1.0 ^b | 1.0 ^b | 0.6 | 0.5 | 0.5 | 0.6 | 0.6 | 0.5 |
| Sedatives | 0.7 | 1.0 ^b | 1.0 ^b | 0.8 | 0.7 | 0.7 | 0.7 | 0.8 | 0.6 | 0.5 | 0.6 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 21.6 ^b | 21.2 ^b | 21.1 ^b | 19.2 ^b | 19.5 ^b | 18.4 ^b | 17.1 ^b | 17.9 ^b | 16.3 ^b | 15.7 ^a | 14.2 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.17A Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|-------|
| ILLCIT DRUGS¹ | 3,607 ^b | 3,663 ^b | 3,610 ^b | 3,342 ^b | 3,356 ^b | 3,137 ^a | 3,230 ^b | 3,104 | 3,022 | 3,037 | 2,921 |
| Marijuana and Hashish | 2,391 ^b | 2,318 ^b | 2,236 ^b | 2,055 | 2,084 | 1,863 | 1,935 | 1,852 | 1,888 | 1,991 | 1,966 |
| Cocaine | 333 ^b | 330 ^b | 272 ^b | 300 ^b | 313 ^b | 287 ^b | 254 ^b | 185 ^a | 173 | 162 | 136 |
| Crack | 102 ^b | 68 ^b | 66 ^b | 60 ^b | 78 ^b | 64 ^b | 48 ^a | 30 | 26 | 31 | 19 |
| Heroin | 60 | 45 | 35 | 28 | 35 | 27 | 30 | 32 | 23 | 37 | 32 |
| Hallucinogens | 680 ^b | 632 ^b | 543 ^b | 480 | 477 | 418 | 470 | 479 | 485 | 435 | 418 |
| LSD | 306 ^b | 191 ^b | 139 | 111 | 94 | 91 | 122 | 109 | 111 | 92 | 121 |
| PCP | 127 ^b | 104 ^b | 77 | 85 ^a | 75 | 52 | 45 | 55 | 49 | 37 | 51 |
| Ecstasy | 444 ^b | 346 ^a | 280 | 218 | 274 | 231 | 301 | 295 | 340 | 307 | 274 |
| Inhalants | 1,179 ^b | 1,281 ^b | 1,333 ^b | 1,298 ^b | 1,257 ^b | 1,203 ^b | 1,245 ^b | 1,174 ^b | 1,030 ^b | 968 | 864 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 1,759 ^b | 1,733 ^b | 1,787 ^b | 1,575 ^b | 1,613 ^b | 1,513 ^b | 1,521 ^b | 1,458 ^b | 1,392 | 1,307 | 1,273 |
| Pain Relievers | 1,412 ^b | 1,407 ^b | 1,500 ^b | 1,305 ^b | 1,335 ^b | 1,257 ^b | 1,286 ^b | 1,215 ^b | 1,191 ^a | 1,123 | 1,040 |
| OxyContin [®] | 112 ^a | 108 ^b | 143 | 140 | 184 | 187 | 185 | 191 | 185 | 204 | 165 |
| Tranquilizers | 450 | 476 ^a | 464 | 424 | 426 | 425 | 403 | 392 | 398 | 385 | 394 |
| Stimulants ³ | 643 ^b | 574 ^b | 519 ^b | 506 ^b | 531 ^b | 428 ^b | 425 ^b | 349 | 301 | 276 | 292 |
| Methamphetamine ³ | 266 ^b | 222 ^b | 206 ^b | 185 ^b | 216 ^b | 140 ^b | 130 | 125 | 81 | 103 | 86 |
| Sedatives | 154 ^a | 123 | 123 | 122 | 126 | 124 | 115 | 121 | 101 | 77 | 102 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 2,559 ^b | 2,607 ^b | 2,634 ^b | 2,456 ^b | 2,424 ^b | 2,271 ^b | 2,334 ^b | 2,250 ^b | 2,136 ^b | 1,999 | 1,881 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.17B Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------|------|
| ILLCIT DRUGS¹ | 29.8 ^b | 30.0 ^b | 29.3 ^b | 26.9 ^b | 27.0 ^b | 25.4 | 26.5 ^b | 25.8 ^a | 25.4 | 24.9 | 24.0 |
| Marijuana and Hashish | 19.7 ^b | 19.0 ^b | 18.1 ^b | 16.6 | 16.8 | 15.1 | 15.9 | 15.4 | 15.9 | 16.3 | 16.1 |
| Cocaine | 2.8 ^b | 2.7 ^b | 2.2 ^b | 2.4 ^b | 2.5 ^b | 2.3 ^b | 2.1 ^b | 1.5 ^a | 1.5 | 1.3 | 1.1 |
| Crack | 0.8 ^b | 0.6 ^b | 0.5 ^b | 0.5 ^b | 0.6 ^b | 0.5 ^b | 0.4 ^a | 0.3 | 0.2 | 0.3 | 0.2 |
| Heroin | 0.5 | 0.4 | 0.3 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.3 | 0.3 |
| Hallucinogens | 5.6 ^b | 5.2 ^b | 4.4 ^b | 3.9 | 3.8 | 3.4 | 3.9 | 4.0 | 4.1 | 3.6 | 3.4 |
| LSD | 2.5 ^b | 1.6 ^b | 1.1 | 0.9 | 0.8 | 0.7 | 1.0 | 0.9 | 0.9 | 0.8 | 1.0 |
| PCP | 1.0 ^b | 0.8 ^b | 0.6 | 0.7 ^a | 0.6 | 0.4 | 0.4 | 0.5 | 0.4 | 0.3 | 0.4 |
| Ecstasy | 3.7 ^b | 2.8 ^a | 2.3 | 1.8 ^a | 2.2 | 1.9 | 2.5 | 2.5 | 2.9 ^a | 2.5 | 2.2 |
| Inhalants | 9.7 ^b | 10.5 ^b | 10.8 ^b | 10.5 ^b | 10.1 ^b | 9.7 ^b | 10.2 ^b | 9.7 ^b | 8.7 ^b | 7.9 | 7.1 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 14.5 ^b | 14.2 ^b | 14.5 ^b | 12.7 ^b | 13.0 ^b | 12.2 ^b | 12.5 ^b | 12.1 ^b | 11.7 ^a | 10.7 | 10.4 |
| Pain Relievers | 11.7 ^b | 11.5 ^b | 12.2 ^b | 10.5 ^b | 10.7 ^b | 10.2 ^b | 10.6 ^b | 10.1 ^b | 10.0 ^b | 9.2 | 8.5 |
| OxyContin [®] | 0.9 ^a | 0.9 ^b | 1.2 | 1.1 | 1.5 | 1.5 | 1.5 | 1.6 | 1.6 | 1.7 | 1.4 |
| Tranquilizers | 3.7 | 3.9 ^a | 3.8 | 3.4 | 3.4 | 3.4 | 3.3 | 3.3 | 3.4 | 3.2 | 3.2 |
| Stimulants ³ | 5.3 ^b | 4.7 ^b | 4.2 ^b | 4.1 ^b | 4.3 ^b | 3.5 ^b | 3.5 ^b | 2.9 | 2.5 | 2.3 | 2.4 |
| Methamphetamine ³ | 2.2 ^b | 1.8 ^b | 1.7 ^b | 1.5 ^b | 1.7 ^b | 1.1 ^b | 1.1 | 1.0 ^a | 0.7 | 0.8 | 0.7 |
| Sedatives | 1.3 ^a | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 0.9 | 1.0 | 0.9 | 0.6 | 0.8 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 21.1 ^b | 21.3 ^b | 21.4 ^b | 19.8 ^b | 19.5 ^b | 18.4 ^b | 19.2 ^b | 18.7 ^b | 18.0 ^b | 16.4 | 15.4 |

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.18A Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------|
| ILLCIT DRUGS¹ | 2,842 ^b | 2,776 ^b | 2,653 ^b | 2,557 ^b | 2,531 ^b | 2,507 ^a | 2,433 | 2,556 ^b | 2,422 | 2,438 | 2,268 |
| Marijuana and Hashish | 2,064 ^b | 1,955 ^a | 1,918 | 1,796 | 1,752 | 1,754 | 1,738 | 1,898 | 1,861 | 1,933 | 1,799 |
| Cocaine | 263 ^b | 222 ^b | 223 ^b | 208 ^b | 179 ^b | 174 ^b | 161 ^b | 146 ^b | 116 | 115 | 88 |
| Crack | 36 ^b | 47 ^b | 31 ^b | 23 | 32 ^a | 25 ^a | 19 | 16 | 8 | 16 | 8 |
| Heroin | 20 | 17 | 32 | 17 | 14 | 16 | 27 | 17 | 13 | 22 | 15 |
| Hallucinogens | 501 ^b | 386 ^b | 389 ^b | 352 ^a | 320 | 342 | 377 ^b | 407 ^b | 382 ^b | 369 ^a | 285 |
| LSD | 183 ^b | 73 | 94 | 92 | 51 ^a | 76 | 119 ^a | 90 | 73 | 98 | 82 |
| PCP | 46 ^a | 44 ^a | 39 | 44 | 26 | 32 | 37 | 26 | 12 | 19 | 20 |
| Ecstasy | 262 ^b | 142 | 141 | 125 | 130 | 165 | 160 | 212 ^b | 206 ^b | 218 ^b | 123 |
| Inhalants | 586 ^b | 554 ^b | 587 ^b | 538 ^b | 520 ^b | 501 ^b | 470 ^b | 458 ^b | 381 ^b | 396 ^b | 271 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 1,091 ^b | 1,090 ^b | 974 ^b | 962 ^b | 1,055 ^b | 973 ^b | 867 | 878 | 806 | 829 | 777 |
| Pain Relievers | 913 ^b | 930 ^b | 837 ^b | 818 ^b | 885 ^b | 819 ^b | 730 | 755 ^a | 688 | 693 | 653 |
| OxyContin [®] | -- | -- | 99 | 101 | 96 | 107 | 122 | 110 | 113 | 102 | 98 |
| Tranquilizers | 249 ^a | 258 ^a | 226 | 229 | 196 | 214 | 226 | 210 | 200 | 216 | 184 |
| Stimulants ³ | 294 ^b | 253 ^b | 210 ^a | 194 | 224 ^b | 174 | 140 | 138 | 133 | 134 | 147 |
| Methamphetamine ³ | 97 ^b | 82 ^b | 69 ^a | 78 ^a | 74 ^a | 39 | 33 | 29 | 44 | 44 | 38 |
| Sedatives | 45 | 67 ^b | 59 ^a | 51 | 56 ^a | 42 | 47 | 70 ^b | 41 | 32 | 29 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 1,683 ^b | 1,637 ^b | 1,537 ^b | 1,479 ^b | 1,543 ^b | 1,450 ^b | 1,383 ^b | 1,376 ^b | 1,265 ^b | 1,263 ^b | 1,087 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.18B Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|------|
| ILLCIT DRUGS¹ | 22.5 ^b | 21.7 ^b | 20.6 ^b | 19.7 ^b | 19.5 ^a | 19.4 ^a | 19.1 ^a | 20.3 ^b | 19.4 ^a | 19.1 | 17.8 |
| Marijuana and Hashish | 16.3 ^b | 15.3 | 14.9 | 13.9 | 13.5 | 13.6 | 13.7 | 15.1 | 14.9 | 15.1 | 14.1 |
| Cocaine | 2.1 ^b | 1.7 ^b | 1.7 ^b | 1.6 ^b | 1.4 ^b | 1.4 ^b | 1.3 ^b | 1.2 ^b | 0.9 | 0.9 | 0.7 |
| Crack | 0.3 ^b | 0.4 ^b | 0.2 ^b | 0.2 | 0.2 ^a | 0.2 ^a | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Heroin | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 |
| Hallucinogens | 4.0 ^b | 3.0 ^b | 3.0 ^b | 2.7 | 2.5 | 2.7 | 3.0 ^b | 3.2 ^b | 3.1 ^b | 2.9 ^a | 2.2 |
| LSD | 1.5 ^b | 0.6 | 0.7 | 0.7 | 0.4 ^a | 0.6 | 0.9 ^a | 0.7 | 0.6 | 0.8 | 0.6 |
| PCP | 0.4 ^a | 0.3 ^a | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.1 | 0.2 | 0.2 |
| Ecstasy | 2.1 ^b | 1.1 | 1.1 | 1.0 | 1.0 | 1.3 | 1.3 | 1.7 ^b | 1.7 ^b | 1.7 ^b | 1.0 |
| Inhalants | 4.6 ^b | 4.3 ^b | 4.6 ^b | 4.2 ^b | 4.0 ^b | 3.9 ^b | 3.7 ^b | 3.6 ^b | 3.1 ^b | 3.1 ^b | 2.1 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 8.6 ^b | 8.5 ^b | 7.6 ^b | 7.4 ^b | 8.1 ^b | 7.5 ^b | 6.8 | 7.0 ^a | 6.5 | 6.5 | 6.1 |
| Pain Relievers | 7.2 ^b | 7.3 ^b | 6.5 ^b | 6.3 ^b | 6.8 ^b | 6.4 ^b | 5.7 | 6.0 ^a | 5.5 | 5.4 | 5.1 |
| OxyContin [®] | -- | -- | 0.8 | 0.8 | 0.7 | 0.8 | 1.0 | 0.9 | 0.9 | 0.8 | 0.8 |
| Tranquilizers | 2.0 ^a | 2.0 ^a | 1.8 | 1.8 | 1.5 | 1.7 | 1.8 | 1.7 | 1.6 | 1.7 | 1.4 |
| Stimulants ³ | 2.3 ^b | 2.0 ^b | 1.6 ^a | 1.5 | 1.7 ^b | 1.4 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 |
| Methamphetamine ³ | 0.8 ^b | 0.6 ^b | 0.5 ^a | 0.6 ^a | 0.6 ^a | 0.3 | 0.3 | 0.2 | 0.4 | 0.3 | 0.3 |
| Sedatives | 0.4 | 0.5 ^b | 0.5 ^a | 0.4 | 0.4 ^a | 0.3 | 0.4 | 0.6 ^b | 0.3 | 0.2 | 0.2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 13.3 ^b | 12.8 ^b | 11.9 ^b | 11.4 ^b | 11.9 ^b | 11.2 ^b | 10.9 ^b | 10.9 ^b | 10.2 ^b | 9.9 ^b | 8.5 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.19A Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|------------------|-------|
| ILLCIT DRUGS¹ | 2,652 ^b | 2,679 ^b | 2,648 ^b | 2,484 ^b | 2,447 ^b | 2,231 | 2,318 | 2,282 | 2,322 | 2,297 | 2,201 |
| Marijuana and Hashish | 1,841 ^b | 1,790 ^b | 1,738 ^a | 1,587 | 1,603 | 1,404 ^a | 1,520 | 1,464 | 1,553 | 1,620 | 1,565 |
| Cocaine | 245 ^b | 216 ^b | 188 ^b | 215 ^b | 232 ^b | 211 ^b | 145 ^a | 96 | 124 | 113 | 92 |
| Crack | 64 ^b | 48 ^b | 32 ^a | 31 ^a | 41 ^b | 40 ^b | 16 | 13 | 10 | 12 | 11 |
| Heroin | 31 | 19 | 16 | 20 | 23 | 8 | 14 | 17 | 15 | 32 | 16 |
| Hallucinogens | 444 ^b | 392 ^b | 364 ^b | 313 | 333 | 287 | 312 | 330 | 365 ^b | 279 | 269 |
| LSD | 131 ^b | 76 | 63 | 59 | 55 | 50 | 66 | 53 | 62 | 61 | 70 |
| PCP | 64 ^b | 44 | 33 | 41 | 33 | 24 | 25 | 22 | 23 | 20 | 28 |
| Ecstasy | 285 ^b | 188 | 169 | 131 | 177 | 163 | 186 | 204 | 258 ^b | 201 | 164 |
| Inhalants | 492 ^b | 565 ^b | 581 ^b | 606 ^b | 589 ^b | 494 ^b | 516 ^b | 502 ^b | 504 ^b | 424 | 376 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 1,194 ^b | 1,205 ^b | 1,254 ^b | 1,139 ^b | 1,129 ^b | 1,064 ^b | 1,052 ^b | 1,017 ^b | 1,006 ^a | 920 | 865 |
| Pain Relievers | 957 ^b | 987 ^b | 1,028 ^b | 922 ^b | 937 ^b | 866 ^b | 883 ^b | 873 ^b | 833 ^b | 791 ^a | 680 |
| OxyContin [®] | -- | -- | 95 | 87 | 105 | 125 | 104 | 115 | 121 | 117 | 88 |
| Tranquilizers | 319 | 328 | 307 | 268 | 288 | 291 | 257 | 252 | 266 | 259 | 268 |
| Stimulants ³ | 371 ^b | 325 ^b | 297 ^b | 312 ^b | 324 ^b | 240 ^a | 233 ^a | 196 | 191 | 165 | 177 |
| Methamphetamine ³ | 148 ^b | 103 ^b | 108 ^b | 100 ^b | 111 ^b | 77 ^a | 62 | 58 | 54 | 62 | 45 |
| Sedatives | 93 ^a | 64 | 75 | 67 | 54 | 68 | 67 | 56 | 70 | 52 | 53 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 1,655 ^b | 1,714 ^b | 1,732 ^b | 1,650 ^b | 1,605 ^b | 1,471 ^b | 1,477 ^b | 1,474 ^b | 1,493 ^b | 1,302 | 1,210 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.19B Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|------|
| ILLCIT DRUGS¹ | 21.9 ^b | 21.9 ^b | 21.5 ^b | 20.0 ^b | 19.7 ^a | 18.1 | 19.0 | 18.9 | 19.5 ^a | 18.8 | 18.0 |
| Marijuana and Hashish | 15.2 ^b | 14.6 ^b | 14.1 ^a | 12.8 | 12.9 | 11.4 ^a | 12.5 | 12.2 | 13.1 | 13.3 | 12.8 |
| Cocaine | 2.0 ^b | 1.8 ^b | 1.5 ^b | 1.7 ^b | 1.9 ^b | 1.7 ^b | 1.2 ^a | 0.8 | 1.0 | 0.9 | 0.8 |
| Crack | 0.5 ^b | 0.4 ^b | 0.3 ^a | 0.3 ^a | 0.3 ^b | 0.3 ^b | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Heroin | 0.3 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.1 |
| Hallucinogens | 3.7 ^b | 3.2 ^b | 3.0 ^b | 2.5 | 2.7 | 2.3 | 2.6 | 2.7 | 3.1 ^b | 2.3 | 2.2 |
| LSD | 1.1 ^b | 0.6 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 | 0.4 | 0.5 | 0.5 | 0.6 |
| PCP | 0.5 ^b | 0.4 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Ecstasy | 2.4 ^b | 1.5 | 1.4 | 1.1 | 1.4 | 1.3 | 1.5 | 1.7 | 2.2 ^b | 1.7 | 1.3 |
| Inhalants | 4.1 ^b | 4.6 ^b | 4.7 ^b | 4.9 ^b | 4.7 ^b | 4.0 ^b | 4.2 ^b | 4.2 ^b | 4.2 ^b | 3.5 | 3.1 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 9.9 ^b | 9.9 ^b | 10.2 ^b | 9.2 ^b | 9.1 ^b | 8.6 ^b | 8.6 ^b | 8.4 ^b | 8.5 ^b | 7.5 | 7.1 |
| Pain Relievers | 7.9 ^b | 8.1 ^b | 8.3 ^b | 7.4 ^b | 7.5 ^b | 7.0 ^b | 7.2 ^b | 7.2 ^b | 7.0 ^b | 6.5 ^a | 5.6 |
| OxyContin [®] | -- | -- | 0.8 | 0.7 | 0.8 | 1.0 | 0.8 | 1.0 | 1.0 | 1.0 | 0.7 |
| Tranquilizers | 2.6 | 2.7 | 2.5 | 2.2 | 2.3 | 2.4 | 2.1 | 2.1 | 2.2 | 2.1 | 2.2 |
| Stimulants ³ | 3.1 ^b | 2.7 ^b | 2.4 ^b | 2.5 ^b | 2.6 ^b | 1.9 ^a | 1.9 ^a | 1.6 | 1.6 | 1.3 | 1.5 |
| Methamphetamine ³ | 1.2 ^b | 0.8 ^b | 0.9 ^b | 0.8 ^b | 0.9 ^b | 0.6 ^a | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 |
| Sedatives | 0.8 ^b | 0.5 | 0.6 | 0.5 | 0.4 | 0.6 | 0.5 | 0.5 | 0.6 | 0.4 | 0.4 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 13.7 ^b | 14.0 ^b | 14.0 ^b | 13.3 ^b | 12.9 ^b | 11.9 ^b | 12.1 ^b | 12.2 ^b | 12.6 ^b | 10.7 | 9.9 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.20A Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|------------------|------------------|------------------|------------------|------------------|-----------------|--------------------|-------|
| ILLCIT DRUGS¹ | 1,560 ^b | 1,457 ^b | 1,360 ^a | 1,313 | 1,277 | 1,296 | 1,205 | 1,344 | 1,299 | 1,382 ^a | 1,217 |
| Marijuana and Hashish | 1,152 ^b | 1,093 ^a | 1,037 | 967 | 893 | 973 | 929 | 1,056 | 1,042 | 1,151 ^b | 951 |
| Cocaine | 77 ^b | 70 ^b | 75 ^b | 79 ^b | 49 ^b | 48 ^b | 62 ^b | 43 ^b | 37 ^a | 23 | 13 |
| Crack | 10 ^a | 22 ^b | 9 ^a | 14 ^a | 5 | 4 | 6 ^a | 5 | * | 2 | 1 |
| Heroin | 6 | 8 | 11 | 11 | 3 | 3 | 8 | 8 | 2 | 8 | * |
| Hallucinogens | 130 | 138 ^a | 120 | 98 | 86 | 103 | 148 ^b | 128 | 116 | 134 | 92 |
| LSD | 37 ^b | 27 ^a | 23 | 24 | 10 | 24 | 36 ^b | 21 | 30 ^a | 26 ^a | 10 |
| PCP | 11 | 14 | 7 | 8 | 6 | 9 | 10 | 8 | 4 | 3 | 4 |
| Ecstasy | 58 ^a | 37 | 45 | 32 | 38 | 34 | 58 ^a | 65 ^a | 50 | 57 ^a | 31 |
| Inhalants | 159 ^b | 139 ^a | 141 ^b | 160 ^b | 156 ^b | 158 ^b | 136 ^a | 120 | 115 | 98 | 86 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 460 ^b | 478 ^b | 413 ^b | 400 ^a | 403 ^a | 392 ^a | 310 | 351 | 282 | 302 | 308 |
| Pain Relievers | 376 ^b | 385 ^b | 337 ^b | 325 ^a | 331 ^a | 325 ^a | 248 | 302 | 246 | 244 | 250 |
| OxyContin [®] | -- | -- | 29 | 17 | 19 | 25 | 33 | 40 | 27 | 29 | 29 |
| Tranquilizers | 70 | 111 ^b | 76 | 84 | 46 | 84 | 63 | 58 | 61 | 74 | 52 |
| Stimulants ³ | 106 ^b | 107 ^b | 89 ^a | 79 | 81 ^a | 48 | 54 | 53 | 49 | 43 | 51 |
| Methamphetamine ³ | 40 ^a | 32 ^b | 24 | 39 ^b | 24 ^a | 10 | 16 | 18 | 8 | 13 | 9 |
| Sedatives | 6 | 27 ^a | 4 | 9 | 21 ^a | 15 | 8 | 25 ^a | 14 | 10 | 7 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 705 ^b | 682 ^b | 626 ^b | 616 ^b | 604 ^b | 584 ^b | 536 ^a | 539 ^a | 466 | 484 | 436 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.20B Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|------------------|------------------|------------------|------------------|-------------------|------------------|-------------------|------|
| ILLCIT DRUGS¹ | 12.3 ^b | 11.4 ^b | 10.6 ^a | 10.1 | 9.8 | 10.1 | 9.5 | 10.7 ^a | 10.4 | 10.8 ^a | 9.6 |
| Marijuana and Hashish | 9.1 ^b | 8.6 ^a | 8.1 | 7.5 | 6.9 | 7.5 | 7.3 | 8.4 ^a | 8.4 | 9.0 ^b | 7.5 |
| Cocaine | 0.6 ^b | 0.5 ^b | 0.6 ^b | 0.6 ^b | 0.4 ^b | 0.4 ^b | 0.5 ^b | 0.3 ^b | 0.3 ^a | 0.2 | 0.1 |
| Crack | 0.1 ^a | 0.2 ^b | 0.1 ^a | 0.1 ^a | 0.0 | 0.0 | 0.0 ^a | 0.0 | * | 0.0 | 0.0 |
| Heroin | 0.0 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | * |
| Hallucinogens | 1.0 ^a | 1.1 ^a | 0.9 | 0.8 | 0.7 | 0.8 | 1.2 ^b | 1.0 | 0.9 | 1.0 | 0.7 |
| LSD | 0.3 ^b | 0.2 ^a | 0.2 | 0.2 | 0.1 | 0.2 | 0.3 ^b | 0.2 | 0.2 ^a | 0.2 ^a | 0.1 |
| PCP | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 |
| Ecstasy | 0.5 ^a | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.5 ^a | 0.5 ^a | 0.4 | 0.4 ^a | 0.2 |
| Inhalants | 1.3 ^b | 1.1 ^a | 1.1 ^b | 1.2 ^b | 1.2 ^b | 1.2 ^b | 1.1 ^a | 1.0 | 0.9 | 0.8 | 0.7 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 3.6 ^b | 3.7 ^b | 3.2 ^b | 3.1 ^a | 3.1 ^a | 3.0 ^a | 2.4 | 2.8 | 2.3 | 2.4 | 2.4 |
| Pain Relievers | 3.0 ^b | 3.0 ^b | 2.6 ^a | 2.5 ^a | 2.5 ^a | 2.5 ^a | 1.9 | 2.4 | 2.0 | 1.9 | 2.0 |
| OxyContin [®] | -- | -- | 0.2 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 |
| Tranquilizers | 0.6 | 0.9 ^b | 0.6 | 0.7 | 0.4 | 0.6 | 0.5 | 0.5 | 0.5 | 0.6 | 0.4 |
| Stimulants ³ | 0.8 ^b | 0.8 ^b | 0.7 ^a | 0.6 | 0.6 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 | 0.4 |
| Methamphetamine ³ | 0.3 ^a | 0.3 ^b | 0.2 | 0.3 ^b | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Sedatives | 0.0 | 0.2 ^a | 0.0 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 ^a | 0.1 | 0.1 | 0.1 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 5.6 ^b | 5.3 ^b | 4.9 ^b | 4.8 ^b | 4.7 ^b | 4.5 ^b | 4.2 ^a | 4.3 ^a | 3.7 | 3.8 | 3.4 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.21A Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Numbers in Thousands, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|--------------------|--------------------|--------------------|------------------|------------------|------------------|------------------|------------------|------------------|-----------------|-------|
| ILLCIT DRUGS¹ | 1,318 ^a | 1,355 ^b | 1,314 ^a | 1,198 | 1,211 | 1,119 | 1,107 | 1,141 | 1,169 | 1,140 | 1,160 |
| Marijuana and Hashish | 871 | 878 | 872 | 763 | 802 | 721 ^a | 739 ^a | 757 | 760 | 816 | 849 |
| Cocaine | 68 ^b | 83 ^b | 51 ^b | 63 ^b | 61 ^b | 56 ^b | 36 | 27 | 24 | 44 ^a | 17 |
| Crack | 16 | 10 | 6 | 12 | 7 | 12 | 1 | 3 | 3 | 1 | * |
| Heroin | 5 | 10 | 4 | 4 | 13 | * | 5 | 5 | 6 | 6 | * |
| Hallucinogens | 126 ^b | 122 ^b | 91 | 107 ^a | 96 | 71 | 97 | 89 | 99 | 82 | 68 |
| LSD | 23 | 23 | 15 | 8 | 21 | 6 | 16 | 14 | 16 | 8 | 16 |
| PCP | 11 | 6 | 4 | 9 | 6 | * | 5 | 4 | 3 | 1 | * |
| Ecstasy | 66 | 54 | 32 | 41 | 45 | 31 | 47 | 50 | 73 | 52 | 43 |
| Inhalants | 143 | 181 ^b | 171 ^b | 153 ^a | 184 ^b | 136 | 134 | 124 | 147 ^a | 124 | 102 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 527 ^b | 519 ^b | 501 ^b | 439 | 443 | 438 | 401 | 424 | 440 | 386 | 385 |
| Pain Relievers | 418 ^b | 416 ^b | 414 ^b | 361 | 361 | 346 | 319 | 356 | 362 | 319 | 308 |
| OxyContin [®] | -- | -- | 37 | 17 | 10 | 22 | 17 | 29 | 20 | 21 | 20 |
| Tranquilizers | 117 | 115 | 86 | 76 | 79 | 97 | 88 | 86 | 73 | 88 | 87 |
| Stimulants ³ | 102 ^a | 115 ^b | 91 | 93 | 90 | 78 | 58 | 74 | 58 | 55 | 67 |
| Methamphetamine ³ | 31 | 38 | 34 | 32 | 29 | 24 | 13 | 18 | 16 | 23 | 22 |
| Sedatives | 41 ^a | 20 | 22 | 18 | 24 | 19 | 27 | 15 | 21 | 14 | 20 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 715 ^b | 735 ^b | 702 ^b | 637 ^b | 648 ^b | 598 ^a | 557 | 583 ^a | 626 ^b | 541 | 482 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

Table 7.21B Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Percentages, 2002-2012

| Drug | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|--|-------------------|-------------------|-------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------|
| ILLCIT DRUGS¹ | 10.9 ^b | 11.1 ^b | 10.6 ^a | 9.7 | 9.8 | 9.1 | 9.1 | 9.5 | 9.8 | 9.3 | 9.5 |
| Marijuana and Hashish | 7.2 | 7.2 | 7.1 | 6.2 | 6.5 | 5.8 ^a | 6.1 ^a | 6.3 | 6.4 | 6.7 | 7.0 |
| Cocaine | 0.6 ^b | 0.7 ^b | 0.4 ^b | 0.5 ^b | 0.5 ^b | 0.5 ^b | 0.3 | 0.2 | 0.2 | 0.4 ^a | 0.1 |
| Crack | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | * |
| Heroin | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | * | 0.0 | 0.0 | 0.1 | 0.1 | * |
| Hallucinogens | 1.0 ^b | 1.0 ^b | 0.7 | 0.9 ^a | 0.8 | 0.6 | 0.8 | 0.7 | 0.8 | 0.7 | 0.6 |
| LSD | 0.2 | 0.2 | 0.1 | 0.1 | 0.2 | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| PCP | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | * | 0.0 | 0.0 | 0.0 | 0.0 | * |
| Ecstasy | 0.5 | 0.4 | 0.3 | 0.3 | 0.4 | 0.3 | 0.4 | 0.4 | 0.6 ^a | 0.4 | 0.3 |
| Inhalants | 1.2 ^a | 1.5 ^b | 1.4 ^b | 1.2 ^a | 1.5 ^b | 1.1 | 1.1 | 1.0 | 1.2 ^a | 1.0 | 0.8 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 4.4 ^b | 4.2 ^b | 4.1 ^b | 3.5 | 3.6 | 3.5 | 3.3 | 3.5 | 3.7 | 3.2 | 3.2 |
| Pain Relievers | 3.5 ^b | 3.4 ^b | 3.4 ^b | 2.9 | 2.9 | 2.8 | 2.6 | 3.0 | 3.0 | 2.6 | 2.5 |
| OxyContin [®] | -- | -- | 0.3 | 0.1 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Tranquilizers | 1.0 | 0.9 | 0.7 | 0.6 | 0.6 | 0.8 | 0.7 | 0.7 | 0.6 | 0.7 | 0.7 |
| Stimulants ³ | 0.8 ^a | 0.9 ^b | 0.7 | 0.8 | 0.7 | 0.6 | 0.5 | 0.6 | 0.5 | 0.5 | 0.5 |
| Methamphetamine ³ | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 |
| Sedatives | 0.3 ^a | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 5.9 ^b | 6.0 ^b | 5.7 ^b | 5.1 ^b | 5.2 ^b | 4.8 ^a | 4.6 | 4.8 ^a | 5.3 ^b | 4.4 | 4.0 |

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.