

**Table 7.22A GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	1,635	1,326	1,198	1,443	1,698	1,389	1,446
<b>AGE GROUP</b>							
12 to 17	40	21	21	30	21	17	20
18 to 25	475 <sup>b</sup>	307 <sup>b</sup>	202	255 <sup>b</sup>	217	142	150
26 or Older	1,120	998	976	1,158	1,459	1,230	1,276
<b>GENDER</b>							
Male	986	864	765	965	1,211	976	964
Female	650	462	433	479	486	413	482
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	130	68	100	99	170	108	143
<b>AGE GROUP</b>							
12 to 17	16	7	8	14	3	11	15
18 to 25	46	42	22	31	30	23	33
26 or Older	68	19	70	55	137	74	96
<b>GENDER</b>							
Male	105	55	81	78	123	96	101
Female	26	14	19	21	47	12	42
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	28	24	28	18	70	43	66
<b>AGE GROUP</b>							
12 to 17	5	3	1	5	*	1	11
18 to 25	6	21	5	4	6	3	12
26 or Older	17	*	22	*	63	39	43
<b>GENDER</b>							
Male	24	22	22	8	40	43	32
Female	*	*	*	9	30	*	34

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.22B GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.7	0.5	0.5	0.6	0.7	0.5	0.6
<b>AGE GROUP</b>							
12 to 17	0.2	0.1	0.1	0.1	0.1	0.1	0.1
18 to 25	1.5 <sup>b</sup>	0.9 <sup>b</sup>	0.6 <sup>a</sup>	0.8 <sup>b</sup>	0.6 <sup>a</sup>	0.4	0.4
26 or Older	0.6	0.5	0.5	0.6	0.7	0.6	0.6
<b>GENDER</b>							
Male	0.8	0.7	0.6	0.8	1.0	0.8	0.8
Female	0.5 <sup>a</sup>	0.4	0.3	0.4	0.4	0.3	0.4
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.1	0.0	0.0	0.0	0.1	0.0	0.1
<b>AGE GROUP</b>							
12 to 17	0.1	0.0	0.0	0.1	0.0	0.0	0.1
18 to 25	0.1	0.1	0.1	0.1	0.1	0.1	0.1
26 or Older	0.0	0.0	0.0	0.0	0.1	0.0	0.0
<b>GENDER</b>							
Male	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>AGE GROUP</b>							
12 to 17	0.0	0.0	0.0	0.0	*	0.0	0.0
18 to 25	0.0	0.1	0.0	0.0	0.0	0.0	0.0
26 or Older	0.0	*	0.0	*	0.0	0.0	0.0
<b>GENDER</b>							
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	*	*	0.0	0.0	*	0.0

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.23A Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	4,704 <sup>b</sup>	4,951 <sup>b</sup>	5,366 <sup>b</sup>	6,751 <sup>b</sup>	7,936 <sup>b</sup>	8,198 <sup>b</sup>	9,624
<b>AGE GROUP</b>							
12 to 17	701	673	613	631	664	621	665
18 to 25	2,778 <sup>b</sup>	2,868 <sup>b</sup>	3,111 <sup>b</sup>	3,701 <sup>b</sup>	4,038 <sup>a</sup>	4,079 <sup>a</sup>	4,392
26 or Older	1,224 <sup>b</sup>	1,410 <sup>b</sup>	1,642 <sup>b</sup>	2,419 <sup>b</sup>	3,234 <sup>b</sup>	3,498 <sup>b</sup>	4,567
<b>GENDER</b>							
Male	2,662 <sup>b</sup>	2,798 <sup>b</sup>	2,959 <sup>b</sup>	4,012 <sup>b</sup>	4,707 <sup>a</sup>	4,327 <sup>b</sup>	5,382
Female	2,042 <sup>b</sup>	2,153 <sup>b</sup>	2,407 <sup>b</sup>	2,739 <sup>b</sup>	3,228 <sup>b</sup>	3,870	4,241
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	1,991 <sup>b</sup>	2,003 <sup>b</sup>	2,099 <sup>b</sup>	2,779 <sup>b</sup>	2,904 <sup>b</sup>	3,076 <sup>a</sup>	3,509
<b>AGE GROUP</b>							
12 to 17	434	395	382	398	414	409	391
18 to 25	1,212 <sup>b</sup>	1,216 <sup>b</sup>	1,218 <sup>b</sup>	1,608 <sup>b</sup>	1,774	1,766	1,961
26 or Older	345 <sup>b</sup>	391 <sup>b</sup>	499 <sup>b</sup>	773 <sup>b</sup>	716 <sup>b</sup>	902	1,157
<b>GENDER</b>							
Male	1,055 <sup>b</sup>	1,157 <sup>b</sup>	1,155 <sup>b</sup>	1,649	1,661	1,684	1,934
Female	937 <sup>b</sup>	846 <sup>b</sup>	944 <sup>b</sup>	1,129 <sup>b</sup>	1,243 <sup>b</sup>	1,392	1,575
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	551 <sup>b</sup>	529 <sup>b</sup>	613 <sup>b</sup>	858	921	836	948
<b>AGE GROUP</b>							
12 to 17	119	134	102	118	126	122	103
18 to 25	339 <sup>b</sup>	344 <sup>b</sup>	346 <sup>b</sup>	466	501	490	548
26 or Older	94 <sup>b</sup>	51 <sup>b</sup>	165	274	293	224	297
<b>GENDER</b>							
Male	299 <sup>b</sup>	300 <sup>b</sup>	344 <sup>a</sup>	512	547	502	507
Female	252 <sup>b</sup>	229 <sup>b</sup>	269 <sup>b</sup>	346	373	334	441

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.23B Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	1.9 <sup>b</sup>	2.0 <sup>b</sup>	2.1 <sup>b</sup>	2.7 <sup>b</sup>	3.1 <sup>b</sup>	3.2 <sup>b</sup>	3.7
<b>AGE GROUP</b>							
12 to 17	2.8	2.7	2.5	2.6	2.7	2.5	2.7
18 to 25	8.5 <sup>b</sup>	8.8 <sup>b</sup>	9.4 <sup>b</sup>	11.0 <sup>b</sup>	11.9	11.9	12.7
26 or Older	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.9 <sup>b</sup>	1.2 <sup>b</sup>	1.7 <sup>b</sup>	1.8 <sup>b</sup>	2.3
<b>GENDER</b>							
Male	2.2 <sup>b</sup>	2.3 <sup>b</sup>	2.4 <sup>b</sup>	3.3 <sup>b</sup>	3.8	3.5 <sup>b</sup>	4.3
Female	1.6 <sup>b</sup>	1.7 <sup>b</sup>	1.9 <sup>b</sup>	2.1 <sup>b</sup>	2.5 <sup>b</sup>	2.9	3.2
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	1.1 <sup>b</sup>	1.1 <sup>b</sup>	1.2 <sup>a</sup>	1.3
<b>AGE GROUP</b>							
12 to 17	1.7	1.6	1.5	1.6	1.7	1.6	1.6
18 to 25	3.7 <sup>b</sup>	3.7 <sup>b</sup>	3.7 <sup>b</sup>	4.8 <sup>a</sup>	5.2	5.1	5.7
26 or Older	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.4 <sup>a</sup>	0.4 <sup>b</sup>	0.5	0.6
<b>GENDER</b>							
Male	0.9 <sup>b</sup>	1.0 <sup>b</sup>	1.0 <sup>b</sup>	1.3	1.3	1.4	1.5
Female	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.9 <sup>b</sup>	1.0 <sup>a</sup>	1.0	1.2
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3	0.4	0.3	0.4
<b>AGE GROUP</b>							
12 to 17	0.5	0.5	0.4	0.5	0.5	0.5	0.4
18 to 25	1.0 <sup>b</sup>	1.1 <sup>b</sup>	1.1 <sup>b</sup>	1.4	1.5	1.4	1.6
26 or Older	0.0 <sup>b</sup>	0.0 <sup>b</sup>	0.1	0.1	0.2	0.1	0.1
<b>GENDER</b>							
Male	0.3 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>a</sup>	0.4	0.4	0.4	0.4
Female	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3	0.3	0.3	0.3

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.24A Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	4,440 <sup>b</sup>	5,197 <sup>b</sup>	5,525 <sup>b</sup>	6,231	6,444	6,123	6,537
<b>AGE GROUP</b>							
12 to 17	315	297	268	287	231	273	273
18 to 25	1,187	1,087	1,182	1,284	1,288	1,113	1,204
26 or Older	2,939 <sup>b</sup>	3,813 <sup>b</sup>	4,075 <sup>b</sup>	4,660	4,925	4,738	5,060
<b>GENDER</b>							
Male	2,141 <sup>b</sup>	2,366 <sup>b</sup>	2,345 <sup>b</sup>	2,932	3,227	2,703	3,029
Female	2,299 <sup>b</sup>	2,831 <sup>a</sup>	3,180	3,299	3,217	3,421	3,508
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	2,034	2,048	2,065	2,163	2,264	1,904	1,844
<b>AGE GROUP</b>							
12 to 17	174 <sup>a</sup>	152	139	142	123	150	121
18 to 25	536 <sup>a</sup>	476	480	530 <sup>a</sup>	453	382	414
26 or Older	1,324	1,420	1,446	1,492	1,688	1,371	1,309
<b>GENDER</b>							
Male	997	995	940	1,089	1,160	789	814
Female	1,037	1,053	1,125	1,074	1,104	1,115	1,030
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	544	377	442	613	815	486	527
<b>AGE GROUP</b>							
12 to 17	46	30	28	35	35	35	30
18 to 25	98	108	133	132	109	92	92
26 or Older	400	238	281	446	671	359	405
<b>GENDER</b>							
Male	329	135	206	308	460 <sup>a</sup>	205	191
Female	215	241	236	305	355	281	336

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.24B Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	1.8 <sup>b</sup>	2.1 <sup>b</sup>	2.2 <sup>a</sup>	2.5	2.5	2.4	2.5
<b>AGE GROUP</b>							
12 to 17	1.2	1.2	1.1	1.2	0.9	1.1	1.1
18 to 25	3.6	3.3	3.6	3.8	3.8	3.2	3.5
26 or Older	1.6 <sup>b</sup>	2.0 <sup>b</sup>	2.1 <sup>a</sup>	2.4	2.5	2.4	2.5
<b>GENDER</b>							
Male	1.8 <sup>b</sup>	2.0 <sup>a</sup>	1.9 <sup>a</sup>	2.4	2.6	2.2	2.4
Female	1.8 <sup>b</sup>	2.2 <sup>a</sup>	2.5	2.5	2.5	2.6	2.6
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.8	0.8	0.8	0.9	0.9	0.7	0.7
<b>AGE GROUP</b>							
12 to 17	0.7 <sup>a</sup>	0.6	0.6	0.6	0.5	0.6	0.5
18 to 25	1.6 <sup>b</sup>	1.5	1.5	1.6 <sup>a</sup>	1.3	1.1	1.2
26 or Older	0.7	0.7	0.8	0.8	0.9	0.7	0.7
<b>GENDER</b>							
Male	0.8	0.8	0.8	0.9 <sup>a</sup>	0.9 <sup>a</sup>	0.6	0.6
Female	0.8	0.8	0.9	0.8	0.8	0.8	0.8
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.2	0.2	0.2	0.2	0.3	0.2	0.2
<b>AGE GROUP</b>							
12 to 17	0.2	0.1	0.1	0.1	0.1	0.1	0.1
18 to 25	0.3	0.3	0.4	0.4	0.3	0.3	0.3
26 or Older	0.2	0.1	0.1	0.2	0.3	0.2	0.2
<b>GENDER</b>							
Male	0.3	0.1	0.2	0.3	0.4 <sup>a</sup>	0.2	0.2
Female	0.2	0.2	0.2	0.2	0.3	0.2	0.3

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.25A Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	7,770	7,108 <sup>b</sup>	7,532 <sup>a</sup>	7,419 <sup>a</sup>	8,044	7,678	8,311
<b>AGE GROUP</b>							
12 to 17	932 <sup>b</sup>	855 <sup>b</sup>	795	750	760	704	699
18 to 25	2,134 <sup>a</sup>	2,040 <sup>b</sup>	2,183	2,325	2,324	2,270	2,373
26 or Older	4,705	4,213 <sup>b</sup>	4,554	4,344 <sup>a</sup>	4,959	4,704	5,239
<b>GENDER</b>							
Male	4,879	4,211 <sup>b</sup>	4,428 <sup>a</sup>	4,262 <sup>b</sup>	5,030	4,696	5,150
Female	2,891	2,898	3,103	3,157	3,014	2,982	3,162
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	1,582	1,354	1,434	1,470	1,781 <sup>a</sup>	1,577	1,308
<b>AGE GROUP</b>							
12 to 17	486 <sup>b</sup>	433 <sup>b</sup>	404 <sup>a</sup>	373	374	332	313
18 to 25	511	482	462	485	489	519	487
26 or Older	586	439	568	611	918 <sup>a</sup>	726	508
<b>GENDER</b>							
Male	887 <sup>b</sup>	690	779	727	1,102 <sup>b</sup>	807	631
Female	696	664	655	743	679	771	678
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	423	403	404	412	467	395	410
<b>AGE GROUP</b>							
12 to 17	123	125	104	108	94	77	87
18 to 25	107	130	107	124	127	94	117
26 or Older	193	147	192	181	246	224	205
<b>GENDER</b>							
Male	263	205	220	195	248	189	157
Female	160	197	184	217	219	206	252

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.25B Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	3.2	2.9 <sup>a</sup>	3.0	2.9	3.2	3.0	3.2
<b>AGE GROUP</b>							
12 to 17	3.7 <sup>b</sup>	3.4 <sup>a</sup>	3.2	3.0	3.1	2.8	2.8
18 to 25	6.5	6.2	6.6	6.9	6.8	6.6	6.9
26 or Older	2.5	2.2 <sup>a</sup>	2.4	2.2 <sup>a</sup>	2.5	2.4	2.6
<b>GENDER</b>							
Male	4.1	3.5 <sup>a</sup>	3.7	3.5 <sup>a</sup>	4.1	3.8	4.1
Female	2.3	2.3	2.4	2.4	2.3	2.2	2.4
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.6 <sup>a</sup>	0.5	0.6	0.6	0.7 <sup>b</sup>	0.6	0.5
<b>AGE GROUP</b>							
12 to 17	1.9 <sup>b</sup>	1.7 <sup>b</sup>	1.6 <sup>a</sup>	1.5	1.5	1.3	1.3
18 to 25	1.6	1.5	1.4	1.4	1.4	1.5	1.4
26 or Older	0.3	0.2	0.3	0.3	0.5 <sup>a</sup>	0.4	0.3
<b>GENDER</b>							
Male	0.7 <sup>b</sup>	0.6	0.6	0.6	0.9 <sup>b</sup>	0.6	0.5
Female	0.5	0.5	0.5	0.6	0.5	0.6	0.5
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.2	0.2	0.2	0.2	0.2	0.2	0.2
<b>AGE GROUP</b>							
12 to 17	0.5	0.5	0.4	0.4	0.4	0.3	0.4
18 to 25	0.3	0.4	0.3	0.4	0.4	0.3	0.3
26 or Older	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>GENDER</b>							
Male	0.2 <sup>a</sup>	0.2	0.2	0.2	0.2	0.2	0.1
Female	0.1	0.2	0.1	0.2	0.2	0.2	0.2

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.26A Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	2,283	2,191 <sup>a</sup>	2,148 <sup>a</sup>	2,309	2,411	2,379	2,669
<b>AGE GROUP</b>							
12 to 17	83	71	74	78	80	88	67
18 to 25	919 <sup>b</sup>	796 <sup>b</sup>	636 <sup>a</sup>	595	555	503	496
26 or Older	1,281 <sup>b</sup>	1,324 <sup>b</sup>	1,439 <sup>b</sup>	1,637 <sup>a</sup>	1,776	1,788	2,106
<b>GENDER</b>							
Male	1,468 <sup>a</sup>	1,436 <sup>b</sup>	1,354 <sup>b</sup>	1,547	1,747	1,552 <sup>a</sup>	1,911
Female	814	755	794	762	663	827	758
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	203	212	150 <sup>a</sup>	202	246	336	280
<b>AGE GROUP</b>							
12 to 17	33	32	34	35	51	54	36
18 to 25	74	122	70	92	136	152	98
26 or Older	95	58	46 <sup>a</sup>	75	59	130	146
<b>GENDER</b>							
Male	141	143	105 <sup>a</sup>	115 <sup>a</sup>	175	224	206
Female	62	69	45	88	71	112	74
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	53	39	27	86	61	82	53
<b>AGE GROUP</b>							
12 to 17	11	12	10	17	14	25	8
18 to 25	14	17	17	26	22	29	22
26 or Older	28	10	*	43	24	28	23
<b>GENDER</b>							
Male	34	27	16	42	39	38	20
Female	19	12	10	43	21	44	33

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.26B Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.9	0.9	0.9 <sup>a</sup>	0.9	1.0	0.9	1.0
<b>AGE GROUP</b>							
12 to 17	0.3	0.3	0.3	0.3	0.3	0.4	0.3
18 to 25	2.8 <sup>b</sup>	2.4 <sup>b</sup>	1.9 <sup>b</sup>	1.8 <sup>a</sup>	1.6	1.5	1.4
26 or Older	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.8	0.9	0.9	1.1
<b>GENDER</b>							
Male	1.2	1.2 <sup>a</sup>	1.1 <sup>b</sup>	1.3	1.4	1.2	1.5
Female	0.6	0.6	0.6	0.6	0.5	0.6	0.6
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.1	0.1	0.1 <sup>a</sup>	0.1	0.1	0.1	0.1
<b>AGE GROUP</b>							
12 to 17	0.1	0.1	0.1	0.1	0.2	0.2	0.1
18 to 25	0.2	0.4	0.2	0.3	0.4	0.4	0.3
26 or Older	0.1	0.0	0.0 <sup>a</sup>	0.0	0.0	0.1	0.1
<b>GENDER</b>							
Male	0.1	0.1	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.2	0.2
Female	0.0	0.1	0.0	0.1	0.1	0.1	0.1
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>AGE GROUP</b>							
12 to 17	0.0	0.0	0.0	0.1	0.1	0.1	0.0
18 to 25	0.0	0.1	0.1	0.1	0.1	0.1	0.1
26 or Older	0.0	0.0	*	0.0	0.0	0.0	0.0
<b>GENDER</b>							
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.27A DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2012**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012
<b>LIFETIME USE</b>							
<b>TOTAL</b>	688 <sup>b</sup>	678 <sup>b</sup>	731 <sup>b</sup>	885 <sup>b</sup>	1,212	1,082 <sup>b</sup>	1,475
<b>AGE GROUP</b>							
12 to 17	45 <sup>b</sup>	44 <sup>b</sup>	49 <sup>b</sup>	58 <sup>a</sup>	61 <sup>a</sup>	65 <sup>a</sup>	98
18 to 25	307 <sup>b</sup>	270 <sup>b</sup>	280 <sup>b</sup>	367 <sup>b</sup>	444 <sup>b</sup>	483 <sup>a</sup>	619
26 or Older	336 <sup>b</sup>	363 <sup>b</sup>	402 <sup>b</sup>	460 <sup>a</sup>	707	535	757
<b>GENDER</b>							
Male	508 <sup>b</sup>	521 <sup>b</sup>	474 <sup>b</sup>	644 <sup>b</sup>	883	825 <sup>a</sup>	1,104
Female	180 <sup>b</sup>	157 <sup>b</sup>	258	241 <sup>a</sup>	329	257	371
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	104 <sup>b</sup>	82 <sup>b</sup>	150 <sup>b</sup>	194 <sup>b</sup>	234	265	325
<b>AGE GROUP</b>							
12 to 17	16 <sup>b</sup>	20 <sup>b</sup>	29	38	38	36	52
18 to 25	64 <sup>b</sup>	52 <sup>b</sup>	73 <sup>b</sup>	111 <sup>b</sup>	118 <sup>b</sup>	183	198
26 or Older	24	10 <sup>a</sup>	47	46	79	47	75
<b>GENDER</b>							
Male	83 <sup>b</sup>	63 <sup>b</sup>	111 <sup>b</sup>	117 <sup>b</sup>	161 <sup>a</sup>	213	255
Female	21 <sup>b</sup>	19 <sup>b</sup>	39	77	73	53	70
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	20 <sup>b</sup>	*	46	57	81	47	75
<b>AGE GROUP</b>							
12 to 17	*	*	8	11	17	7	17
18 to 25	15 <sup>a</sup>	10 <sup>b</sup>	19	12 <sup>b</sup>	21	25	46
26 or Older	*	*	19	34	43	14	12
<b>GENDER</b>							
Male	13 <sup>a</sup>	7 <sup>b</sup>	40	22	50	27	47
Female	7 <sup>a</sup>	*	*	35	31	20	29

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.27B DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.5	0.4 <sup>b</sup>	0.6
<b>AGE GROUP</b>							
12 to 17	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.4
18 to 25	0.9 <sup>b</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	1.1 <sup>b</sup>	1.3 <sup>b</sup>	1.4 <sup>a</sup>	1.8
26 or Older	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.4	0.3	0.4
<b>GENDER</b>							
Male	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.5 <sup>b</sup>	0.7	0.7 <sup>a</sup>	0.9
Female	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.2	0.2	0.3	0.2	0.3
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.0 <sup>b</sup>	0.0 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1	0.1	0.1
<b>AGE GROUP</b>							
12 to 17	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1	0.2	0.2	0.1	0.2
18 to 25	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.5	0.6
26 or Older	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0
<b>GENDER</b>							
Male	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>a</sup>	0.2	0.2
Female	0.0 <sup>b</sup>	0.0 <sup>b</sup>	0.0	0.1	0.1	0.0	0.1
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.0 <sup>b</sup>	*	0.0	0.0	0.0	0.0	0.0
<b>AGE GROUP</b>							
12 to 17	*	*	0.0	0.0	0.1	0.0	0.1
18 to 25	0.0 <sup>a</sup>	0.0 <sup>b</sup>	0.1	0.0 <sup>b</sup>	0.1	0.1	0.1
26 or Older	*	*	0.0	0.0	0.0	0.0	0.0
<b>GENDER</b>							
Male	0.0 <sup>a</sup>	0.0 <sup>b</sup>	0.0	0.0	0.0	0.0	0.0
Female	0.0 <sup>a</sup>	*	*	0.0	0.0	0.0	0.0

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.28A Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	1,802 <sup>b</sup>	2,522 <sup>b</sup>	3,201 <sup>b</sup>	4,406 <sup>b</sup>	5,235	5,285	5,223
<b>AGE GROUP</b>							
12 to 17	229	298	394 <sup>b</sup>	418 <sup>b</sup>	399 <sup>b</sup>	375 <sup>b</sup>	267
18 to 25	1,188 <sup>b</sup>	1,599 <sup>b</sup>	1,962 <sup>b</sup>	2,698	2,999	3,015	2,788
26 or Older	386 <sup>b</sup>	624 <sup>b</sup>	846 <sup>b</sup>	1,289 <sup>b</sup>	1,837	1,895	2,168
<b>GENDER</b>							
Male	1,436 <sup>b</sup>	1,873 <sup>b</sup>	2,296 <sup>b</sup>	3,391 <sup>a</sup>	3,907	3,733	3,891
Female	366 <sup>b</sup>	649 <sup>b</sup>	905 <sup>b</sup>	1,015 <sup>b</sup>	1,329	1,551	1,332
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	761	1,007 <sup>b</sup>	1,213 <sup>b</sup>	1,283 <sup>b</sup>	1,191 <sup>b</sup>	871 <sup>b</sup>	648
<b>AGE GROUP</b>							
12 to 17	149	192 <sup>b</sup>	264 <sup>b</sup>	244 <sup>b</sup>	229 <sup>b</sup>	208 <sup>b</sup>	129
18 to 25	552 <sup>b</sup>	656 <sup>b</sup>	738 <sup>b</sup>	784 <sup>b</sup>	651 <sup>b</sup>	522 <sup>b</sup>	378
26 or Older	60 <sup>a</sup>	158	211	255 <sup>a</sup>	311 <sup>a</sup>	141	140
<b>GENDER</b>							
Male	600 <sup>a</sup>	737 <sup>b</sup>	877 <sup>b</sup>	965 <sup>b</sup>	929 <sup>b</sup>	629 <sup>b</sup>	463
Female	161	270 <sup>a</sup>	337 <sup>b</sup>	318 <sup>b</sup>	262 <sup>a</sup>	242	184
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	136	200 <sup>a</sup>	198 <sup>a</sup>	194	162	136	122
<b>AGE GROUP</b>							
12 to 17	28	42	59 <sup>b</sup>	50 <sup>a</sup>	44	37	24
18 to 25	104 <sup>b</sup>	134 <sup>b</sup>	129 <sup>b</sup>	95 <sup>a</sup>	93 <sup>a</sup>	85	48
26 or Older	*	24	10	49	25	13	50
<b>GENDER</b>							
Male	112	151	160 <sup>a</sup>	134	134	95	88
Female	24	49	37	60	28	41	33

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.28B Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.7 <sup>b</sup>	1.0 <sup>b</sup>	1.3 <sup>b</sup>	1.7 <sup>b</sup>	2.1	2.1	2.0
<b>AGE GROUP</b>							
12 to 17	0.9	1.2	1.6 <sup>b</sup>	1.7 <sup>b</sup>	1.6 <sup>b</sup>	1.5 <sup>b</sup>	1.1
18 to 25	3.6 <sup>b</sup>	4.9 <sup>b</sup>	6.0 <sup>b</sup>	8.0	8.8	8.8	8.1
26 or Older	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.7 <sup>b</sup>	0.9	1.0	1.1
<b>GENDER</b>							
Male	1.2 <sup>b</sup>	1.6 <sup>b</sup>	1.9 <sup>b</sup>	2.8	3.2	3.0	3.1
Female	0.3 <sup>b</sup>	0.5 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	1.0	1.2	1.0
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.3 <sup>a</sup>	0.4 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.3 <sup>b</sup>	0.2
<b>AGE GROUP</b>							
12 to 17	0.6	0.8 <sup>a</sup>	1.1 <sup>b</sup>	1.0 <sup>b</sup>	0.9 <sup>b</sup>	0.8 <sup>b</sup>	0.5
18 to 25	1.7 <sup>b</sup>	2.0 <sup>b</sup>	2.2 <sup>b</sup>	2.3 <sup>b</sup>	1.9 <sup>b</sup>	1.5 <sup>b</sup>	1.1
26 or Older	0.0	0.1	0.1	0.1 <sup>a</sup>	0.2 <sup>a</sup>	0.1	0.1
<b>GENDER</b>							
Male	0.5 <sup>b</sup>	0.6 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.5 <sup>b</sup>	0.4
Female	0.1	0.2 <sup>a</sup>	0.3 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.2	0.1
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.1	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.0
<b>AGE GROUP</b>							
12 to 17	0.1	0.2	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.2	0.1	0.1
18 to 25	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>a</sup>	0.2	0.1
26 or Older	*	0.0	0.0	0.0	0.0	0.0	0.0
<b>GENDER</b>							
Male	0.1	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.1	0.1
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.