

**Table 7.22C GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	118	106	106	119	155	121	121
<b>AGE GROUP</b>							
12 to 17	8	7	5	8	6	5	7
18 to 25	37	30	22	25	27	20	22
26 or Older	111	100	100	117	150	120	117
<b>GENDER</b>							
Male	88	87	87	98	130	101	106
Female	70	59	55	63	68	63	66
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	31	13	31	26	47	37	49
<b>AGE GROUP</b>							
12 to 17	5	3	3	5	2	4	7
18 to 25	10	10	8	10	9	8	10
26 or Older	28	11	30	24	46	35	45
<b>GENDER</b>							
Male	29	12	30	24	43	36	41
Female	7	4	9	9	19	6	28
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	18	9	18	9	29	28	34
<b>AGE GROUP</b>							
12 to 17	3	3	1	3	*	1	6
18 to 25	4	8	3	3	3	3	7
26 or Older	17	*	17	*	29	28	29
<b>GENDER</b>							
Male	18	9	17	4	23	28	20
Female	*	*	*	8	18	*	28

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.22D GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.05	0.04	0.04	0.05	0.06	0.05	0.05
<b>AGE GROUP</b>							
12 to 17	0.03	0.03	0.02	0.03	0.02	0.02	0.03
18 to 25	0.11	0.09	0.07	0.08	0.08	0.06	0.06
26 or Older	0.06	0.05	0.05	0.06	0.08	0.06	0.06
<b>GENDER</b>							
Male	0.07	0.07	0.07	0.08	0.11	0.08	0.08
Female	0.06	0.05	0.04	0.05	0.05	0.05	0.05
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.01	0.01	0.01	0.01	0.02	0.01	0.02
<b>AGE GROUP</b>							
12 to 17	0.02	0.01	0.01	0.02	0.01	0.02	0.03
18 to 25	0.03	0.03	0.02	0.03	0.03	0.02	0.03
26 or Older	0.01	0.01	0.02	0.01	0.02	0.02	0.02
<b>GENDER</b>							
Male	0.02	0.01	0.02	0.02	0.03	0.03	0.03
Female	0.01	0.00	0.01	0.01	0.01	0.00	0.02
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.01	0.00	0.01	0.00	0.01	0.01	0.01
<b>AGE GROUP</b>							
12 to 17	0.01	0.01	0.00	0.01	*	0.00	0.02
18 to 25	0.01	0.03	0.01	0.01	0.01	0.01	0.02
26 or Older	0.01	*	0.01	*	0.01	0.01	0.01
<b>GENDER</b>							
Male	0.01	0.01	0.01	0.00	0.02	0.02	0.02
Female	*	*	*	0.01	0.01	*	0.02

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.22P GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.0980	0.7450	0.2286	0.7884	0.1429	0.8000	
<b>AGE GROUP</b>							
12 to 17	0.0726	0.8926	0.9347	0.3056	0.8302	0.7692	
18 to 25	0.0000	0.0000	0.0496	0.0010	0.0426	0.8267	
26 or Older	0.6246	0.1686	0.1026	0.6379	0.2541	0.8454	
<b>GENDER</b>							
Male	0.5941	0.6746	0.2117	0.8388	0.1168	0.8800	
Female	0.0334	0.9610	0.7524	0.8700	0.8364	0.4772	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.9264	0.1599	0.5028	0.4649	0.6789	0.5798	
<b>AGE GROUP</b>							
12 to 17	0.8769	0.2777	0.3648	0.9756	0.1124	0.6547	
18 to 25	0.3094	0.4247	0.4719	0.9251	0.8561	0.4488	
26 or Older	0.6764	0.1088	0.6762	0.4469	0.5353	0.7208	
<b>GENDER</b>							
Male	0.8500	0.3035	0.7414	0.6672	0.7095	0.9393	
Female	0.5973	0.3251	0.4392	0.4848	0.8658	0.2957	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.3564	0.2511	0.3457	0.1767	0.9007	0.6151	
<b>AGE GROUP</b>							
12 to 17	0.3681	0.2633	0.1219	0.4302	*	0.1285	
18 to 25	0.4599	0.4097	0.3321	0.3022	0.4038	0.2201	
26 or Older	0.4842	*	0.5741	*	0.5934	0.9367	
<b>GENDER</b>							
Male	0.8224	0.6914	0.7276	0.2582	0.7775	0.7442	
Female	*	*	*	0.3954	0.9240	*	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.23C Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	165	170	180	220	257	232	286
<b>AGE GROUP</b>							
12 to 17	34	37	31	32	37	34	37
18 to 25	98	97	95	112	117	113	122
26 or Older	112	116	139	164	204	194	233
<b>GENDER</b>							
Male	123	125	127	174	197	161	214
Female	96	102	118	113	139	158	172
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	92	99	102	135	128	134	152
<b>AGE GROUP</b>							
12 to 17	27	28	25	25	29	27	26
18 to 25	66	69	65	82	84	86	88
26 or Older	53	58	71	90	76	91	112
<b>GENDER</b>							
Male	62	72	74	107	97	93	106
Female	60	61	65	70	67	92	96
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	43	45	55	65	73	65	70
<b>AGE GROUP</b>							
12 to 17	14	17	13	13	16	12	14
18 to 25	32	37	31	36	46	41	43
26 or Older	24	15	42	53	49	47	55
<b>GENDER</b>							
Male	33	32	41	57	57	45	53
Female	27	30	33	35	37	42	45

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.23D Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.07	0.07	0.07	0.09	0.10	0.09	0.11
<b>AGE GROUP</b>							
12 to 17	0.14	0.15	0.12	0.13	0.15	0.13	0.15
18 to 25	0.30	0.30	0.29	0.33	0.34	0.33	0.35
26 or Older	0.06	0.06	0.07	0.08	0.10	0.10	0.12
<b>GENDER</b>							
Male	0.10	0.10	0.10	0.14	0.16	0.13	0.17
Female	0.08	0.08	0.09	0.09	0.11	0.12	0.13
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.04	0.04	0.04	0.05	0.05	0.05	0.06
<b>AGE GROUP</b>							
12 to 17	0.11	0.11	0.10	0.10	0.12	0.11	0.10
18 to 25	0.20	0.21	0.20	0.24	0.25	0.25	0.25
26 or Older	0.03	0.03	0.04	0.05	0.04	0.05	0.06
<b>GENDER</b>							
Male	0.05	0.06	0.06	0.09	0.08	0.07	0.08
Female	0.05	0.05	0.05	0.05	0.05	0.07	0.07
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.02	0.02	0.02	0.03	0.03	0.03	0.03
<b>AGE GROUP</b>							
12 to 17	0.05	0.07	0.05	0.05	0.06	0.05	0.06
18 to 25	0.10	0.11	0.09	0.11	0.13	0.12	0.12
26 or Older	0.01	0.01	0.02	0.03	0.03	0.02	0.03
<b>GENDER</b>							
Male	0.03	0.03	0.03	0.05	0.05	0.04	0.04
Female	0.02	0.02	0.03	0.03	0.03	0.03	0.03

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.23P Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	
<b>AGE GROUP</b>							
12 to 17	0.6454	1.0000	0.2881	0.6070	0.7791	0.3531	
18 to 25	0.0000	0.0000	0.0000	0.0008	0.0781	0.0571	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0001	0.0008	
<b>GENDER</b>							
Male	0.0000	0.0000	0.0000	0.0000	0.0527	0.0001	
Female	0.0000	0.0000	0.0000	0.0000	0.0000	0.1324	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.0000	0.0000	0.0000	0.0024	0.0075	0.0320	
<b>AGE GROUP</b>							
12 to 17	0.3498	0.9851	0.8084	0.7512	0.4037	0.6617	
18 to 25	0.0000	0.0000	0.0000	0.0138	0.1721	0.0882	
26 or Older	0.0000	0.0000	0.0000	0.0145	0.0026	0.0873	
<b>GENDER</b>							
Male	0.0000	0.0000	0.0000	0.1268	0.0980	0.0838	
Female	0.0000	0.0000	0.0000	0.0008	0.0136	0.1729	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.0000	0.0000	0.0007	0.5222	0.9669	0.2591	
<b>AGE GROUP</b>							
12 to 17	0.4863	0.1847	0.9754	0.3640	0.2088	0.3114	
18 to 25	0.0001	0.0013	0.0006	0.2312	0.5030	0.3193	
26 or Older	0.0012	0.0000	0.0760	0.8603	0.9529	0.3325	
<b>GENDER</b>							
Male	0.0020	0.0018	0.0316	0.8053	0.5134	1.0000	
Female	0.0009	0.0002	0.0050	0.1531	0.3356	0.0889	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.24C Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	206	216	233	265	289	264	277
<b>AGE GROUP</b>							
12 to 17	24	24	21	25	21	22	24
18 to 25	61	54	54	55	62	54	63
26 or Older	190	207	221	254	279	258	265
<b>GENDER</b>							
Male	143	149	149	185	213	166	188
Female	140	162	181	178	193	204	202
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	138	140	151	149	184	138	152
<b>AGE GROUP</b>							
12 to 17	18	16	16	18	15	17	15
18 to 25	42	38	35	37	36	32	35
26 or Older	130	133	144	141	177	133	147
<b>GENDER</b>							
Male	102	98	95	106	142	86	97
Female	93	98	112	103	114	104	115
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	85	54	58	83	133	72	88
<b>AGE GROUP</b>							
12 to 17	10	7	6	7	8	7	7
18 to 25	17	16	18	18	16	16	18
26 or Older	81	51	55	79	132	71	88
<b>GENDER</b>							
Male	70	27	38	49	113	53	48
Female	41	45	44	66	73	49	74

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.24D Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.08	0.09	0.09	0.11	0.11	0.10	0.11
<b>AGE GROUP</b>							
12 to 17	0.10	0.09	0.08	0.10	0.09	0.09	0.10
18 to 25	0.19	0.17	0.16	0.16	0.18	0.16	0.18
26 or Older	0.10	0.11	0.12	0.13	0.14	0.13	0.13
<b>GENDER</b>							
Male	0.12	0.12	0.12	0.15	0.17	0.13	0.15
Female	0.11	0.13	0.14	0.14	0.15	0.15	0.15
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.06	0.06	0.06	0.06	0.07	0.05	0.06
<b>AGE GROUP</b>							
12 to 17	0.07	0.06	0.06	0.07	0.06	0.07	0.06
18 to 25	0.13	0.12	0.11	0.11	0.11	0.09	0.10
26 or Older	0.07	0.07	0.07	0.07	0.09	0.07	0.07
<b>GENDER</b>							
Male	0.09	0.08	0.08	0.09	0.12	0.07	0.08
Female	0.07	0.08	0.09	0.08	0.09	0.08	0.09
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.03	0.02	0.02	0.03	0.05	0.03	0.03
<b>AGE GROUP</b>							
12 to 17	0.04	0.03	0.02	0.03	0.03	0.03	0.03
18 to 25	0.05	0.05	0.05	0.05	0.05	0.05	0.05
26 or Older	0.04	0.03	0.03	0.04	0.07	0.04	0.04
<b>GENDER</b>							
Male	0.06	0.02	0.03	0.04	0.09	0.04	0.04
Female	0.03	0.04	0.03	0.05	0.06	0.04	0.06

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.



**Table 7.24P Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:  
P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.0000	0.0024	0.0381	0.7961	0.8610	0.3502	
<b>AGE GROUP</b>							
12 to 17	0.2901	0.5400	0.8877	0.6014	0.2760	0.9923	
18 to 25	0.5751	0.5129	0.6548	0.1733	0.2474	0.3410	
26 or Older	0.0000	0.0025	0.0260	0.5313	1.0000	0.4653	
<b>GENDER</b>							
Male	0.0013	0.0267	0.0172	0.9690	0.3582	0.2320	
Female	0.0000	0.0478	0.4934	0.7446	0.5029	0.8446	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.1507	0.1401	0.1731	0.0750	0.0541	0.7012	
<b>AGE GROUP</b>							
12 to 17	0.0273	0.1831	0.4032	0.3373	0.8083	0.1831	
18 to 25	0.0053	0.1137	0.0807	0.0133	0.3705	0.5593	
26 or Older	0.6084	0.3327	0.3499	0.2601	0.0745	0.6938	
<b>GENDER</b>							
Male	0.0992	0.1080	0.2416	0.0356	0.0400	0.8978	
Female	0.6529	0.6098	0.3955	0.6014	0.5199	0.5395	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.7039	0.2053	0.5450	0.3931	0.0596	0.7430	
<b>AGE GROUP</b>							
12 to 17	0.2237	0.9845	0.8550	0.6205	0.5942	0.6527	
18 to 25	0.6625	0.3864	0.0694	0.0905	0.4625	0.9776	
26 or Older	0.8551	0.1306	0.3015	0.6359	0.0806	0.7048	
<b>GENDER</b>							
Male	0.0776	0.3743	0.7152	0.0680	0.0255	0.8021	
Female	0.2083	0.3337	0.3265	0.8391	0.7772	0.5550	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.25C Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	283	260	274	258	288	254	286
<b>AGE GROUP</b>							
12 to 17	40	41	36	36	39	37	38
18 to 25	77	74	71	83	80	79	86
26 or Older	264	244	260	235	269	242	271
<b>GENDER</b>							
Male	229	203	223	193	236	215	244
Female	151	163	169	163	157	145	168
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	99	85	98	111	162	111	99
<b>AGE GROUP</b>							
12 to 17	30	29	26	25	27	25	27
18 to 25	38	37	31	35	37	40	40
26 or Older	87	72	91	100	158	100	85
<b>GENDER</b>							
Male	76	55	81	75	147	86	59
Female	64	66	56	80	70	71	78
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	51	46	53	60	67	60	70
<b>AGE GROUP</b>							
12 to 17	13	15	14	13	12	11	16
18 to 25	15	19	16	17	21	15	24
26 or Older	47	39	49	57	63	56	64
<b>GENDER</b>							
Male	45	34	43	37	52	43	31
Female	26	31	31	47	43	42	63

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.25D Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.11	0.10	0.11	0.10	0.11	0.10	0.11
<b>AGE GROUP</b>							
12 to 17	0.16	0.16	0.15	0.15	0.16	0.15	0.15
18 to 25	0.24	0.23	0.22	0.25	0.23	0.23	0.25
26 or Older	0.14	0.13	0.14	0.12	0.14	0.12	0.14
<b>GENDER</b>							
Male	0.19	0.17	0.18	0.16	0.19	0.17	0.19
Female	0.12	0.13	0.13	0.13	0.12	0.11	0.13
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.04	0.03	0.04	0.04	0.06	0.04	0.04
<b>AGE GROUP</b>							
12 to 17	0.12	0.12	0.10	0.10	0.11	0.10	0.11
18 to 25	0.12	0.11	0.10	0.11	0.11	0.12	0.12
26 or Older	0.05	0.04	0.05	0.05	0.08	0.05	0.04
<b>GENDER</b>							
Male	0.06	0.05	0.07	0.06	0.12	0.07	0.05
Female	0.05	0.05	0.04	0.06	0.05	0.05	0.06
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.02	0.02	0.02	0.02	0.03	0.02	0.03
<b>AGE GROUP</b>							
12 to 17	0.05	0.06	0.05	0.05	0.05	0.05	0.06
18 to 25	0.05	0.06	0.05	0.05	0.06	0.04	0.07
26 or Older	0.02	0.02	0.03	0.03	0.03	0.03	0.03
<b>GENDER</b>							
Male	0.04	0.03	0.04	0.03	0.04	0.03	0.02
Female	0.02	0.02	0.02	0.04	0.03	0.03	0.05

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.25P Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.8134	0.0326	0.2387	0.0977	0.8773	0.1475	
<b>AGE GROUP</b>							
12 to 17	0.0001	0.0114	0.0664	0.2526	0.1381	0.9407	
18 to 25	0.2904	0.0681	0.4677	0.8626	0.9094	0.4800	
26 or Older	0.5807	0.0363	0.2097	0.0455	0.7048	0.1826	
<b>GENDER</b>							
Male	0.9910	0.0217	0.0969	0.0138	0.9541	0.2087	
Female	0.6839	0.6450	0.7483	0.6537	0.8089	0.4982	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.0128	0.3971	0.1945	0.1607	0.0094	0.0579	
<b>AGE GROUP</b>							
12 to 17	0.0001	0.0042	0.0144	0.0819	0.0616	0.6104	
18 to 25	0.3613	0.7006	0.9694	0.8254	0.8663	0.5326	
26 or Older	0.3492	0.6926	0.5027	0.3464	0.0206	0.0856	
<b>GENDER</b>							
Male	0.0027	0.2670	0.0790	0.2296	0.0023	0.0805	
Female	0.5705	0.8447	0.9572	0.4143	0.8398	0.3387	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.6798	0.8815	0.9062	0.8641	0.4850	0.9086	
<b>AGE GROUP</b>							
12 to 17	0.1114	0.0968	0.4151	0.2702	0.6511	0.5940	
18 to 25	0.8831	0.5144	0.8715	0.7400	0.7195	0.4348	
26 or Older	0.9894	0.5099	0.9608	0.8353	0.6135	0.8052	
<b>GENDER</b>							
Male	0.0333	0.2169	0.1950	0.3749	0.1180	0.5337	
Female	0.2215	0.5161	0.3990	0.7264	0.7301	0.5576	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.26C Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	145	140	136	151	166	152	161
<b>AGE GROUP</b>							
12 to 17	14	12	12	12	15	13	10
18 to 25	50	47	40	38	39	41	40
26 or Older	133	129	126	145	160	147	156
<b>GENDER</b>							
Male	123	115	108	124	152	120	138
Female	69	68	73	76	64	90	75
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	38	31	28	32	36	49	42
<b>AGE GROUP</b>							
12 to 17	8	7	9	8	11	11	7
18 to 25	14	19	14	16	22	26	17
26 or Older	34	24	23	27	26	40	40
<b>GENDER</b>							
Male	35	27	25	23	33	39	37
Female	14	16	11	23	14	28	21
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	18	11	7	24	22	26	14
<b>AGE GROUP</b>							
12 to 17	4	5	5	6	5	9	4
18 to 25	6	6	6	9	8	9	6
26 or Older	16	8	*	21	20	22	12
<b>GENDER</b>							
Male	14	10	6	13	21	12	6
Female	10	4	4	20	8	23	12

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.26D Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.06	0.06	0.05	0.06	0.07	0.06	0.06
<b>AGE GROUP</b>							
12 to 17	0.05	0.05	0.05	0.05	0.06	0.05	0.04
18 to 25	0.15	0.14	0.12	0.11	0.11	0.12	0.11
26 or Older	0.07	0.07	0.07	0.07	0.08	0.07	0.08
<b>GENDER</b>							
Male	0.10	0.10	0.09	0.10	0.12	0.10	0.11
Female	0.05	0.05	0.06	0.06	0.05	0.07	0.06
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.02	0.01	0.01	0.01	0.01	0.02	0.02
<b>AGE GROUP</b>							
12 to 17	0.03	0.03	0.04	0.03	0.05	0.04	0.03
18 to 25	0.04	0.06	0.04	0.05	0.07	0.07	0.05
26 or Older	0.02	0.01	0.01	0.01	0.01	0.02	0.02
<b>GENDER</b>							
Male	0.03	0.02	0.02	0.02	0.03	0.03	0.03
Female	0.01	0.01	0.01	0.02	0.01	0.02	0.02
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.01	0.00	0.00	0.01	0.01	0.01	0.01
<b>AGE GROUP</b>							
12 to 17	0.02	0.02	0.02	0.03	0.02	0.04	0.02
18 to 25	0.02	0.02	0.02	0.03	0.02	0.03	0.02
26 or Older	0.01	0.00	*	0.01	0.01	0.01	0.01
<b>GENDER</b>							
Male	0.01	0.01	0.00	0.01	0.02	0.01	0.01
Female	0.01	0.00	0.00	0.02	0.01	0.02	0.01

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.26P Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.2439	0.0841	0.0465	0.2094	0.3989	0.2355	
<b>AGE GROUP</b>							
12 to 17	0.3728	0.8176	0.6564	0.4346	0.4076	0.1958	
18 to 25	0.0000	0.0000	0.0040	0.0370	0.2169	0.8436	
26 or Older	0.0004	0.0005	0.0039	0.0599	0.2119	0.1647	
<b>GENDER</b>							
Male	0.0557	0.0231	0.0051	0.0911	0.5346	0.0598	
Female	0.3106	0.7357	0.5204	0.7778	0.4685	0.5306	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.2550	0.2702	0.0159	0.1843	0.6219	0.3846	
<b>AGE GROUP</b>							
12 to 17	0.7027	0.6531	0.8875	0.9833	0.2573	0.1832	
18 to 25	0.3804	0.2359	0.2516	0.8910	0.1504	0.0632	
26 or Older	0.4180	0.0762	0.0368	0.1627	0.0800	0.8087	
<b>GENDER</b>							
Male	0.2682	0.2220	0.0319	0.0464	0.5796	0.6940	
Female	0.7360	0.9463	0.2630	0.6094	0.9762	0.2814	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.9061	0.4940	0.1126	0.2032	0.7389	0.3205	
<b>AGE GROUP</b>							
12 to 17	0.5743	0.5114	0.7123	0.2300	0.3524	0.0751	
18 to 25	0.4048	0.6080	0.5706	0.6929	0.9677	0.5751	
26 or Older	0.7521	0.3799	*	0.3786	0.9246	0.8267	
<b>GENDER</b>							
Male	0.3209	0.5009	0.7547	0.1004	0.3630	0.1901	
Female	0.4111	0.1184	0.0869	0.6297	0.4350	0.6611	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.27C DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	62	71	74	79	101	84	108
<b>AGE GROUP</b>							
12 to 17	9	10	10	10	10	10	12
18 to 25	29	24	28	33	36	36	43
26 or Older	54	65	68	73	94	74	98
<b>GENDER</b>							
Male	55	66	60	71	87	70	92
Female	29	23	42	33	53	50	53
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	21	16	27	29	35	32	36
<b>AGE GROUP</b>							
12 to 17	4	6	8	8	8	7	9
18 to 25	15	12	14	19	16	24	25
26 or Older	14	7	21	21	31	19	29
<b>GENDER</b>							
Male	21	14	23	19	28	27	33
Female	5	6	14	22	21	16	13
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	7	*	18	20	28	17	16
<b>AGE GROUP</b>							
12 to 17	*	*	4	4	7	3	6
18 to 25	5	5	9	4	7	8	11
26 or Older	*	*	15	19	27	14	9
<b>GENDER</b>							
Male	6	4	17	8	22	8	13
Female	4	*	*	18	18	15	9

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.



**Table 7.27D DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.03	0.03	0.03	0.03	0.04	0.03	0.04
<b>AGE GROUP</b>							
12 to 17	0.04	0.04	0.04	0.04	0.04	0.04	0.05
18 to 25	0.09	0.07	0.09	0.10	0.10	0.10	0.13
26 or Older	0.03	0.03	0.04	0.04	0.05	0.04	0.05
<b>GENDER</b>							
Male	0.05	0.05	0.05	0.06	0.07	0.06	0.07
Female	0.02	0.02	0.03	0.03	0.04	0.04	0.04
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.01	0.01	0.01	0.01	0.01	0.01	0.01
<b>AGE GROUP</b>							
12 to 17	0.02	0.02	0.03	0.03	0.03	0.03	0.04
18 to 25	0.05	0.04	0.04	0.06	0.05	0.07	0.07
26 or Older	0.01	0.00	0.01	0.01	0.02	0.01	0.01
<b>GENDER</b>							
Male	0.02	0.01	0.02	0.02	0.02	0.02	0.03
Female	0.00	0.00	0.01	0.02	0.02	0.01	0.01
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.00	*	0.01	0.01	0.01	0.01	0.01
<b>AGE GROUP</b>							
12 to 17	*	*	0.02	0.02	0.03	0.01	0.02
18 to 25	0.02	0.01	0.03	0.01	0.02	0.02	0.03
26 or Older	*	*	0.01	0.01	0.01	0.01	0.00
<b>GENDER</b>							
Male	0.00	0.00	0.01	0.01	0.02	0.01	0.01
Female	0.00	*	*	0.01	0.01	0.01	0.01

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.27P DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.0000	0.0000	0.0000	0.0000	0.1247	0.0058	
<b>AGE GROUP</b>							
12 to 17	0.0011	0.0005	0.0021	0.0170	0.0287	0.0324	
18 to 25	0.0000	0.0000	0.0000	0.0000	0.0033	0.0129	
26 or Older	0.0006	0.0023	0.0060	0.0221	0.8208	0.0926	
<b>GENDER</b>							
Male	0.0000	0.0000	0.0000	0.0001	0.1205	0.0154	
Female	0.0033	0.0005	0.1331	0.0536	0.6791	0.1567	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.0000	0.0000	0.0001	0.0081	0.0919	0.2246	
<b>AGE GROUP</b>							
12 to 17	0.0002	0.0020	0.0547	0.2413	0.2641	0.1431	
18 to 25	0.0000	0.0000	0.0000	0.0080	0.0084	0.6908	
26 or Older	0.1258	0.0309	0.4875	0.4468	0.8855	0.4241	
<b>GENDER</b>							
Male	0.0000	0.0000	0.0002	0.0005	0.0372	0.3391	
Female	0.0005	0.0004	0.1269	0.7114	0.8473	0.4092	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.0017	*	0.2617	0.5341	0.8113	0.2212	
<b>AGE GROUP</b>							
12 to 17	*	*	0.1704	0.3966	0.9793	0.1085	
18 to 25	0.0134	0.0032	0.0703	0.0044	0.0540	0.1262	
26 or Older	*	*	0.6489	0.2800	0.2607	0.8686	
<b>GENDER</b>							
Male	0.0170	0.0030	0.8328	0.1089	0.8590	0.1913	
Female	0.0275	*	*	0.7105	0.8748	0.6189	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.28C Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	92	110	129	159	189	175	182
<b>AGE GROUP</b>							
12 to 17	20	23	27	28	28	25	23
18 to 25	62	68	75	89	95	92	94
26 or Older	59	78	98	120	155	139	152
<b>GENDER</b>							
Male	81	91	104	142	159	142	167
Female	34	54	62	63	99	97	77
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	48	68	70	72	77	52	51
<b>AGE GROUP</b>							
12 to 17	16	18	23	22	22	19	16
18 to 25	38	45	47	47	42	39	33
26 or Older	22	43	43	46	59	29	34
<b>GENDER</b>							
Male	41	55	55	58	73	43	46
Female	21	34	37	37	28	28	23
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	18	26	24	28	28	20	26
<b>AGE GROUP</b>							
12 to 17	7	9	10	9	9	8	7
18 to 25	16	21	21	15	17	18	11
26 or Older	*	11	8	22	20	7	22
<b>GENDER</b>							
Male	17	22	22	19	27	17	24
Female	7	11	8	20	9	10	10

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.28D Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2012**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.04	0.04	0.05	0.06	0.07	0.07	0.07
<b>AGE GROUP</b>							
12 to 17	0.08	0.09	0.11	0.12	0.12	0.10	0.09
18 to 25	0.19	0.21	0.23	0.26	0.28	0.27	0.27
26 or Older	0.03	0.04	0.05	0.06	0.08	0.07	0.08
<b>GENDER</b>							
Male	0.07	0.08	0.09	0.12	0.13	0.11	0.13
Female	0.03	0.04	0.05	0.05	0.08	0.07	0.06
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.02	0.03	0.03	0.03	0.03	0.02	0.02
<b>AGE GROUP</b>							
12 to 17	0.06	0.07	0.09	0.09	0.09	0.07	0.06
18 to 25	0.12	0.14	0.14	0.14	0.12	0.11	0.10
26 or Older	0.01	0.02	0.02	0.02	0.03	0.01	0.02
<b>GENDER</b>							
Male	0.03	0.05	0.05	0.05	0.06	0.03	0.04
Female	0.02	0.03	0.03	0.03	0.02	0.02	0.02
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.01	0.01	0.01	0.01	0.01	0.01	0.01
<b>AGE GROUP</b>							
12 to 17	0.03	0.03	0.04	0.04	0.04	0.03	0.03
18 to 25	0.05	0.06	0.06	0.04	0.05	0.05	0.03
26 or Older	*	0.01	0.00	0.01	0.01	0.00	0.01
<b>GENDER</b>							
Male	0.01	0.02	0.02	0.02	0.02	0.01	0.02
Female	0.01	0.01	0.01	0.02	0.01	0.01	0.01

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.

**Table 7.28P Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2006-2011**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>
<b>LIFETIME USE</b>							
<b>TOTAL</b>	0.0000	0.0000	0.0000	0.0074	0.5924	0.6510	
<b>AGE GROUP</b>							
12 to 17	0.1667	0.3912	0.0004	0.0000	0.0001	0.0022	
18 to 25	0.0000	0.0000	0.0000	0.9463	0.0566	0.0570	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.2171	0.2139	
<b>GENDER</b>							
Male	0.0000	0.0000	0.0000	0.0771	0.6908	0.5913	
Female	0.0000	0.0000	0.0001	0.0059	0.7694	0.0572	
<b>PAST YEAR USE</b>							
<b>TOTAL</b>	0.0305	0.0000	0.0000	0.0000	0.0000	0.0017	
<b>AGE GROUP</b>							
12 to 17	0.4309	0.0114	0.0000	0.0000	0.0001	0.0015	
18 to 25	0.0001	0.0000	0.0000	0.0000	0.0000	0.0054	
26 or Older	0.0638	0.6375	0.1416	0.0344	0.0105	0.9484	
<b>GENDER</b>							
Male	0.0070	0.0000	0.0000	0.0000	0.0000	0.0057	
Female	0.6606	0.0177	0.0002	0.0010	0.0188	0.0933	
<b>PAST MONTH USE</b>							
<b>TOTAL</b>	0.4899	0.0157	0.0199	0.0415	0.2397	0.6448	
<b>AGE GROUP</b>							
12 to 17	0.7494	0.1208	0.0051	0.0214	0.0746	0.2011	
18 to 25	0.0023	0.0001	0.0004	0.0093	0.0160	0.0717	
26 or Older	*	0.3382	0.0971	0.9649	0.4333	0.1230	
<b>GENDER</b>							
Male	0.3086	0.0349	0.0167	0.1093	0.1747	0.8073	
Female	0.5139	0.2393	0.6656	0.2092	0.7472	0.5926	

\*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2012.