

**Table 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	832	803	794	845	850	844	871	831	852	846	882
Cigarettes	795	789	768	777	807	809	810	797	780	816	839
Smokeless Tobacco	279	284	245	267	284	283	312	291	314	278	298
Cigars	350	331	323	354	352	337	366	361	363	373	388
Pipe Tobacco	166	131	147	146	167	170	147	151	147	144	184
<b>ALCOHOL</b>	989	919	957	975	968	1,019	983	957	990	1,006	1,016
Binge Alcohol Use <sup>2</sup>	732	682	698	701	746	770	735	783	752	734	797
Heavy Alcohol Use <sup>2</sup>	392	380	382	379	427	429	414	406	422	423	448
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	572	573	570	576	585	598	628	602	626	581	618
Cigarettes	527	545	537	515	545	577	565	565	536	537	556
Smokeless Tobacco	264	256	234	241	273	252	296	283	302	264	281
Cigars	327	305	290	324	320	312	333	336	321	336	351
Pipe Tobacco	139	127	139	141	160	168	131	144	137	135	170
<b>ALCOHOL</b>	656	599	626	621	610	666	632	651	675	674	675
Binge Alcohol Use <sup>2</sup>	540	517	513	529	559	585	586	605	577	567	625
Heavy Alcohol Use <sup>2</sup>	346	321	319	321	354	374	357	349	353	355	382
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	513	507	497	543	511	511	558	505	535	542	554
Cigarettes	501	488	491	538	503	493	553	498	526	532	539
Smokeless Tobacco	100	126	65	87	76	113	75	51	78	82	103
Cigars	114	123	122	112	131	116	114	130	153	142	146
Pipe Tobacco	90	35	43	42	46	30	55	37	54	46	61
<b>ALCOHOL</b>	644	600	664	637	645	676	673	687	635	684	686
Binge Alcohol Use <sup>2</sup>	415	382	390	402	399	395	398	441	441	441	427
Heavy Alcohol Use <sup>2</sup>	177	174	174	163	169	171	192	193	201	207	203

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.35	0.34	0.33	0.35	0.35	0.34	0.35	0.33	0.34	0.33	0.34
Cigarettes	0.34	0.33	0.32	0.32	0.33	0.33	0.32	0.32	0.31	0.32	0.32
Smokeless Tobacco	0.12	0.12	0.10	0.11	0.12	0.11	0.12	0.12	0.12	0.11	0.11
Cigars	0.15	0.14	0.13	0.15	0.14	0.14	0.15	0.14	0.14	0.14	0.15
Pipe Tobacco	0.07	0.06	0.06	0.06	0.07	0.07	0.06	0.06	0.06	0.06	0.07
<b>ALCOHOL</b>	0.42	0.39	0.40	0.40	0.39	0.41	0.39	0.38	0.39	0.39	0.39
Binge Alcohol Use <sup>2</sup>	0.31	0.29	0.29	0.29	0.30	0.31	0.29	0.31	0.30	0.29	0.31
Heavy Alcohol Use <sup>2</sup>	0.17	0.16	0.16	0.16	0.17	0.17	0.17	0.16	0.17	0.16	0.17
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.50	0.50	0.49	0.49	0.49	0.50	0.52	0.49	0.51	0.47	0.49
Cigarettes	0.46	0.47	0.46	0.44	0.46	0.48	0.47	0.46	0.43	0.43	0.44
Smokeless Tobacco	0.23	0.22	0.20	0.20	0.23	0.21	0.24	0.23	0.25	0.21	0.22
Cigars	0.29	0.27	0.25	0.27	0.27	0.26	0.27	0.27	0.26	0.27	0.28
Pipe Tobacco	0.12	0.11	0.12	0.12	0.13	0.14	0.11	0.12	0.11	0.11	0.14
<b>ALCOHOL</b>	0.58	0.52	0.54	0.53	0.51	0.55	0.52	0.53	0.55	0.54	0.54
Binge Alcohol Use <sup>2</sup>	0.48	0.45	0.44	0.45	0.47	0.49	0.48	0.49	0.47	0.45	0.50
Heavy Alcohol Use <sup>2</sup>	0.30	0.28	0.27	0.27	0.30	0.31	0.29	0.29	0.29	0.28	0.30
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.42	0.41	0.40	0.43	0.40	0.40	0.43	0.39	0.41	0.41	0.41
Cigarettes	0.41	0.40	0.40	0.43	0.40	0.39	0.43	0.38	0.40	0.40	0.40
Smokeless Tobacco	0.08	0.10	0.05	0.07	0.06	0.09	0.06	0.04	0.06	0.06	0.08
Cigars	0.09	0.10	0.10	0.09	0.10	0.09	0.09	0.10	0.12	0.11	0.11
Pipe Tobacco	0.07	0.03	0.03	0.03	0.04	0.02	0.04	0.03	0.04	0.03	0.05
<b>ALCOHOL</b>	0.53	0.49	0.54	0.51	0.51	0.53	0.52	0.53	0.49	0.51	0.51
Binge Alcohol Use <sup>2</sup>	0.34	0.31	0.31	0.32	0.32	0.31	0.31	0.34	0.34	0.33	0.32
Heavy Alcohol Use <sup>2</sup>	0.15	0.14	0.14	0.13	0.13	0.13	0.15	0.15	0.15	0.16	0.15

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2012 Versus 2002-2011**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0266	0.1282	0.6014	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0058	0.0515	0.8854	
Smokeless Tobacco	0.3230	0.1757	0.0012	0.0519	0.4202	0.1540	0.9516	0.7461	0.8772	0.0804	
Cigars	0.2116	0.2363	0.0065	0.0342	0.0509	0.2938	0.5206	0.5079	0.7177	0.4289	
Pipe Tobacco	0.0451	0.0012	0.0252	0.4343	0.7446	0.1489	0.0201	0.1270	0.1760	0.1109	
<b>ALCOHOL</b>	0.0458	0.0002	0.0011	0.6047	0.0400	0.0906	0.3700	0.6771	0.5721	0.5527	
Binge Alcohol Use <sup>2</sup>	0.8607	0.4348	0.6399	0.4673	0.8400	0.3604	0.3197	0.0848	0.7584	0.4605	
Heavy Alcohol Use <sup>2</sup>	0.4048	0.2932	0.0933	0.8379	0.1493	0.1887	0.0872	0.2409	0.6120	0.0933	
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0001	0.0001	0.0000	0.0010	0.0279	0.4617	0.3090	0.3000	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0105	0.2995	0.2207	0.6251	
Smokeless Tobacco	0.2469	0.0676	0.0013	0.0310	0.5717	0.1470	0.8728	0.9010	0.9269	0.0551	
Cigars	0.0345	0.2165	0.0011	0.0061	0.0544	0.1234	0.2022	0.5507	0.9397	0.3368	
Pipe Tobacco	0.0885	0.0218	0.1445	0.8791	0.6786	0.5291	0.0276	0.3683	0.2295	0.1824	
<b>ALCOHOL</b>	0.2775	0.2911	0.6099	0.0436	0.5270	0.8928	0.1445	0.1912	0.2953	0.7798	
Binge Alcohol Use <sup>2</sup>	0.2413	0.4326	0.3137	0.8270	0.1839	0.0583	0.0695	0.0741	0.4717	0.5309	
Heavy Alcohol Use <sup>2</sup>	0.0487	0.2299	0.0739	0.2990	0.0524	0.0991	0.0714	0.2938	0.6374	0.1842	
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0001	0.0000	0.0000	0.0031	0.0027	0.0095	0.2485	0.7270	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0008	0.0017	0.0993	0.7659	
Smokeless Tobacco	0.6858	0.3926	0.5133	0.9797	0.3209	0.7245	0.6230	0.3395	0.9191	0.8057	
Cigars	0.0606	0.8595	0.5550	0.2548	0.6388	0.2581	0.0913	0.8502	0.5886	0.8054	
Pipe Tobacco	0.2750	0.0005	0.0051	0.0390	0.0176	0.0009	0.3529	0.0222	0.4361	0.2995	
<b>ALCOHOL</b>	0.0000	0.0000	0.0000	0.0046	0.0002	0.0081	0.0053	0.0536	0.0439	0.2373	
Binge Alcohol Use <sup>2</sup>	0.0613	0.0121	0.0232	0.1032	0.1078	0.2431	0.3401	0.5930	0.5448	0.6774	
Heavy Alcohol Use <sup>2</sup>	0.0607	0.9284	0.7189	0.1241	0.6347	0.7123	0.7745	0.6020	0.9375	0.2658	

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.30C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	81	80	80	78	72	76	69	71	69	68	61
Cigarettes	74	73	76	71	66	67	61	64	63	60	55
Smokeless Tobacco	27	30	32	29	31	36	30	32	30	30	31
Cigars	47	44	44	46	41	45	41	39	36	41	32
Pipe Tobacco	14	18	19	17	18	19	17	21	17	18	18
<b>ALCOHOL</b>	80	82	81	81	82	87	79	79	80	77	78
Binge Alcohol Use <sup>2</sup>	67	67	73	65	69	69	57	62	62	55	56
Heavy Alcohol Use <sup>2</sup>	31	33	36	32	33	37	30	30	28	25	25
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	59	56	55	59	54	57	50	52	52	53	47
Cigarettes	53	47	52	53	46	48	42	44	45	46	41
Smokeless Tobacco	26	29	30	28	29	34	28	30	28	28	29
Cigars	37	36	37	38	34	37	34	33	31	31	26
Pipe Tobacco	13	17	15	14	15	16	12	18	14	14	12
<b>ALCOHOL</b>	55	57	59	57	57	61	53	58	57	53	55
Binge Alcohol Use <sup>2</sup>	46	46	52	48	47	51	41	48	44	42	41
Heavy Alcohol Use <sup>2</sup>	24	23	27	27	26	28	23	23	23	22	19
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	51	52	53	50	47	48	46	46	43	43	37
Cigarettes	49	50	52	48	46	46	44	44	40	40	34
Smokeless Tobacco	9	6	11	8	10	8	10	9	12	10	8
Cigars	26	23	25	24	24	22	21	25	20	24	19
Pipe Tobacco	9	7	13	11	10	11	12	11	10	12	12
<b>ALCOHOL</b>	58	60	59	58	59	61	56	52	54	53	54
Binge Alcohol Use <sup>2</sup>	46	46	50	43	49	45	42	40	43	37	38
Heavy Alcohol Use <sup>2</sup>	19	23	23	20	21	23	20	19	17	15	16

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.30D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.33	0.32	0.32	0.31	0.29	0.30	0.28	0.29	0.28	0.27	0.25
Cigarettes	0.30	0.29	0.30	0.28	0.26	0.27	0.25	0.26	0.26	0.24	0.22
Smokeless Tobacco	0.11	0.12	0.13	0.12	0.12	0.14	0.12	0.13	0.12	0.12	0.12
Cigars	0.19	0.17	0.18	0.18	0.16	0.18	0.16	0.16	0.15	0.16	0.13
Pipe Tobacco	0.06	0.07	0.08	0.07	0.07	0.08	0.07	0.09	0.07	0.07	0.07
<b>ALCOHOL</b>	0.32	0.33	0.32	0.32	0.32	0.34	0.32	0.32	0.33	0.31	0.31
Binge Alcohol Use <sup>2</sup>	0.27	0.27	0.29	0.26	0.27	0.27	0.23	0.25	0.25	0.22	0.22
Heavy Alcohol Use <sup>2</sup>	0.12	0.13	0.14	0.13	0.13	0.15	0.12	0.12	0.11	0.10	0.10
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.47	0.44	0.43	0.46	0.41	0.45	0.39	0.42	0.42	0.41	0.37
Cigarettes	0.42	0.37	0.40	0.41	0.36	0.37	0.33	0.35	0.36	0.36	0.32
Smokeless Tobacco	0.21	0.23	0.23	0.22	0.22	0.26	0.22	0.24	0.23	0.22	0.23
Cigars	0.30	0.28	0.28	0.29	0.26	0.29	0.27	0.27	0.25	0.25	0.20
Pipe Tobacco	0.10	0.13	0.11	0.11	0.12	0.12	0.10	0.14	0.11	0.11	0.09
<b>ALCOHOL</b>	0.43	0.45	0.45	0.44	0.44	0.47	0.41	0.46	0.46	0.42	0.43
Binge Alcohol Use <sup>2</sup>	0.37	0.36	0.40	0.37	0.36	0.40	0.32	0.38	0.35	0.33	0.32
Heavy Alcohol Use <sup>2</sup>	0.19	0.18	0.21	0.21	0.20	0.22	0.18	0.18	0.18	0.17	0.15
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.42	0.42	0.43	0.40	0.38	0.39	0.38	0.38	0.36	0.35	0.30
Cigarettes	0.40	0.41	0.42	0.39	0.37	0.37	0.36	0.37	0.34	0.33	0.28
Smokeless Tobacco	0.08	0.05	0.09	0.07	0.08	0.07	0.08	0.08	0.11	0.08	0.06
Cigars	0.21	0.19	0.20	0.19	0.19	0.18	0.17	0.21	0.17	0.20	0.16
Pipe Tobacco	0.07	0.06	0.10	0.09	0.08	0.09	0.10	0.09	0.09	0.10	0.10
<b>ALCOHOL</b>	0.48	0.49	0.48	0.47	0.47	0.49	0.46	0.43	0.45	0.43	0.44
Binge Alcohol Use <sup>2</sup>	0.38	0.38	0.41	0.35	0.39	0.37	0.35	0.33	0.36	0.30	0.31
Heavy Alcohol Use <sup>2</sup>	0.16	0.19	0.19	0.16	0.17	0.18	0.16	0.15	0.14	0.12	0.13

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.30P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2002-2011**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	
Smokeless Tobacco	0.5518	0.8872	0.2610	0.9831	0.0797	0.0369	0.3486	0.1010	0.1155	0.9810	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.0000	
Pipe Tobacco	0.2450	0.8510	0.6620	0.7110	0.9855	0.5764	0.5922	0.0133	0.8116	0.5068	
<b>ALCOHOL</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1119	0.3572	
Binge Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0650	0.5661	
Heavy Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0143	0.1412	
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0048	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0033	
Smokeless Tobacco	0.4375	0.8606	0.2752	0.9588	0.0990	0.0360	0.4631	0.1698	0.3476	0.8071	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0099	0.0051	
Pipe Tobacco	0.8856	0.0984	0.2308	0.2433	0.1976	0.1567	0.4742	0.0005	0.4516	0.4477	
<b>ALCOHOL</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0040	0.0000	0.0799	0.2691	
Binge Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0000	0.0509	0.3038	
Heavy Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0040	0.0545	
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0197	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0201	
Smokeless Tobacco	0.6172	0.1695	0.7588	0.9122	0.5000	0.7124	0.4017	0.2075	0.0496	0.3920	
Cigars	0.0001	0.0000	0.0000	0.0002	0.0000	0.0028	0.0159	0.0000	0.0438	0.0016	
Pipe Tobacco	0.0546	0.0058	0.5295	0.0704	0.1205	0.3835	0.9607	0.5855	0.2115	0.8437	
<b>ALCOHOL</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0029	0.0583	0.5959	0.8087	
Binge Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0004	0.0252	0.5134	0.7836	
Heavy Alcohol Use <sup>2</sup>	0.0005	0.0000	0.0001	0.0028	0.0017	0.0070	0.0474	0.0016	0.7709	0.8153	

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.31C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	150	151	161	155	160	162	153	169	166	167	164
Cigarettes	150	151	159	152	156	159	149	162	160	162	161
Smokeless Tobacco	59	56	60	64	65	63	68	78	81	70	70
Cigars	83	84	97	92	94	92	96	99	101	100	93
Pipe Tobacco	24	25	30	37	33	32	34	40	41	48	39
<b>ALCOHOL</b>	164	160	164	166	168	170	163	174	169	184	170
Binge Alcohol Use <sup>2</sup>	161	155	166	171	169	169	167	182	168	189	175
Heavy Alcohol Use <sup>2</sup>	112	114	120	120	125	114	124	128	127	125	121
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	100	106	106	110	109	116	109	121	121	113	118
Cigarettes	99	109	105	107	108	116	106	121	113	111	113
Smokeless Tobacco	56	55	58	60	62	61	63	73	75	65	65
Cigars	73	72	83	79	78	81	80	81	84	89	80
Pipe Tobacco	22	21	27	33	30	30	29	35	33	39	34
<b>ALCOHOL</b>	111	100	110	101	108	110	109	117	112	119	118
Binge Alcohol Use <sup>2</sup>	110	105	113	111	115	116	118	130	121	126	124
Heavy Alcohol Use <sup>2</sup>	87	91	88	96	95	94	95	99	101	87	97
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	101	93	102	101	102	97	101	98	96	105	104
Cigarettes	102	92	101	101	98	95	100	96	97	102	102
Smokeless Tobacco	8	11	10	12	13	13	14	18	20	14	14
Cigars	38	43	44	41	44	43	47	45	49	50	46
Pipe Tobacco	12	12	12	16	11	14	17	15	23	26	19
<b>ALCOHOL</b>	102	114	104	112	112	114	107	110	109	123	118
Binge Alcohol Use <sup>2</sup>	99	100	96	104	103	110	102	105	101	116	110
Heavy Alcohol Use <sup>2</sup>	55	59	60	56	63	60	65	61	59	72	63

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.31D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.48	0.48	0.50	0.48	0.49	0.50	0.47	0.50	0.49	0.49	0.47
Cigarettes	0.48	0.47	0.49	0.47	0.48	0.49	0.45	0.48	0.47	0.47	0.47
Smokeless Tobacco	0.19	0.18	0.19	0.20	0.20	0.19	0.20	0.23	0.24	0.20	0.20
Cigars	0.27	0.26	0.30	0.28	0.29	0.28	0.29	0.29	0.30	0.29	0.27
Pipe Tobacco	0.08	0.08	0.09	0.11	0.10	0.10	0.10	0.12	0.12	0.14	0.11
<b>ALCOHOL</b>	0.53	0.50	0.51	0.51	0.51	0.52	0.49	0.52	0.50	0.54	0.49
Binge Alcohol Use <sup>2</sup>	0.52	0.49	0.52	0.53	0.52	0.52	0.51	0.54	0.49	0.55	0.51
Heavy Alcohol Use <sup>2</sup>	0.36	0.36	0.37	0.37	0.38	0.35	0.38	0.38	0.37	0.36	0.35
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.64	0.67	0.66	0.67	0.66	0.70	0.66	0.72	0.70	0.66	0.68
Cigarettes	0.64	0.69	0.65	0.66	0.66	0.70	0.64	0.71	0.65	0.65	0.65
Smokeless Tobacco	0.36	0.34	0.36	0.37	0.38	0.37	0.38	0.43	0.43	0.38	0.38
Cigars	0.47	0.45	0.51	0.48	0.48	0.49	0.48	0.48	0.49	0.52	0.46
Pipe Tobacco	0.14	0.13	0.17	0.20	0.18	0.18	0.17	0.21	0.19	0.23	0.20
<b>ALCOHOL</b>	0.71	0.63	0.68	0.62	0.66	0.67	0.66	0.69	0.65	0.69	0.68
Binge Alcohol Use <sup>2</sup>	0.71	0.66	0.70	0.68	0.70	0.70	0.71	0.77	0.70	0.73	0.71
Heavy Alcohol Use <sup>2</sup>	0.56	0.57	0.54	0.59	0.58	0.57	0.57	0.59	0.59	0.51	0.56
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.65	0.59	0.64	0.63	0.63	0.59	0.62	0.59	0.57	0.61	0.60
Cigarettes	0.66	0.58	0.63	0.63	0.60	0.58	0.61	0.58	0.58	0.60	0.59
Smokeless Tobacco	0.05	0.07	0.06	0.08	0.08	0.08	0.08	0.11	0.12	0.08	0.08
Cigars	0.25	0.27	0.27	0.26	0.27	0.26	0.29	0.27	0.29	0.29	0.27
Pipe Tobacco	0.08	0.08	0.08	0.10	0.07	0.08	0.10	0.09	0.14	0.15	0.11
<b>ALCOHOL</b>	0.66	0.72	0.65	0.69	0.69	0.70	0.65	0.66	0.65	0.72	0.68
Binge Alcohol Use <sup>2</sup>	0.64	0.63	0.60	0.65	0.64	0.68	0.62	0.63	0.60	0.68	0.64
Heavy Alcohol Use <sup>2</sup>	0.35	0.37	0.37	0.35	0.39	0.37	0.39	0.36	0.35	0.42	0.37

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.



**Table 7.31P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2002-2011**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0433	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0110	
Smokeless Tobacco	0.0136	0.0017	0.0316	0.1505	0.3086	0.3184	0.5380	0.0570	0.0055	0.7737	
Cigars	0.4765	0.0738	0.0000	0.0018	0.0006	0.0031	0.0837	0.0534	0.1563	0.7892	
Pipe Tobacco	0.0000	0.0000	0.0001	0.0278	0.0012	0.0001	0.0068	0.7973	0.9497	0.8088	
<b>ALCOHOL</b>	0.7255	0.1061	0.6958	0.3193	0.0101	0.1590	0.2138	0.0279	0.0916	0.4827	
Binge Alcohol Use <sup>2</sup>	0.0483	0.0033	0.0172	0.0012	0.0001	0.0010	0.0144	0.0018	0.1616	0.6487	
Heavy Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0339	0.0833	0.2155	
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0009	0.0320	0.0014	0.0229	0.3809	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0014	0.0001	0.0978	0.0527	
Smokeless Tobacco	0.0322	0.0020	0.0512	0.1499	0.3241	0.2569	0.5591	0.1192	0.0129	0.7135	
Cigars	0.5231	0.1326	0.0000	0.0023	0.0007	0.0012	0.1054	0.0769	0.4978	0.5805	
Pipe Tobacco	0.0000	0.0000	0.0198	0.1592	0.0505	0.0042	0.0147	0.7771	0.3434	0.8321	
<b>ALCOHOL</b>	0.0174	0.0000	0.0334	0.0001	0.0009	0.0138	0.1824	0.0023	0.0026	0.6857	
Binge Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0098	0.0002	0.0202	0.5499	
Heavy Alcohol Use <sup>2</sup>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0018	0.0018	0.0396	0.2389	0.1600	
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0234	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0573	
Smokeless Tobacco	0.0156	0.3443	0.0778	0.5942	0.4237	0.9961	0.5472	0.0512	0.1937	0.6715	
Cigars	0.6983	0.3346	0.0837	0.2082	0.3076	0.9153	0.5150	0.4081	0.1516	0.1154	
Pipe Tobacco	0.0003	0.0001	0.0000	0.0345	0.0005	0.0017	0.2343	0.2247	0.1141	0.3955	
<b>ALCOHOL</b>	0.0529	0.0747	0.1066	0.0310	0.6109	0.7344	0.6328	0.8842	0.5276	0.5527	
Binge Alcohol Use <sup>2</sup>	0.1002	0.1229	0.2974	0.1632	0.3057	0.4992	0.3750	0.4809	0.5158	0.9432	
Heavy Alcohol Use <sup>2</sup>	0.1566	0.0516	0.0918	0.0856	0.0001	0.0022	0.0232	0.3697	0.1880	0.9168	

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.32C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	797	748	752	798	799	801	849	784	813	805	841
Cigarettes	754	738	718	726	760	763	785	753	747	771	796
Smokeless Tobacco	270	274	236	256	267	268	303	274	302	266	286
Cigars	331	332	311	335	332	312	352	341	340	352	372
Pipe Tobacco	160	126	142	141	162	166	141	141	141	135	176
<b>ALCOHOL</b>	953	887	933	936	914	986	958	931	946	978	972
Binge Alcohol Use <sup>2</sup>	702	639	654	650	697	714	703	751	715	699	749
Heavy Alcohol Use <sup>2</sup>	355	349	356	347	392	402	391	377	398	396	420
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	553	546	552	554	566	575	613	574	599	560	605
Cigarettes	506	521	512	493	527	551	546	538	514	514	545
Smokeless Tobacco	252	247	226	232	258	238	286	267	290	252	267
Cigars	311	302	280	309	305	290	325	318	306	321	334
Pipe Tobacco	132	124	135	136	156	164	128	136	134	129	165
<b>ALCOHOL</b>	630	577	610	598	583	640	616	631	645	665	652
Binge Alcohol Use <sup>2</sup>	527	492	492	507	541	562	565	585	552	552	594
Heavy Alcohol Use <sup>2</sup>	323	306	307	302	333	355	340	326	337	343	363
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	488	477	469	515	477	486	543	485	516	520	525
Cigarettes	475	460	466	508	472	469	539	481	509	510	512
Smokeless Tobacco	99	125	63	85	74	111	73	49	72	80	102
Cigars	99	117	107	98	116	104	101	116	140	130	135
Pipe Tobacco	89	32	39	37	43	26	50	32	47	36	57
<b>ALCOHOL</b>	628	582	654	615	614	658	656	668	620	663	662
Binge Alcohol Use <sup>2</sup>	393	347	361	370	356	364	382	422	427	406	406
Heavy Alcohol Use <sup>2</sup>	158	155	160	146	147	157	181	181	190	184	192

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.32D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.44	0.41	0.41	0.43	0.43	0.42	0.44	0.40	0.42	0.41	0.42
Cigarettes	0.42	0.41	0.39	0.39	0.40	0.40	0.41	0.39	0.38	0.39	0.40
Smokeless Tobacco	0.15	0.15	0.13	0.14	0.14	0.14	0.16	0.14	0.15	0.13	0.14
Cigars	0.18	0.18	0.17	0.18	0.18	0.16	0.18	0.18	0.17	0.18	0.19
Pipe Tobacco	0.09	0.07	0.08	0.08	0.09	0.09	0.07	0.07	0.07	0.07	0.09
<b>ALCOHOL</b>	0.53	0.49	0.51	0.51	0.49	0.52	0.50	0.48	0.48	0.49	0.48
Binge Alcohol Use <sup>2</sup>	0.39	0.35	0.36	0.35	0.37	0.38	0.37	0.39	0.37	0.35	0.37
Heavy Alcohol Use <sup>2</sup>	0.20	0.19	0.19	0.19	0.21	0.21	0.20	0.19	0.20	0.20	0.21
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.65	0.63	0.63	0.62	0.63	0.63	0.67	0.62	0.64	0.59	0.63
Cigarettes	0.59	0.60	0.59	0.56	0.59	0.61	0.59	0.58	0.55	0.54	0.57
Smokeless Tobacco	0.29	0.29	0.26	0.26	0.29	0.26	0.31	0.29	0.31	0.27	0.28
Cigars	0.36	0.35	0.32	0.35	0.34	0.32	0.35	0.34	0.33	0.34	0.35
Pipe Tobacco	0.15	0.14	0.15	0.15	0.17	0.18	0.14	0.15	0.14	0.14	0.17
<b>ALCOHOL</b>	0.74	0.67	0.70	0.67	0.65	0.70	0.67	0.68	0.69	0.70	0.68
Binge Alcohol Use <sup>2</sup>	0.62	0.57	0.56	0.57	0.60	0.62	0.61	0.63	0.59	0.58	0.62
Heavy Alcohol Use <sup>2</sup>	0.38	0.35	0.35	0.34	0.37	0.39	0.37	0.35	0.36	0.36	0.38
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.52	0.50	0.49	0.53	0.49	0.49	0.54	0.48	0.51	0.50	0.50
Cigarettes	0.51	0.49	0.49	0.53	0.48	0.47	0.54	0.48	0.50	0.49	0.49
Smokeless Tobacco	0.11	0.13	0.07	0.09	0.08	0.11	0.07	0.05	0.07	0.08	0.10
Cigars	0.11	0.12	0.11	0.10	0.12	0.11	0.10	0.12	0.14	0.13	0.13
Pipe Tobacco	0.10	0.03	0.04	0.04	0.04	0.03	0.05	0.03	0.05	0.03	0.05
<b>ALCOHOL</b>	0.67	0.61	0.68	0.64	0.63	0.66	0.66	0.66	0.61	0.64	0.63
Binge Alcohol Use <sup>2</sup>	0.42	0.37	0.38	0.38	0.36	0.37	0.38	0.42	0.42	0.39	0.39
Heavy Alcohol Use <sup>2</sup>	0.17	0.16	0.17	0.15	0.15	0.16	0.18	0.18	0.19	0.18	0.18

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.32P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2002-2011**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0002	0.0095	0.0011	0.0001	0.0072	0.0280	0.5611	0.7466	0.2191	
Cigarettes	0.0000	0.0001	0.0019	0.0005	0.0000	0.0015	0.0127	0.2758	0.4080	0.3516	
Smokeless Tobacco	0.7587	0.5314	0.0032	0.1019	0.4586	0.1528	0.8811	0.3052	0.4555	0.0778	
Cigars	0.8026	0.8792	0.7363	0.5277	0.6879	0.6106	0.7434	0.6080	0.7038	0.1815	
Pipe Tobacco	0.3519	0.0467	0.1266	0.8237	0.7092	0.4623	0.0627	0.0610	0.1542	0.0683	
<b>ALCOHOL</b>	0.0204	0.0000	0.0002	0.4504	0.0058	0.0324	0.2413	0.2899	0.2987	0.4684	
Binge Alcohol Use <sup>2</sup>	0.2535	0.0332	0.0694	0.0426	0.2231	0.8554	0.8007	0.4682	0.8256	0.3716	
Heavy Alcohol Use <sup>2</sup>	0.4346	0.4331	0.8949	0.0790	0.6101	0.9162	0.6089	0.7963	0.8458	0.1315	
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0064	0.0163	0.0081	0.0001	0.0214	0.1136	0.6319	0.8174	0.1488	
Cigarettes	0.0000	0.0016	0.0041	0.0084	0.0002	0.0061	0.0758	0.8429	0.5701	0.2703	
Smokeless Tobacco	0.5991	0.2584	0.0031	0.0662	0.6330	0.1524	0.7890	0.4952	0.5081	0.0596	
Cigars	0.1573	0.8276	0.2072	0.1200	0.5087	0.9097	0.6250	0.7695	0.7115	0.2659	
Pipe Tobacco	0.3358	0.1373	0.2689	0.9212	0.4517	0.8441	0.0696	0.1657	0.2686	0.1669	
<b>ALCOHOL</b>	0.5095	0.7660	0.9456	0.1277	0.9893	0.6975	0.2075	0.6018	0.7145	0.8363	
Binge Alcohol Use <sup>2</sup>	0.9820	0.4962	0.7560	0.2046	0.8902	0.3971	0.2048	0.4725	0.9868	0.4183	
Heavy Alcohol Use <sup>2</sup>	0.5561	0.7517	0.8531	0.4517	0.6063	0.5003	0.3798	0.7887	0.9348	0.2610	
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0025	0.0035	0.1707	0.0244	0.0425	0.1051	0.0925	0.1633	0.8630	0.7951	
Cigarettes	0.0009	0.0042	0.0862	0.0104	0.0182	0.0517	0.0488	0.0627	0.5303	0.8203	
Smokeless Tobacco	0.5318	0.2927	0.6468	0.9263	0.3378	0.7004	0.6204	0.1287	0.5572	0.9180	
Cigars	0.0025	0.3613	0.0260	0.0150	0.5377	0.0719	0.0115	0.4168	0.8148	0.3948	
Pipe Tobacco	0.8809	0.0614	0.1114	0.2894	0.2479	0.0234	0.5253	0.0566	0.2226	0.1038	
<b>ALCOHOL</b>	0.0000	0.0000	0.0000	0.0069	0.0000	0.0057	0.0026	0.0386	0.0459	0.1938	
Binge Alcohol Use <sup>2</sup>	0.0440	0.0039	0.0043	0.0719	0.0095	0.0909	0.1257	0.8425	0.5838	0.6652	
Heavy Alcohol Use <sup>2</sup>	0.0051	0.2809	0.5289	0.0111	0.0220	0.0901	0.5343	0.9816	0.7004	0.2530	

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.33C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	829	784	793	839	838	838	872	822	840	845	874
Cigarettes	790	774	765	772	799	803	809	790	772	814	832
Smokeless Tobacco	276	283	243	264	281	280	311	288	310	275	298
Cigars	346	333	323	351	348	331	363	358	358	369	387
Pipe Tobacco	163	130	145	145	165	169	146	149	147	143	183
<b>ALCOHOL</b>	972	909	950	969	954	1,011	977	945	974	1,006	1,004
Binge Alcohol Use <sup>2</sup>	730	675	699	701	742	766	737	784	748	736	792
Heavy Alcohol Use <sup>2</sup>	389	376	383	378	425	426	414	404	421	423	448
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	565	561	568	576	581	594	630	597	616	580	613
Cigarettes	521	537	532	514	542	573	565	564	529	535	552
Smokeless Tobacco	260	255	233	239	271	249	295	281	299	261	280
Cigars	322	306	291	324	316	306	332	332	318	337	349
Pipe Tobacco	136	127	138	141	159	168	131	143	136	135	169
<b>ALCOHOL</b>	643	590	616	616	599	658	626	644	666	678	666
Binge Alcohol Use <sup>2</sup>	540	517	511	529	561	585	588	606	573	568	621
Heavy Alcohol Use <sup>2</sup>	346	320	318	322	352	373	358	349	352	355	382
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	509	498	494	541	506	505	553	501	532	539	552
Cigarettes	497	481	488	537	497	487	548	494	524	529	538
Smokeless Tobacco	99	126	64	87	75	112	74	51	76	82	103
Cigars	110	121	119	109	128	112	113	127	150	140	145
Pipe Tobacco	90	35	41	40	45	28	54	35	53	45	60
<b>ALCOHOL</b>	640	593	662	635	641	673	667	679	634	677	682
Binge Alcohol Use <sup>2</sup>	411	375	389	401	395	392	399	442	444	443	426
Heavy Alcohol Use <sup>2</sup>	175	170	173	162	168	169	192	192	201	207	202

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.33D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2012**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.39	0.37	0.37	0.39	0.38	0.38	0.39	0.36	0.37	0.36	0.37
Cigarettes	0.38	0.36	0.36	0.35	0.36	0.36	0.36	0.35	0.34	0.35	0.35
Smokeless Tobacco	0.13	0.13	0.11	0.12	0.13	0.13	0.14	0.13	0.14	0.12	0.13
Cigars	0.16	0.16	0.15	0.16	0.16	0.15	0.16	0.16	0.16	0.16	0.16
Pipe Tobacco	0.08	0.06	0.07	0.07	0.07	0.08	0.06	0.07	0.06	0.06	0.08
<b>ALCOHOL</b>	0.46	0.43	0.44	0.44	0.43	0.45	0.43	0.42	0.42	0.43	0.43
Binge Alcohol Use <sup>2</sup>	0.35	0.32	0.32	0.32	0.34	0.34	0.33	0.34	0.33	0.32	0.34
Heavy Alcohol Use <sup>2</sup>	0.19	0.18	0.18	0.17	0.19	0.19	0.18	0.18	0.18	0.18	0.19
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.56	0.55	0.55	0.55	0.55	0.55	0.58	0.54	0.56	0.52	0.54
Cigarettes	0.52	0.53	0.51	0.49	0.51	0.53	0.52	0.51	0.48	0.48	0.49
Smokeless Tobacco	0.26	0.25	0.22	0.23	0.25	0.23	0.27	0.26	0.27	0.23	0.25
Cigars	0.32	0.30	0.28	0.31	0.30	0.29	0.31	0.30	0.29	0.30	0.31
Pipe Tobacco	0.13	0.12	0.13	0.13	0.15	0.16	0.12	0.13	0.12	0.12	0.15
<b>ALCOHOL</b>	0.64	0.58	0.59	0.59	0.56	0.61	0.58	0.59	0.60	0.61	0.59
Binge Alcohol Use <sup>2</sup>	0.53	0.51	0.49	0.50	0.53	0.55	0.54	0.55	0.52	0.51	0.55
Heavy Alcohol Use <sup>2</sup>	0.34	0.31	0.31	0.31	0.33	0.35	0.33	0.32	0.32	0.32	0.34
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.47	0.45	0.44	0.48	0.44	0.44	0.47	0.43	0.45	0.45	0.45
Cigarettes	0.45	0.44	0.44	0.48	0.44	0.42	0.47	0.42	0.44	0.44	0.44
Smokeless Tobacco	0.09	0.11	0.06	0.08	0.07	0.10	0.06	0.04	0.06	0.07	0.08
Cigars	0.10	0.11	0.11	0.10	0.11	0.10	0.10	0.11	0.13	0.12	0.12
Pipe Tobacco	0.08	0.03	0.04	0.04	0.04	0.02	0.05	0.03	0.04	0.04	0.05
<b>ALCOHOL</b>	0.59	0.54	0.59	0.56	0.56	0.58	0.57	0.58	0.54	0.56	0.56
Binge Alcohol Use <sup>2</sup>	0.38	0.34	0.35	0.36	0.35	0.34	0.34	0.38	0.37	0.37	0.35
Heavy Alcohol Use <sup>2</sup>	0.16	0.15	0.16	0.14	0.15	0.15	0.16	0.16	0.17	0.17	0.17

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.

**Table 7.33P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: P Values from Tests of Differences of Percentages, 2012 Versus 2002-2011**

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>TOTAL</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0024	0.0997	0.2596	0.4449	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0211	0.1148	0.7093	
Smokeless Tobacco	0.3975	0.2073	0.0009	0.0597	0.3507	0.1036	1.0000	0.6268	1.0000	0.0797	
Cigars	0.6743	0.7260	0.0833	0.1455	0.1909	0.7802	0.9133	0.9790	0.9517	0.2381	
Pipe Tobacco	0.0590	0.0012	0.0220	0.4737	0.7593	0.1354	0.0166	0.0652	0.1799	0.0936	
<b>ALCOHOL</b>	0.0280	0.0001	0.0005	0.5816	0.0271	0.0680	0.3471	0.5313	0.4859	0.5585	
Binge Alcohol Use <sup>2</sup>	0.5227	0.2046	0.2865	0.2868	0.8074	0.5730	0.4479	0.1545	0.8664	0.4576	
Heavy Alcohol Use <sup>2</sup>	0.6017	0.4807	0.1932	0.9358	0.2450	0.3029	0.1284	0.3763	0.7203	0.0811	
<b>MALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0002	0.0006	0.0004	0.0000	0.0039	0.0523	0.8357	0.4865	0.2158	
Cigarettes	0.0000	0.0000	0.0000	0.0001	0.0000	0.0005	0.0206	0.5018	0.3490	0.4917	
Smokeless Tobacco	0.3260	0.0793	0.0010	0.0377	0.5047	0.0999	0.9019	0.7979	1.0000	0.0589	
Cigars	0.1297	0.5408	0.0126	0.0239	0.1376	0.3641	0.4010	0.8766	0.8883	0.2366	
Pipe Tobacco	0.0916	0.0149	0.1255	0.8403	0.7285	0.4708	0.0238	0.2076	0.2021	0.1628	
<b>ALCOHOL</b>	0.2872	0.2909	0.6493	0.0333	0.5492	0.9579	0.1421	0.2885	0.3678	0.7870	
Binge Alcohol Use <sup>2</sup>	0.3629	0.5841	0.5014	0.9510	0.2784	0.1054	0.0862	0.1334	0.5676	0.5118	
Heavy Alcohol Use <sup>2</sup>	0.0786	0.3074	0.1246	0.4113	0.0816	0.1522	0.0990	0.3994	0.7649	0.1577	
<b>FEMALE</b>											
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0018	0.0001	0.0004	0.0108	0.0083	0.0236	0.3788	0.8402	
Cigarettes	0.0000	0.0000	0.0003	0.0000	0.0000	0.0026	0.0027	0.0052	0.1700	0.8734	
Smokeless Tobacco	0.7153	0.3444	0.4871	0.9854	0.2835	0.7028	0.5601	0.2672	0.7483	0.8702	
Cigars	0.0082	0.6157	0.1608	0.0738	0.8442	0.0993	0.0368	0.6480	0.7827	0.8141	
Pipe Tobacco	0.3917	0.0026	0.0056	0.0863	0.0334	0.0012	0.3215	0.0261	0.5852	0.2623	
<b>ALCOHOL</b>	0.0000	0.0000	0.0000	0.0028	0.0001	0.0060	0.0043	0.0476	0.0407	0.2476	
Binge Alcohol Use <sup>2</sup>	0.0201	0.0023	0.0041	0.0462	0.0375	0.1526	0.2111	0.7025	0.5092	0.7090	
Heavy Alcohol Use <sup>2</sup>	0.0316	0.7473	0.9873	0.0777	0.4714	0.5555	0.8853	0.7866	0.9536	0.2774	

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2012.