

**Table 1.74A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	9,208 <sup>a</sup>	8,214	885 <sup>a</sup>	625	378 <sup>a</sup>	228
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,643	1,388	213	138	101	63
New England	521	353	83	46	57	21
Middle Atlantic	1,122	1,035	130	93	44	42
Midwest	1,757	1,698	185	164	79	76
East North Central	1,126	1,247	160	137	64	66
West North Central	631 <sup>a</sup>	451	25	27	15	9
South	3,329	2,870	342	245	165	76
South Atlantic	1,975	1,578	153	136	62	43
East South Central	542	473	79	55	39	9
West South Central	813	819	110	53	64	23
West	2,479	2,258	144	78	33	14
Mountain	844	692	51	22	9	7
Pacific	1,635	1,566	93	57	24	7
<b>COUNTY TYPE</b>						
Large Metro	4,619	4,231	469	302	202 <sup>a</sup>	99
Small Metro	3,145	2,702	349	245	155	100
250K – 1 Mil. Pop.	2,095	1,863	249	136	135	67
< 250K Pop.	1,050	840	99	109	20	33
Nonmetro	1,444	1,280	67	79	21	30
Urbanized	671	535	39	45	14	19
Less Urbanized	709	654	26	26	7	3
Completely Rural	64	91	1	7	*	7

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.74B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	3.6 <sup>a</sup>	3.2	0.3 <sup>a</sup>	0.2	0.1 <sup>a</sup>	0.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	3.5	3.0	0.5	0.3	0.2	0.1
New England	4.3	2.9	0.7	0.4	0.5	0.2
Middle Atlantic	3.3	3.0	0.4	0.3	0.1	0.1
Midwest	3.2	3.0	0.3	0.3	0.1	0.1
East North Central	2.9	3.2	0.4	0.4	0.2	0.2
West North Central	3.8 <sup>a</sup>	2.7	0.1	0.2	0.1	0.1
South	3.6	3.0	0.4	0.3	0.2	0.1
South Atlantic	4.0	3.1	0.3	0.3	0.1	0.1
East South Central	3.6	3.1	0.5	0.4	0.3	0.1
West South Central	2.8	2.8	0.4	0.2	0.2	0.1
West	4.2	3.8	0.2	0.1	0.1	0.0
Mountain	4.7	3.8	0.3	0.1	0.0	0.0
Pacific	4.0	3.8	0.2	0.1	0.1	0.0
<b>COUNTY TYPE</b>						
Large Metro	3.4	3.1	0.3 <sup>a</sup>	0.2	0.1 <sup>a</sup>	0.1
Small Metro	4.1 <sup>a</sup>	3.4	0.4	0.3	0.2	0.1
250K – 1 Mil. Pop.	4.1	3.4	0.5 <sup>a</sup>	0.2	0.3	0.1
< 250K Pop.	4.0	3.2	0.4	0.4	0.1	0.1
Nonmetro	3.6	3.2	0.2	0.2	0.1	0.1
Urbanized	3.9	3.2	0.2	0.3	0.1	0.1
Less Urbanized	3.6	3.4	0.1	0.1	0.0	0.0
Completely Rural	1.7	2.2	0.0	0.2	*	0.2

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.