

Table 1.76A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	21,778	20,523	2,048	1,861	690	624
GEOGRAPHIC DIVISION						
Northeast	4,088	4,323	451	351	147	110
New England	1,121	1,174	104	104	36	33
Middle Atlantic	2,968	3,149	347	247	111	77
Midwest	4,425	4,181	397	351	117	116
East North Central	3,243	3,034	289 ^a	208	101 ^a	52
West North Central	1,182	1,147	108	143	17	64
South	7,025	6,411	667	533	284	208
South Atlantic	3,934	3,356	379	294	207	121
East South Central	1,010	1,039	111	71	30	27
West South Central	2,082	2,016	177	168	47	60
West	6,239	5,608	533	626	141	190
Mountain	1,931	1,710	144	212	56	54
Pacific	4,308	3,898	389	415	85	135
COUNTY TYPE						
Large Metro	12,091	11,277	1,199	974	439	290
Small Metro	6,882	6,706	589	683	194	270
250K – 1 Mil. Pop.	4,725	4,805	396	495	138	212
< 250K Pop.	2,157	1,901	192	188	56	58
Nonmetro	2,805	2,541	260 ^a	203	58	64
Urbanized	1,349	1,190	107	100	24	39
Less Urbanized	1,255	1,145	123	89	27	23
Completely Rural	201	206	30	15	7	2

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.76B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	8.6 ^a	8.0	0.8	0.7	0.3	0.2
GEOGRAPHIC DIVISION						
Northeast	8.8	9.2	1.0	0.7	0.3	0.2
New England	9.1	9.5	0.9	0.8	0.3	0.3
Middle Atlantic	8.7	9.1	1.0	0.7	0.3	0.2
Midwest	8.0	7.5	0.7	0.6	0.2	0.2
East North Central	8.4	7.8	0.8 ^a	0.5	0.3 ^a	0.1
West North Central	7.0	6.7	0.6	0.8	0.1	0.4
South	7.6	6.7	0.7	0.6	0.3	0.2
South Atlantic	8.0 ^a	6.7	0.8	0.6	0.4	0.2
East South Central	6.7	6.8	0.7 ^a	0.5	0.2	0.2
West South Central	7.2	6.8	0.6	0.6	0.2	0.2
West	10.6	9.4	0.9	1.0	0.2	0.3
Mountain	10.7	9.4	0.8	1.2	0.3	0.3
Pacific	10.6	9.4	1.0	1.0	0.2	0.3
COUNTY TYPE						
Large Metro	8.9	8.2	0.9	0.7	0.3	0.2
Small Metro	8.9	8.3	0.8	0.8	0.3	0.3
250K – 1 Mil. Pop.	9.2	8.8	0.8	0.9	0.3	0.4
< 250K Pop.	8.3	7.3	0.7	0.7	0.2	0.2
Nonmetro	6.9	6.3	0.6	0.5	0.1	0.2
Urbanized	7.9	7.1	0.6	0.6	0.1	0.2
Less Urbanized	6.4	6.0	0.6	0.5	0.1	0.1
Completely Rural	5.4	4.9	0.8	0.3	0.2	0.0

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.77A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	2,015	1,877	885	820	262	223
GEOGRAPHIC DIVISION						
Northeast	344	303	156	151	44	38
New England	100 ^a	73	46	38	18	8
Middle Atlantic	244	230	110	113	27	29
Midwest	447	406	191	162	58	42
East North Central	320 ^a	267	134 ^a	100	49 ^a	26
West North Central	127	139	56	63	9	16
South	657	632	305	251	102	77
South Atlantic	339	296	152	112	48	33
East South Central	139 ^a	102	76 ^a	45	23	19
West South Central	179 ^a	234	77	94	32	25
West	567	536	234	256	58	66
Mountain	178	148	65	71	14	11
Pacific	389	388	169	184	44	55
COUNTY TYPE						
Large Metro	1,070	1,000	465	442	142	98
Small Metro	650	568	263	253	87	85
250K – 1 Mil. Pop.	464	398	179	196	56	68
< 250K Pop.	186	170	84	57	31	17
Nonmetro	295	310	157	125	33	39
Urbanized	113	120	63	57	10	18
Less Urbanized	151	161	76	59	18	19
Completely Rural	30	28	19	8	5	2

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.77B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	8.3 ^a	7.5	3.6	3.3	1.1	0.9
GEOGRAPHIC DIVISION						
Northeast	8.1	7.1	3.7	3.5	1.0	0.9
New England	9.1 ^a	6.6	4.2	3.4	1.6	0.7
Middle Atlantic	7.8	7.3	3.5	3.6	0.9	0.9
Midwest	8.4	7.5	3.6	3.0	1.1	0.8
East North Central	8.6 ^a	7.0	3.6 ^a	2.6	1.3 ^a	0.7
West North Central	8.0	8.5	3.5	3.8	0.6	1.0
South	7.3	6.8	3.4 ^a	2.7	1.1	0.8
South Atlantic	7.5	6.4	3.4	2.4	1.1	0.7
East South Central	9.6 ^a	6.9	5.3 ^a	3.1	1.6	1.3
West South Central	5.9	7.4	2.6	3.0	1.0	0.8
West	9.7	8.9	4.0	4.3	1.0	1.1
Mountain	10.0	8.0	3.6	3.9	0.8	0.6
Pacific	9.6	9.3	4.2	4.4	1.1	1.3
COUNTY TYPE						
Large Metro	8.2	7.5	3.6	3.3	1.1	0.7
Small Metro	8.6 ^a	7.2	3.5	3.2	1.2	1.1
250K – 1 Mil. Pop.	9.2 ^b	7.4	3.5	3.6	1.1	1.3
< 250K Pop.	7.5	6.9	3.4	2.3	1.2	0.7
Nonmetro	7.8	8.1	4.2	3.3	0.9	1.0
Urbanized	7.4	7.1	4.1	3.4	0.6	1.1
Less Urbanized	7.8	9.1	3.9	3.3	1.0	1.1
Completely Rural	9.5	8.0	5.9 ^a	2.4	1.5	0.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.