

Table 1.78A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	51,832	51,243	16,051 ^a	14,657	6,957 ^a	6,119
GEOGRAPHIC DIVISION						
Northeast	8,572	8,532	2,697	2,466	1,298	1,048
New England	2,346	2,350	758	691	411	278
Middle Atlantic	6,227	6,182	1,938	1,775	887	770
Midwest	10,678	10,685	3,268	3,144	1,383	1,393
East North Central	7,674	7,529	2,423	2,272	1,080	1,039
West North Central	3,004	3,156	845	871	303	354
South	18,348	18,419	5,602	5,334	2,524	2,145
South Atlantic	9,698	9,515	2,951	2,652	1,378 ^a	975
East South Central	3,241	3,095	874	927	404	422
West South Central	5,410	5,810	1,777	1,754	742	749
West	14,234	13,606	4,484 ^a	3,713	1,751	1,534
Mountain	4,523	4,405	1,460	1,239	603	469
Pacific	9,710	9,202	3,024	2,473	1,149	1,065
COUNTY TYPE						
Large Metro	28,305	27,040	8,711 ^a	7,672	3,671 ^a	3,036
Small Metro	16,190	16,833	5,100	4,844	2,214	2,122
250K – 1 Mil. Pop.	10,740	11,692	3,480	3,207	1,517	1,398
< 250K Pop.	5,450	5,142	1,620	1,637	697	724
Nonmetro	7,337	7,369	2,239	2,141	1,072	961
Urbanized	3,415	3,398	1,061	1,101	533	485
Less Urbanized	3,409	3,196	1,061	873	498	405
Completely Rural	513 ^a	775	117	168	41	72

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.78B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	20.4	19.9	6.3 ^b	5.7	2.7 ^b	2.4
GEOGRAPHIC DIVISION						
Northeast	18.4	18.2	5.8	5.3	2.8	2.2
New England	19.2	19.1	6.2	5.6	3.4 ^a	2.3
Middle Atlantic	18.2	17.9	5.7	5.1	2.6	2.2
Midwest	19.3	19.2	5.9	5.6	2.5	2.5
East North Central	19.9	19.5	6.3	5.9	2.8	2.7
West North Central	17.9	18.6	5.0	5.1	1.8	2.1
South	19.7	19.4	6.0	5.6	2.7 ^a	2.3
South Atlantic	19.8	19.0	6.0	5.3	2.8 ^b	1.9
East South Central	21.5	20.2	5.8	6.1	2.7	2.8
West South Central	18.7	19.6	6.2	5.9	2.6	2.5
West	24.2	22.7	7.6 ^b	6.2	3.0	2.6
Mountain	25.1	24.3	8.1	6.8	3.3	2.6
Pacific	23.8	22.1	7.4 ^a	5.9	2.8	2.6
COUNTY TYPE						
Large Metro	20.9	19.8	6.4 ^a	5.6	2.7 ^a	2.2
Small Metro	20.9	20.9	6.6	6.0	2.9	2.6
250K – 1 Mil. Pop.	20.9	21.5	6.8 ^a	5.9	2.9	2.6
< 250K Pop.	21.0	19.7	6.2	6.3	2.7	2.8
Nonmetro	18.1	18.3	5.5	5.3	2.6	2.4
Urbanized	20.0	20.2	6.2	6.6	3.1	2.9
Less Urbanized	17.3	16.7	5.4	4.6	2.5	2.1
Completely Rural	13.8	18.5	3.1	4.0	1.1	1.7

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.79A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	2,591	2,522	1,812	1,749	721	688
GEOGRAPHIC DIVISION						
Northeast	336	375	247	264	94	99
New England	88	89	65	60	23	21
Middle Atlantic	248	286	182	204	71	78
Midwest	564	539	414	384	159	167
East North Central	390	368	286	268	114	119
West North Central	174	172	128	116	45	48
South	975	941	651	652	266	269
South Atlantic	488 ^a	401	326	278	120	112
East South Central	161	173	103	119	48	58
West South Central	326	367	222	255	98	99
West	715	667	500	449	202	153
Mountain	252	209	189 ^a	139	86	53
Pacific	463	458	311	310	116	100
COUNTY TYPE						
Large Metro	1,313	1,284	928	886	360	315
Small Metro	853	803	594	577	248	255
250K – 1 Mil. Pop.	577	546	405	396	166	171
< 250K Pop.	276	257	189	181	82	84
Nonmetro	425	435	291	286	113	118
Urbanized	179	184	133	123	53	53
Less Urbanized	219	214	143	137	51	54
Completely Rural	27	38	16	26	9	11

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.79B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	10.6	10.1	7.4	7.0	3.0	2.8
GEOGRAPHIC DIVISION						
Northeast	7.9	8.8	5.8	6.2	2.2	2.3
New England	8.1	8.0	6.0	5.4	2.1	1.9
Middle Atlantic	7.9	9.0	5.8	6.4	2.3	2.4
Midwest	10.6	9.9	7.8	7.0	3.0	3.1
East North Central	10.4	9.7	7.6	7.0	3.0	3.1
West North Central	11.0	10.5	8.0	7.1	2.8	3.0
South	10.9	10.2	7.3	7.0	3.0	2.9
South Atlantic	10.9 ^a	8.7	7.3	6.0	2.7	2.4
East South Central	11.1	11.7	7.1	8.0	3.3	3.9
West South Central	10.8	11.6	7.4	8.1	3.2	3.1
West	12.3	11.1	8.6	7.5	3.5	2.6
Mountain	14.1	11.3	10.6 ^a	7.5	4.8	2.9
Pacific	11.5	11.0	7.7	7.5	2.9	2.4
COUNTY TYPE						
Large Metro	10.1	9.6	7.1	6.7	2.8	2.4
Small Metro	11.3	10.2	7.9	7.4	3.3	3.3
250K – 1 Mil. Pop.	11.5	10.2	8.0	7.4	3.3	3.2
< 250K Pop.	11.1	10.4	7.6	7.3	3.3	3.4
Nonmetro	11.2	11.4	7.7	7.5	3.0	3.1
Urbanized	11.7	10.9	8.7	7.3	3.5	3.1
Less Urbanized	11.3	12.1	7.4	7.7	2.6	3.0
Completely Rural	8.4	10.6	5.0	7.4	2.9	3.2

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.80A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	9,832 ^a	9,373	4,876 ^b	4,366	1,997 ^b	1,711
GEOGRAPHIC DIVISION						
Northeast	1,666 ^b	1,452	893 ^a	752	351	285
New England	517 ^a	435	263	227	116	89
Middle Atlantic	1,149	1,017	629 ^a	525	235	196
Midwest	2,107	2,031	1,107 ^a	987	472	401
East North Central	1,470	1,411	781	695	349	292
West North Central	637	619	326	293	123	109
South	3,664	3,465	1,763 ^a	1,554	712	614
South Atlantic	1,802	1,731	880	761	345	290
East South Central	605	608	277	287	126	131
West South Central	1,257	1,126	607	506	241	192
West	2,395	2,425	1,113	1,072	462	411
Mountain	749	818	347	390	154	129
Pacific	1,646	1,607	766	682	309	283
COUNTY TYPE						
Large Metro	5,049	4,754	2,520 ^a	2,221	983	886
Small Metro	3,411	3,273	1,703	1,526	736 ^a	598
250K – 1 Mil. Pop.	2,193	2,112	1,082	975	486 ^b	352
< 250K Pop.	1,218	1,161	621	551	250	247
Nonmetro	1,372	1,346	652	619	277	227
Urbanized	670	698	322	324	140	111
Less Urbanized	606	546	288	258	130	104
Completely Rural	96	102	43	37	8	12

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.80B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	28.9 ^b	27.3	14.3 ^b	12.7	5.9 ^b	5.0
GEOGRAPHIC DIVISION						
Northeast	27.6 ^b	23.7	14.8 ^b	12.3	5.8	4.7
New England	31.8 ^a	27.1	16.2	14.2	7.1	5.5
Middle Atlantic	26.0 ^a	22.5	14.2 ^a	11.6	5.3	4.3
Midwest	28.1	27.7	14.8	13.5	6.3	5.5
East North Central	28.6	27.9	15.2	13.7	6.8	5.8
West North Central	27.0	27.2	13.8	12.9	5.2	4.8
South	29.5 ^a	27.5	14.2 ^a	12.3	5.7	4.9
South Atlantic	28.5	26.9	13.9	11.8	5.5	4.5
East South Central	30.4	30.3	13.9	14.3	6.4	6.5
West South Central	30.6 ^a	27.1	14.7 ^a	12.2	5.9	4.6
West	29.5	29.5	13.7	13.0	5.7	5.0
Mountain	30.0	33.1	13.9	15.8	6.1	5.2
Pacific	29.3	27.9	13.6	11.8	5.5	4.9
COUNTY TYPE						
Large Metro	27.8 ^a	25.8	13.9 ^b	12.1	5.4	4.8
Small Metro	30.7	29.4	15.3 ^a	13.7	6.6 ^a	5.4
250K – 1 Mil. Pop.	31.2	29.8	15.4	13.8	6.9 ^b	5.0
< 250K Pop.	29.7	28.6	15.2	13.6	6.1	6.1
Nonmetro	28.6	28.4	13.6	13.0	5.8	4.8
Urbanized	31.7	31.6	15.2	14.7	6.6	5.0
Less Urbanized	26.0	26.1	12.4	12.3	5.6	5.0
Completely Rural	27.6	23.0	12.2	8.3	2.2	2.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.81A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	39,409	39,348	9,362	8,542	4,239	3,720
GEOGRAPHIC DIVISION						
Northeast	6,570	6,704	1,557	1,450	853	664
New England	1,741	1,826	430	403	272	168
Middle Atlantic	4,829	4,878	1,128	1,047	581	496
Midwest	8,007	8,115	1,746	1,772	753	825
East North Central	5,815	5,750	1,356	1,309	618	629
West North Central	2,192	2,365	391	463	135	196
South	13,708	14,014	3,188	3,128	1,546	1,262
South Atlantic	7,407	7,384	1,746	1,613	913 ^a	573
East South Central	2,475	2,313	494	522	230	233
West South Central	3,826	4,317	948	993	403	457
West	11,124	10,514	2,871 ^a	2,192	1,087	969
Mountain	3,522	3,378	924	711	363	287
Pacific	7,602	7,136	1,947	1,481	724	682
COUNTY TYPE						
Large Metro	21,943	21,002	5,263	4,565	2,328	1,835
Small Metro	11,926	12,758	2,804	2,741	1,230	1,268
250K – 1 Mil. Pop.	7,970 ^a	9,034	1,993	1,836	865	875
< 250K Pop.	3,956	3,724	810	905	365	393
Nonmetro	5,540	5,588	1,296	1,236	681	616
Urbanized	2,566	2,516	607	654	340	321
Less Urbanized	2,584	2,436	630	478	317	247
Completely Rural	390 ^a	635	59	105	24	49

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.81B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	20.2	19.8	4.8	4.3	2.2	1.9
GEOGRAPHIC DIVISION						
Northeast	18.1	18.4	4.3	4.0	2.4	1.8
New England	18.3	19.0	4.5	4.2	2.9	1.7
Middle Atlantic	18.1	18.1	4.2	3.9	2.2	1.8
Midwest	18.8	18.9	4.1	4.1	1.8	1.9
East North Central	19.6	19.3	4.6	4.4	2.1	2.1
West North Central	17.0	18.1	3.0	3.5	1.0	1.5
South	19.1	19.1	4.5	4.3	2.2	1.7
South Atlantic	19.4	18.9	4.6	4.1	2.4 ^a	1.5
East South Central	21.2	19.6	4.2	4.4	2.0	2.0
West South Central	17.6	19.3	4.4	4.4	1.9	2.0
West	24.8	23.1	6.4 ^a	4.8	2.4	2.1
Mountain	25.6	24.4	6.7	5.1	2.6	2.1
Pacific	24.4	22.5	6.3	4.7	2.3	2.1
COUNTY TYPE						
Large Metro	21.0	20.0	5.0	4.3	2.2 ^a	1.7
Small Metro	20.3	20.7	4.8	4.4	2.1	2.1
250K – 1 Mil. Pop.	20.2	21.5	5.1	4.4	2.2	2.1
< 250K Pop.	20.4	19.0	4.2	4.6	1.9	2.0
Nonmetro	17.4	17.7	4.1	3.9	2.1	1.9
Urbanized	19.1	19.5	4.5	5.1	2.5	2.5
Less Urbanized	16.8	15.9	4.1	3.1	2.1	1.6
Completely Rural	12.7 ^a	18.7	1.9	3.1	0.8	1.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.82A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	49,241	48,721	14,238 ^a	12,908	6,236 ^a	5,431
GEOGRAPHIC DIVISION						
Northeast	8,236	8,157	2,450	2,202	1,204	949
New England	2,258	2,261	693	631	388	257
Middle Atlantic	5,978	5,896	1,757	1,571	816	692
Midwest	10,114	10,146	2,853	2,760	1,224	1,226
East North Central	7,284	7,161	2,137	2,004	967	921
West North Central	2,829	2,985	717	756	258	305
South	17,373	17,479	4,951	4,682	2,258	1,876
South Atlantic	9,209	9,114	2,625	2,374	1,258 ^a	863
East South Central	3,080	2,921	771	809	356	364
West South Central	5,084	5,443	1,555	1,499	644	649
West	13,519	12,939	3,984 ^a	3,264	1,550	1,380
Mountain	4,271	4,196	1,271	1,101	517	415
Pacific	9,248	8,743	2,713	2,163	1,033	965
COUNTY TYPE						
Large Metro	26,992	25,756	7,784 ^a	6,786	3,311 ^a	2,721
Small Metro	15,337	16,031	4,507	4,267	1,966	1,866
250K – 1 Mil. Pop.	10,164	11,146	3,075	2,811	1,351	1,227
< 250K Pop.	5,174	4,885	1,432	1,456	615	640
Nonmetro	6,912	6,934	1,948	1,855	958	843
Urbanized	3,236	3,214	928	978	480	432
Less Urbanized	3,190	2,982	918	736	447	351
Completely Rural	486 ^a	737	102	142	32	60

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.82B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	21.5	20.9	6.2 ^b	5.5	2.7 ^a	2.3
GEOGRAPHIC DIVISION						
Northeast	19.5	19.1	5.8	5.2	2.8	2.2
New England	20.2	20.2	6.2	5.6	3.5	2.3
Middle Atlantic	19.2	18.8	5.6	5.0	2.6	2.2
Midwest	20.2	20.2	5.7	5.5	2.4	2.4
East North Central	20.9	20.5	6.1	5.7	2.8	2.6
West North Central	18.6	19.4	4.7	4.9	1.7	2.0
South	20.7	20.3	5.9	5.4	2.7 ^a	2.2
South Atlantic	20.7	20.0	5.9	5.2	2.8 ^b	1.9
East South Central	22.6	21.1	5.7	5.8	2.6	2.6
West South Central	19.7	20.5	6.0	5.6	2.5	2.4
West	25.5	24.0	7.5 ^a	6.1	2.9	2.6
Mountain	26.3	25.7	7.8	6.7	3.2	2.5
Pacific	25.2	23.3	7.4 ^a	5.8	2.8	2.6
COUNTY TYPE						
Large Metro	22.0	20.9	6.3 ^a	5.5	2.7 ^a	2.2
Small Metro	21.9	22.0	6.4	5.9	2.8	2.6
250K – 1 Mil. Pop.	21.9	22.7	6.6	5.7	2.9	2.5
< 250K Pop.	22.0	20.7	6.1	6.2	2.6	2.7
Nonmetro	18.8	19.1	5.3	5.1	2.6	2.3
Urbanized	20.8	21.3	6.0	6.5	3.1	2.9
Less Urbanized	18.0	17.1	5.2	4.2	2.5	2.0
Completely Rural	14.3	19.2	3.0	3.7	0.9	1.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.