

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	1,023	1,036	646	603	488	460
12	27	28	23	20	16	15
13	37	35	31	27	22	19
14	38	40	32	34	24	24
15	51	50	44	46	31	32
16	54	57	47	49	36	37
17	62	58	57	49	41	38
18	88	100	78	85	58	65
19	86	96	73	82	57	61
20	92	110	77	91	60	68
21	91	88	73	67	54	53
22	83	103	66	80	52	57
23	85	84	61	63	49	48
24	83	88	61	61	47	46
25	87	94	62	59	45	50
26-29	345	345	221	223	169	173
30-34	352	375	209	207	172	154
35-39	349	367	169	183	121	138
40-44	377	374	175	161	130	119
45-49	419	384	183	167	136	124
50-54	568	594	240	220	196	173
55-59	538	454	218	191	131	153
60-64	365	389	134	132	96	87
65 or Older	370	343	135	109	97	85

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.40	0.40	0.25	0.23	0.19	0.18
12	0.69	0.72	0.59	0.53	0.41	0.39
13	0.81	0.75	0.69	0.60	0.51	0.43
14	0.87	0.84	0.76	0.74	0.60	0.54
15	0.98	0.96	0.91	0.93	0.70	0.71
16	1.07	1.09	0.95	0.98	0.79	0.76
17	1.07	1.08	1.06	1.00	0.84	0.82
18	1.16	1.28	1.16	1.19	1.03	1.04
19	1.25	1.36	1.24	1.34	1.09	1.15
20	1.31	1.32	1.33	1.33	1.15	1.17
21	1.34	1.28	1.26	1.18	1.09	1.03
22	1.19	1.28	1.19	1.31	1.05	1.11
23	1.30	1.27	1.20	1.30	1.07	1.07
24	1.30	1.27	1.22	1.24	1.03	1.02
25	1.31	1.26	1.27	1.18	1.05	1.10
26-29	1.21	1.27	1.10	1.09	0.92	0.91
30-34	1.12	1.15	0.93	0.94	0.80	0.75
35-39	1.18	1.21	0.80	0.86	0.60	0.68
40-44	1.15	1.22	0.81	0.75	0.60	0.56
45-49	1.13	1.15	0.76	0.73	0.59	0.55
50-54	1.64	1.58	1.06	0.91	0.87	0.73
55-59	1.80	1.62	1.06	0.94	0.66	0.77
60-64	1.86	1.62	0.79	0.70	0.58	0.47
65 or Older	0.95	0.85	0.35	0.27	0.25	0.21

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.6146		0.1638		0.4074	
12	0.3956		0.1139		0.4108	
13	0.1746		0.0490		0.2065	
14	0.9144		0.8442		0.5495	
15	0.4684		0.9203		0.7031	
16	0.1994		0.2416		0.3851	
17	0.8878		0.3193		0.8643	
18	0.2212		0.4325		0.8557	
19	0.2859		0.8862		0.4925	
20	0.3700		0.0386		0.1187	
21	0.2303		0.1160		0.2658	
22	0.4016		0.5428		0.5362	
23	0.7670		0.5554		0.6484	
24	0.9256		0.7536		0.3590	
25	0.9026		0.5097		0.6797	
26-29	0.0517		0.1941		0.9444	
30-34	0.1188		0.8256		0.0941	
35-39	0.7154		0.7663		0.9639	
40-44	0.2388		0.2751		0.5181	
45-49	0.0654		0.1807		0.5186	
50-54	0.6193		0.5592		0.6251	
55-59	0.6185		0.8474		0.0664	
60-64	0.3504		0.2488		0.8939	
65 or Older	0.7127		0.7825		0.7959	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	1,019	1,013	559	514	407	409
12	9	11	9	10	4	6
13	21	23	19	19	11	15
14	28	27	26	26	18	19
15	42	45	39	42	28	28
16	47	51	41	45	33	34
17	57	54	53	47	38	36
18	81	95	72	82	54	61
19	81	91	68	78	54	59
20	88	107	74	87	59	66
21	86	83	66	62	49	51
22	80	100	61	71	49	53
23	82	82	57	59	45	45
24	77	85	52	57	42	42
25	83	90	55	54	40	45
26-29	328	325	199	201	152	158
30-34	337	352	186	176	146	133
35-39	323	348	140	153	102	121
40-44	368	351	144	131	111	99
45-49	405	363	152	133	116	101
50-54	546	568	187	188	159	149
55-59	518	433	181	154	119	133
60-64	352	370	106	119	87	85
65 or Older	348	316	113	89	67	77

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.40	0.39	0.22	0.20	0.16	0.16
12	0.23	0.31	0.22	0.28	0.11	0.15
13	0.48	0.53	0.44	0.45	0.27	0.36
14	0.68	0.61	0.63	0.58	0.46	0.45
15	0.87	0.92	0.82	0.88	0.62	0.62
16	0.96	1.01	0.86	0.91	0.73	0.72
17	1.03	1.05	1.01	0.96	0.80	0.80
18	1.18	1.29	1.15	1.18	0.98	0.99
19	1.27	1.37	1.20	1.31	1.04	1.11
20	1.31	1.36	1.30	1.34	1.13	1.15
21	1.32	1.27	1.24	1.14	1.03	1.00
22	1.23	1.30	1.16	1.24	1.02	1.04
23	1.33	1.32	1.16	1.25	0.98	1.02
24	1.29	1.34	1.10	1.18	0.94	0.93
25	1.36	1.30	1.20	1.14	0.94	1.02
26-29	1.24	1.28	1.03	1.01	0.84	0.85
30-34	1.14	1.17	0.85	0.83	0.69	0.65
35-39	1.15	1.23	0.68	0.73	0.51	0.59
40-44	1.19	1.20	0.67	0.62	0.52	0.47
45-49	1.15	1.11	0.64	0.59	0.51	0.46
50-54	1.67	1.59	0.84	0.79	0.71	0.63
55-59	1.86	1.62	0.90	0.78	0.60	0.68
60-64	1.83	1.58	0.64	0.64	0.52	0.46
65 or Older	0.89	0.79	0.29	0.22	0.17	0.19

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.7538		0.9765		0.4881	
12	0.4971		0.8331		0.3885	
13	0.5699		0.9431		0.1525	
14	0.9135		0.6357		0.6995	
15	0.5677		0.4932		0.7228	
16	0.1022		0.1494		0.2544	
17	0.5834		0.4053		0.7901	
18	0.8357		0.8726		0.4697	
19	0.8459		0.6893		0.9693	
20	0.0674		0.0130		0.0794	
21	0.3136		0.4681		0.8269	
22	0.2287		0.4574		0.5359	
23	0.9850		0.7014		0.7492	
24	0.9190		0.8818		0.3310	
25	0.7256		0.8200		0.2622	
26-29	0.0523		0.5791		0.4432	
30-34	0.1753		0.2723		0.1217	
35-39	0.5082		0.7611		0.5818	
40-44	0.1304		0.6518		0.5085	
45-49	0.0175		0.3301		0.7434	
50-54	0.7251		0.6083		0.9851	
55-59	0.5473		0.6826		0.1208	
60-64	0.3536		0.2166		0.7229	
65 or Older	0.5393		0.8963		0.3246	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	682	701	205	168	108	106
12	0	2	0	1	*	*
13	5	3	1	3	*	1
14	6	7	6	6	1	1
15	11	17	11	10	6	7
16	15	12	13	10	7	6
17	17	14	13	12	6	7
18	29	29	21	22	14	15
19	31	32	23	24	12	12
20	40	39	23	29	13	19
21	43	37	29	26	16	15
22	43	42	30	25	12	11
23	45	45	23	22	16	11
24	45	41	26	23	13	12
25	47	47	24	28	14	10
26-29	184	183	91	60	38	35
30-34	203	203	85	68	55	30
35-39	193	204	69	56	34	43
40-44	228	205	62	50	34	27
45-49	276	247	51	60	37	36
50-54	372	386	62	60	41	43
55-59	295	274	45	28	4	24
60-64	180	195	31	34	17	26
65 or Older	165	173	*	6	*	5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.27	0.27	0.08	0.07	0.04	0.04
12	0.01	0.06	0.01	0.03	*	*
13	0.13	0.08	0.03	0.06	*	0.02
14	0.15	0.15	0.14	0.14	0.02	0.02
15	0.27	0.40	0.26	0.24	0.14	0.17
16	0.36	0.28	0.32	0.24	0.18	0.14
17	0.40	0.33	0.30	0.28	0.14	0.16
18	0.58	0.59	0.42	0.45	0.29	0.32
19	0.68	0.74	0.51	0.54	0.28	0.27
20	0.82	0.83	0.50	0.63	0.27	0.42
21	0.94	0.78	0.66	0.56	0.37	0.33
22	0.93	0.87	0.69	0.54	0.29	0.25
23	1.02	1.03	0.56	0.52	0.39	0.28
24	1.02	0.92	0.61	0.54	0.30	0.30
25	1.09	1.07	0.60	0.68	0.35	0.26
26-29	0.95	0.97	0.53	0.35	0.23	0.21
30-34	0.89	0.94	0.42	0.34	0.27	0.16
35-39	0.89	0.94	0.36	0.28	0.18	0.22
40-44	0.97	0.93	0.30	0.24	0.17	0.13
45-49	1.04	0.98	0.23	0.28	0.17	0.17
50-54	1.46	1.40	0.28	0.26	0.19	0.18
55-59	1.34	1.28	0.23	0.15	0.02	0.13
60-64	1.05	1.04	0.19	0.19	0.10	0.15
65 or Older	0.42	0.43	*	0.01	*	0.01

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.2758		0.0060		0.4122	
12	0.1796		0.5994		*	
13	0.8194		0.0346		*	
14	0.8339		0.8211		0.8510	
15	0.9523		0.3967		0.5641	
16	0.3962		0.4398		0.5516	
17	0.3281		0.9499		0.5361	
18	0.4965		0.7203		0.6846	
19	0.8490		0.9022		0.8247	
20	0.9874		0.0843		0.1220	
21	0.0528		0.6283		0.1652	
22	0.0107		0.0542		0.2679	
23	0.3472		0.9288		0.3814	
24	0.2318		0.8746		0.8479	
25	0.3876		0.9561		0.0915	
26-29	0.1741		0.0034		0.8092	
30-34	0.2506		0.1734		0.0788	
35-39	0.9673		0.5172		0.5235	
40-44	0.0468		0.3316		0.7116	
45-49	0.0050		0.9960		0.7970	
50-54	0.0554		0.6022		0.7160	
55-59	0.5357		0.6300		0.0497	
60-64	0.0959		0.7978		0.8050	
65 or Older	0.5575		*		*	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	379	327	92	69	60	42
12	0	1	0	1	*	*
13	5	*	0	*	*	*
14	3	3	1	3	*	*
15	4	14	3	2	2	2
16	2	4	2	4	0	0
17	8	6	3	4	1	1
18	13	5	11	3	9	2
19	12	9	10	5	7	2
20	16	16	4	8	*	4
21	17	11	5	5	3	1
22	18	15	8	6	6	2
23	18	17	9	6	8	2
24	16	19	5	5	2	1
25	20	19	8	5	3	2
26-29	84	74	31	13	8	7
30-34	105	95	15	19	4	9
35-39	100	108	34	26	18	9
40-44	127	103	41	35	31	11
45-49	149	110	33	32	26	27
50-54	197	185	38	24	35	22
55-59	127	120	4	14	4	12
60-64	80	90	28	16	13	*
65 or Older	34	43	*	5	*	5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.15	0.13	0.04	0.03	0.02	0.02
12	0.01	0.03	0.01	0.03	*	*
13	0.12	*	0.01	*	*	*
14	0.07	0.08	0.02	0.06	*	*
15	0.11	0.32	0.06	0.05	0.04	0.04
16	0.06	0.09	0.04	0.09	0.01	0.01
17	0.19	0.14	0.08	0.10	0.02	0.03
18	0.27	0.12	0.23	0.06	0.18	0.05
19	0.27	0.21	0.22	0.13	0.17	0.04
20	0.35	0.37	0.08	0.17	*	0.08
21	0.41	0.25	0.12	0.11	0.07	0.03
22	0.42	0.34	0.18	0.15	0.14	0.05
23	0.45	0.43	0.21	0.15	0.20	0.04
24	0.39	0.45	0.13	0.11	0.06	0.02
25	0.49	0.46	0.21	0.13	0.07	0.05
26-29	0.49	0.43	0.19	0.08	0.05	0.04
30-34	0.51	0.47	0.07	0.10	0.02	0.04
35-39	0.51	0.54	0.18	0.13	0.09	0.04
40-44	0.59	0.50	0.20	0.17	0.15	0.05
45-49	0.65	0.49	0.15	0.15	0.12	0.13
50-54	0.86	0.76	0.17	0.10	0.16	0.09
55-59	0.65	0.62	0.02	0.07	0.02	0.06
60-64	0.48	0.50	0.17	0.09	0.08	*
65 or Older	0.09	0.11	*	0.01	*	0.01

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0213		0.0208		0.0388	
12	0.6547		0.5780		*	
13	*		*		*	
14	0.8291		0.5545		*	
15	0.2106		0.9899		0.9006	
16	0.4945		0.2634		0.9431	
17	0.3971		0.5309		0.8540	
18	0.0297		0.0750		0.4930	
19	0.4901		0.1868		0.1511	
20	0.9682		0.1651		*	
21	0.0045		0.3643		0.1004	
22	0.0423		0.0928		0.0240	
23	0.1736		0.2642		0.0921	
24	0.8152		0.3613		0.1271	
25	0.8309		0.2843		0.3924	
26-29	0.2000		0.0718		0.5633	
30-34	0.1067		0.7294		0.4078	
35-39	0.6015		0.3071		0.3468	
40-44	0.1202		0.6581		0.2012	
45-49	0.0011		0.3904		0.9073	
50-54	0.3679		0.3473		0.3515	
55-59	0.7093		0.1954		0.4541	
60-64	0.5555		0.5252		*	
65 or Older	0.7264		*		*	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	687	628	178	143	84	60
12	4	5	3	2	0	1
13	10	11	9	5	6	2
14	14	14	13	12	9	9
15	18	20	17	16	10	10
16	23	25	20	22	9	15
17	28	24	25	19	12	10
18	38	40	33	32	19	14
19	42	42	31	33	16	16
20	49	48	33	36	16	14
21	51	49	33	27	19	17
22	49	51	34	30	13	15
23	52	48	28	27	13	10
24	50	48	28	22	17	11
25	52	49	21	25	10	13
26-29	202	201	74	58	39	18
30-34	222	219	73	49	34	26
35-39	203	181	30	51	6	13
40-44	197	191	28	34	18	5
45-49	241	214	27	32	20	15
50-54	330	315	57	26	17	*
55-59	298	243	*	6	*	2
60-64	188	192	12	*	*	*
65 or Older	125	141	10	5	*	*

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.27	0.24	0.07	0.06	0.03	0.02
12	0.11	0.14	0.08	0.06	0.01	0.02
13	0.25	0.25	0.22	0.11	0.16	0.05
14	0.35	0.32	0.32	0.29	0.22	0.21
15	0.41	0.46	0.39	0.38	0.24	0.22
16	0.54	0.55	0.48	0.49	0.22	0.34
17	0.64	0.55	0.58	0.44	0.27	0.23
18	0.73	0.74	0.64	0.61	0.39	0.28
19	0.88	0.88	0.67	0.72	0.37	0.36
20	0.97	0.97	0.68	0.74	0.34	0.32
21	1.10	1.00	0.75	0.59	0.44	0.37
22	1.01	1.02	0.75	0.64	0.30	0.34
23	1.12	1.05	0.68	0.66	0.32	0.25
24	1.08	1.03	0.65	0.53	0.41	0.25
25	1.16	1.14	0.52	0.61	0.26	0.33
26-29	1.03	1.05	0.43	0.33	0.24	0.11
30-34	0.95	0.94	0.36	0.25	0.17	0.13
35-39	0.89	0.86	0.16	0.26	0.03	0.07
40-44	0.86	0.84	0.14	0.16	0.09	0.03
45-49	0.96	0.88	0.12	0.15	0.09	0.07
50-54	1.32	1.23	0.26	0.11	0.08	*
55-59	1.39	1.18	*	0.03	*	0.01
60-64	1.09	1.02	0.07	*	*	*
65 or Older	0.32	0.35	0.03	0.01	*	*

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0495		0.0205		0.0171	
12	0.7639		0.4057		0.3372	
13	0.7423		0.1069		0.1501	
14	0.2492		0.1694		0.2122	
15	0.5843		0.2765		0.6148	
16	0.8055		0.5536		0.0134	
17	0.1203		0.0415		0.2502	
18	0.7121		0.7059		0.1366	
19	0.5574		0.7534		0.5183	
20	0.9092		0.4074		0.6061	
21	0.3576		0.1253		0.4318	
22	0.2721		0.0893		0.5993	
23	0.7722		0.5982		0.2972	
24	0.1916		0.2995		0.2674	
25	0.6009		0.1175		0.5927	
26-29	0.0199		0.0737		0.0517	
30-34	0.4632		0.1041		0.7634	
35-39	0.3977		0.1015		0.3136	
40-44	0.3536		0.4707		0.2424	
45-49	0.1452		0.7539		0.9822	
50-54	0.4432		0.1874		*	
55-59	0.5457		*		*	
60-64	0.5777		*		*	
65 or Older	0.1280		0.6530		*	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	473	491	115	107	83	68
12	23	22	16	14	8	9
13	27	24	21	17	14	10
14	25	26	19	20	10	10
15	32	24	18	15	9	6
16	26	26	16	17	7	6
17	25	24	14	15	8	8
18	30	31	19	18	7	5
19	32	32	16	20	6	17
20	35	33	18	16	5	11
21	36	29	14	14	7	11
22	36	40	13	19	8	8
23	37	35	18	11	12	8
24	32	35	13	8	9	3
25	36	35	6	5	3	2
26-29	140	141	33	33	15	15
30-34	173	136	32	30	20	25
35-39	160	162	24	20	14	9
40-44	158	157	24	25	15	22
45-49	177	170	20	23	19	23
50-54	215	193	75	53	66	15
55-59	188	195	3	23	3	23
60-64	106	99	18	24	18	10
65 or Older	73	109	19	11	*	11

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.19	0.19	0.05	0.04	0.03	0.03
12	0.58	0.56	0.42	0.37	0.21	0.23
13	0.64	0.55	0.50	0.40	0.34	0.24
14	0.62	0.59	0.47	0.46	0.26	0.23
15	0.71	0.53	0.42	0.36	0.22	0.15
16	0.61	0.58	0.38	0.39	0.17	0.14
17	0.54	0.56	0.32	0.34	0.19	0.18
18	0.60	0.61	0.39	0.38	0.16	0.11
19	0.71	0.72	0.37	0.46	0.13	0.39
20	0.73	0.71	0.39	0.35	0.12	0.24
21	0.78	0.64	0.32	0.32	0.16	0.25
22	0.79	0.82	0.31	0.42	0.18	0.19
23	0.86	0.82	0.44	0.26	0.29	0.20
24	0.76	0.77	0.31	0.20	0.22	0.08
25	0.85	0.85	0.15	0.12	0.09	0.06
26-29	0.77	0.78	0.20	0.20	0.09	0.09
30-34	0.79	0.66	0.16	0.15	0.10	0.13
35-39	0.76	0.79	0.12	0.10	0.07	0.05
40-44	0.72	0.73	0.12	0.12	0.07	0.11
45-49	0.75	0.73	0.09	0.10	0.09	0.10
50-54	0.93	0.80	0.34	0.23	0.30	0.06
55-59	0.93	0.98	0.02	0.12	0.02	0.12
60-64	0.63	0.55	0.11	0.13	0.11	0.05
65 or Older	0.19	0.27	0.05	0.03	*	0.03

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0190		0.1784		0.4747	
12	0.1991		0.1828		0.7095	
13	0.1924		0.0965		0.1395	
14	0.6724		0.6900		0.5413	
15	0.0329		0.0853		0.0714	
16	0.6081		0.5521		0.9034	
17	0.9319		0.3979		0.2638	
18	0.1852		0.2465		0.1232	
19	0.1913		0.7369		0.2192	
20	0.4757		0.7018		0.1706	
21	0.0069		0.4921		0.6395	
22	0.3707		0.6854		0.6196	
23	0.8797		0.2449		0.7754	
24	0.9388		0.3569		0.3477	
25	0.5743		0.1336		0.3711	
26-29	0.2376		0.6864		0.8991	
30-34	0.0020		0.4694		0.8410	
35-39	0.3479		0.6980		0.3597	
40-44	0.9727		0.9728		0.6865	
45-49	0.1290		0.7976		0.6673	
50-54	0.6050		0.6642		0.2073	
55-59	0.1320		0.2575		0.2575	
60-64	0.9301		0.8765		0.6608	
65 or Older	0.0635		0.8024		*	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	784	752	397	363	268	231
12	19	19	16	15	14	10
13	22	19	18	14	13	8
14	23	23	19	19	14	12
15	30	28	25	25	16	15
16	35	35	30	30	17	17
17	39	37	33	28	17	17
18	53	54	44	39	28	24
19	55	53	41	41	26	27
20	61	65	47	46	29	26
21	62	57	45	39	28	22
22	61	63	43	45	27	28
23	57	57	37	35	24	21
24	62	63	41	38	24	25
25	66	60	39	34	23	22
26-29	227	246	130	123	79	87
30-34	234	239	126	129	93	85
35-39	216	232	104	112	65	75
40-44	233	225	116	117	75	72
45-49	261	254	108	106	65	68
50-54	386	351	178	126	126	88
55-59	317	300	130	123	57	71
60-64	207	222	82	62	42	29
65 or Older	227	194	89	65	71	35

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.31	0.29	0.16	0.14	0.11	0.09
12	0.50	0.49	0.41	0.38	0.36	0.26
13	0.52	0.44	0.43	0.33	0.31	0.20
14	0.57	0.53	0.46	0.43	0.36	0.29
15	0.66	0.61	0.57	0.56	0.38	0.34
16	0.77	0.73	0.68	0.65	0.40	0.39
17	0.84	0.80	0.72	0.62	0.40	0.40
18	0.94	0.92	0.82	0.72	0.56	0.49
19	1.08	1.05	0.87	0.86	0.57	0.59
20	1.17	1.16	0.96	0.95	0.63	0.57
21	1.18	1.07	0.90	0.82	0.61	0.48
22	1.15	1.19	0.93	0.92	0.62	0.61
23	1.18	1.22	0.85	0.85	0.58	0.53
24	1.24	1.28	0.94	0.87	0.56	0.59
25	1.27	1.25	0.88	0.81	0.58	0.53
26-29	1.11	1.16	0.73	0.69	0.46	0.51
30-34	0.99	1.00	0.60	0.63	0.46	0.43
35-39	0.96	1.02	0.52	0.56	0.34	0.38
40-44	1.02	1.00	0.56	0.56	0.36	0.35
45-49	0.99	0.98	0.47	0.48	0.29	0.31
50-54	1.51	1.31	0.79	0.53	0.57	0.38
55-59	1.43	1.34	0.66	0.63	0.30	0.37
60-64	1.17	1.14	0.49	0.34	0.25	0.16
65 or Older	0.58	0.48	0.23	0.16	0.18	0.09

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.2064		0.0026		0.0078	
12	0.5944		0.5798		0.1922	
13	0.2716		0.0491		0.0427	
14	0.5596		0.8595		0.3997	
15	0.1344		0.5495		0.6641	
16	0.9503		0.7688		0.7665	
17	0.8387		0.3362		0.4287	
18	0.0642		0.0102		0.0541	
19	0.0296		0.1208		0.3929	
20	0.9768		0.7840		0.7740	
21	0.0869		0.0068		0.0025	
22	0.2498		0.1374		0.2396	
23	0.4279		0.9549		0.7409	
24	0.5778		0.6157		0.9280	
25	0.1502		0.0365		0.2645	
26-29	0.3996		0.1255		0.6029	
30-34	0.8035		0.7224		0.4913	
35-39	0.2446		0.7323		0.7853	
40-44	0.6209		0.7507		0.8220	
45-49	0.3703		0.4040		0.3588	
50-54	0.7781		0.0982		0.3884	
55-59	0.3751		0.7798		0.5534	
60-64	0.2882		0.9965		0.5972	
65 or Older	0.3670		0.2119		0.0488	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	610	598	336	302	215	189
12	19	17	16	14	13	9
13	21	17	17	13	12	7
14	22	22	17	16	13	12
15	29	26	24	23	15	13
16	33	33	28	28	15	16
17	37	33	31	26	16	14
18	47	47	38	33	25	21
19	50	47	38	35	22	24
20	55	54	41	38	26	20
21	53	51	38	35	23	18
22	55	56	40	38	24	23
23	51	52	31	32	20	18
24	57	53	36	34	19	19
25	59	54	33	31	22	18
26-29	200	226	122	104	68	75
30-34	205	206	123	113	88	71
35-39	191	190	88	96	59	62
40-44	189	178	95	96	64	60
45-49	184	193	95	87	60	58
50-54	261	254	142	94	102	69
55-59	211	219	116	109	39	54
60-64	161	134	76	56	29	28
65 or Older	131	149	56	61	47	32

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.24	0.23	0.13	0.12	0.08	0.07
12	0.48	0.46	0.40	0.36	0.35	0.24
13	0.49	0.40	0.41	0.31	0.29	0.17
14	0.54	0.49	0.44	0.37	0.34	0.27
15	0.64	0.58	0.55	0.52	0.36	0.31
16	0.74	0.70	0.64	0.61	0.35	0.36
17	0.78	0.72	0.67	0.58	0.38	0.34
18	0.87	0.85	0.73	0.64	0.51	0.43
19	1.01	0.94	0.82	0.75	0.49	0.54
20	1.07	1.04	0.85	0.84	0.56	0.43
21	1.06	0.98	0.80	0.75	0.52	0.41
22	1.09	1.08	0.86	0.79	0.56	0.51
23	1.11	1.15	0.74	0.78	0.50	0.45
24	1.18	1.13	0.82	0.79	0.46	0.46
25	1.18	1.18	0.78	0.74	0.54	0.45
26-29	1.04	1.09	0.69	0.60	0.40	0.44
30-34	0.90	0.94	0.59	0.56	0.44	0.36
35-39	0.87	0.88	0.45	0.48	0.30	0.32
40-44	0.85	0.81	0.45	0.46	0.31	0.29
45-49	0.76	0.83	0.42	0.40	0.27	0.27
50-54	1.11	1.01	0.64	0.40	0.46	0.30
55-59	1.03	1.06	0.59	0.56	0.20	0.28
60-64	0.94	0.73	0.46	0.31	0.18	0.15
65 or Older	0.34	0.37	0.14	0.15	0.12	0.08

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.1653		0.0051		0.0151	
12	0.4726		0.5241		0.1667	
13	0.2215		0.0543		0.0515	
14	0.7777		0.7893		0.3903	
15	0.0444		0.4764		0.7742	
16	0.9121		0.5842		0.3672	
17	0.7738		0.7174		0.6393	
18	0.1086		0.0615		0.0231	
19	0.0116		0.0784		0.8384	
20	0.5260		0.7973		0.2797	
21	0.1217		0.0198		0.0317	
22	0.4594		0.1058		0.1936	
23	0.8223		0.6649		0.4794	
24	0.3359		0.5349		0.5185	
25	0.1531		0.1344		0.0824	
26-29	0.5386		0.0339		0.8638	
30-34	0.5628		0.8657		0.5637	
35-39	0.0580		0.5282		0.2932	
40-44	0.5895		0.6861		0.8788	
45-49	0.9148		0.2466		0.3711	
50-54	0.3503		0.1982		0.5032	
55-59	0.1966		0.9049		0.3404	
60-64	0.3074		0.8829		0.6608	
65 or Older	0.3268		0.7909		0.4761	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.