

Table 1.1C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 1,023 | 1,036 | 646 | 603 | 488 | 460 |
| Marijuana and Hashish | 1,019 | 1,013 | 559 | 514 | 407 | 409 |
| Cocaine | 682 | 701 | 205 | 168 | 108 | 106 |
| Crack | 379 | 327 | 92 | 69 | 60 | 42 |
| Heroin | 249 | 227 | 75 | 72 | 43 | 48 |
| Hallucinogens | 687 | 628 | 178 | 143 | 84 | 60 |
| LSD | 560 | 530 | 72 | 60 | 20 | 23 |
| PCP | 321 | 307 | 24 | 20 | 21 | 10 |
| Ecstasy | 430 | 367 | 128 | 107 | 61 | 45 |
| Inhalants | 473 | 491 | 115 | 107 | 83 | 68 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 784 | 752 | 397 | 363 | 268 | 231 |
| Pain Relievers | 610 | 598 | 336 | 302 | 215 | 189 |
| OxyContin [®] | 246 | 211 | 116 | 104 | 67 | 54 |
| Tranquilizers | 494 | 494 | 207 | 211 | 138 | 133 |
| Stimulants ³ | 557 | 500 | 158 | 151 | 86 | 89 |
| Methamphetamine ³ | 452 | 410 | 105 | 118 | 53 | 72 |
| Sedatives | 355 | 353 | 103 | 64 | 72 | 49 |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 900 | 889 | 446 | 404 | 300 | 256 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.1D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.40 | 0.40 | 0.25 | 0.23 | 0.19 | 0.18 |
| Marijuana and Hashish | 0.40 | 0.39 | 0.22 | 0.20 | 0.16 | 0.16 |
| Cocaine | 0.27 | 0.27 | 0.08 | 0.07 | 0.04 | 0.04 |
| Crack | 0.15 | 0.13 | 0.04 | 0.03 | 0.02 | 0.02 |
| Heroin | 0.10 | 0.09 | 0.03 | 0.03 | 0.02 | 0.02 |
| Hallucinogens | 0.27 | 0.24 | 0.07 | 0.06 | 0.03 | 0.02 |
| LSD | 0.22 | 0.21 | 0.03 | 0.02 | 0.01 | 0.01 |
| PCP | 0.13 | 0.12 | 0.01 | 0.01 | 0.01 | 0.00 |
| Ecstasy | 0.17 | 0.14 | 0.05 | 0.04 | 0.02 | 0.02 |
| Inhalants | 0.19 | 0.19 | 0.05 | 0.04 | 0.03 | 0.03 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.31 | 0.29 | 0.16 | 0.14 | 0.11 | 0.09 |
| Pain Relievers | 0.24 | 0.23 | 0.13 | 0.12 | 0.08 | 0.07 |
| OxyContin [®] | 0.10 | 0.08 | 0.05 | 0.04 | 0.03 | 0.02 |
| Tranquilizers | 0.19 | 0.19 | 0.08 | 0.08 | 0.05 | 0.05 |
| Stimulants ³ | 0.22 | 0.19 | 0.06 | 0.06 | 0.03 | 0.03 |
| Methamphetamine ³ | 0.18 | 0.16 | 0.04 | 0.05 | 0.02 | 0.03 |
| Sedatives | 0.14 | 0.14 | 0.04 | 0.02 | 0.03 | 0.02 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.35 | 0.35 | 0.18 | 0.16 | 0.12 | 0.10 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.1P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.6146 | | 0.1638 | | 0.4074 | |
| Marijuana and Hashish | 0.7538 | | 0.9765 | | 0.4881 | |
| Cocaine | 0.2758 | | 0.0060 | | 0.4122 | |
| Crack | 0.0213 | | 0.0208 | | 0.0388 | |
| Heroin | 0.8940 | | 0.9195 | | 0.5623 | |
| Hallucinogens | 0.0495 | | 0.0205 | | 0.0171 | |
| LSD | 0.3138 | | 0.8704 | | 0.8597 | |
| PCP | 0.5683 | | 0.5027 | | 0.6568 | |
| Ecstasy | 0.0042 | | 0.1184 | | 0.0420 | |
| Inhalants | 0.0190 | | 0.1784 | | 0.4747 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.2064 | | 0.0026 | | 0.0078 | |
| Pain Relievers | 0.1653 | | 0.0051 | | 0.0151 | |
| OxyContin [®] | 0.2809 | | 0.0647 | | 0.1071 | |
| Tranquilizers | 0.1904 | | 0.0634 | | 0.0651 | |
| Stimulants ³ | 0.0218 | | 0.2586 | | 0.3150 | |
| Methamphetamine ³ | 0.0239 | | 0.7112 | | 0.3941 | |
| Sedatives | 0.6020 | | 0.0012 | | 0.0892 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.0783 | | 0.0013 | | 0.0048 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.2C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 45 | 46 | 37 | 34 | 26 | 24 |
| Marijuana and Hashish | 22 | 25 | 21 | 22 | 12 | 16 |
| Cocaine | 5 | 4 | 1 | 3 | * | 1 |
| Crack | 5 | 1 | 1 | 1 | * | * |
| Heroin | 4 | 5 | 4 | 5 | 2 | * |
| Hallucinogens | 11 | 12 | 9 | 5 | 7 | 2 |
| LSD | 4 | 4 | 2 | 2 | 1 | 1 |
| PCP | 3 | 2 | 1 | 2 | * | * |
| Ecstasy | 7 | 9 | 7 | 4 | 6 | 2 |
| Inhalants | 35 | 32 | 26 | 22 | 17 | 13 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 29 | 26 | 24 | 20 | 18 | 13 |
| Pain Relievers | 27 | 23 | 22 | 19 | 17 | 12 |
| OxyContin [®] | 6 | 6 | 5 | 4 | 1 | 2 |
| Tranquilizers | 8 | 8 | 7 | 7 | 4 | 4 |
| Stimulants ³ | 10 | 12 | 6 | 9 | 4 | 5 |
| Methamphetamine ³ | 4 | 9 | 3 | 6 | 1 | 2 |
| Sedatives | 8 | 7 | 4 | 5 | 3 | 2 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 43 | 41 | 35 | 29 | 24 | 18 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.2D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.53 | 0.53 | 0.44 | 0.41 | 0.32 | 0.30 |
| Marijuana and Hashish | 0.27 | 0.32 | 0.25 | 0.27 | 0.15 | 0.20 |
| Cocaine | 0.07 | 0.05 | 0.02 | 0.04 | * | 0.01 |
| Crack | 0.06 | 0.01 | 0.01 | 0.01 | * | * |
| Heroin | 0.05 | 0.06 | 0.05 | 0.06 | 0.03 | * |
| Hallucinogens | 0.14 | 0.15 | 0.12 | 0.07 | 0.08 | 0.03 |
| LSD | 0.04 | 0.05 | 0.02 | 0.02 | 0.01 | 0.01 |
| PCP | 0.04 | 0.02 | 0.02 | 0.02 | * | * |
| Ecstasy | 0.09 | 0.12 | 0.09 | 0.05 | 0.08 | 0.02 |
| Inhalants | 0.43 | 0.39 | 0.33 | 0.27 | 0.21 | 0.17 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.34 | 0.32 | 0.29 | 0.25 | 0.22 | 0.16 |
| Pain Relievers | 0.32 | 0.29 | 0.27 | 0.24 | 0.21 | 0.15 |
| OxyContin [®] | 0.08 | 0.07 | 0.07 | 0.05 | 0.01 | 0.03 |
| Tranquilizers | 0.10 | 0.10 | 0.09 | 0.08 | 0.05 | 0.05 |
| Stimulants ³ | 0.12 | 0.16 | 0.08 | 0.12 | 0.06 | 0.06 |
| Methamphetamine ³ | 0.05 | 0.12 | 0.04 | 0.08 | 0.01 | 0.02 |
| Sedatives | 0.10 | 0.09 | 0.06 | 0.07 | 0.04 | 0.02 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.50 | 0.48 | 0.42 | 0.35 | 0.30 | 0.23 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.2P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.1268 | | 0.0134 | | 0.1369 | |
| Marijuana and Hashish | 0.3771 | | 0.8137 | | 0.0921 | |
| Cocaine | 0.4894 | | 0.0296 | | * | |
| Crack | 0.3315 | | 0.9100 | | * | |
| Heroin | 0.9799 | | 0.7638 | | * | |
| Hallucinogens | 0.6951 | | 0.0697 | | 0.2040 | |
| LSD | 0.4864 | | 0.8050 | | 0.4351 | |
| PCP | 0.6776 | | 0.3233 | | * | |
| Ecstasy | 0.5387 | | 0.3296 | | 0.1249 | |
| Inhalants | 0.0726 | | 0.0379 | | 0.1645 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.2374 | | 0.0684 | | 0.0143 | |
| Pain Relievers | 0.1559 | | 0.0650 | | 0.0136 | |
| OxyContin [®] | 0.8328 | | 0.7100 | | 0.2819 | |
| Tranquilizers | 0.5879 | | 0.8670 | | 0.5820 | |
| Stimulants ³ | 0.2972 | | 0.2378 | | 0.5259 | |
| Methamphetamine ³ | 0.0236 | | 0.1620 | | 0.1691 | |
| Sedatives | 0.9124 | | 0.6135 | | 0.5680 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.0545 | | 0.0036 | | 0.0031 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.3C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 67 | 65 | 56 | 59 | 41 | 40 |
| Marijuana and Hashish | 52 | 53 | 48 | 50 | 34 | 34 |
| Cocaine | 13 | 19 | 12 | 12 | 6 | 7 |
| Crack | 5 | 14 | 3 | 3 | 2 | 2 |
| Heroin | 4 | 7 | 3 | 7 | 1 | 4 |
| Hallucinogens | 23 | 24 | 21 | 21 | 14 | 13 |
| LSD | 11 | 11 | 9 | 10 | 4 | 6 |
| PCP | 8 | 6 | 7 | 6 | 2 | 1 |
| Ecstasy | 18 | 18 | 17 | 15 | 11 | 8 |
| Inhalants | 41 | 37 | 26 | 25 | 14 | 12 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 40 | 36 | 33 | 31 | 22 | 19 |
| Pain Relievers | 38 | 33 | 31 | 28 | 20 | 17 |
| OxyContin [®] | 14 | 12 | 13 | 10 | 7 | 4 |
| Tranquilizers | 19 | 19 | 17 | 17 | 9 | 7 |
| Stimulants ³ | 17 | 16 | 15 | 13 | 9 | 7 |
| Methamphetamine ³ | 12 | 9 | 10 | 9 | 6 | 6 |
| Sedatives | 12 | 11 | 11 | 8 | 7 | 5 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 55 | 52 | 42 | 42 | 28 | 25 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.3D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.69 | 0.65 | 0.62 | 0.61 | 0.48 | 0.44 |
| Marijuana and Hashish | 0.58 | 0.57 | 0.54 | 0.54 | 0.40 | 0.39 |
| Cocaine | 0.16 | 0.22 | 0.15 | 0.15 | 0.07 | 0.08 |
| Crack | 0.06 | 0.16 | 0.03 | 0.04 | 0.02 | 0.02 |
| Heroin | 0.05 | 0.08 | 0.03 | 0.08 | 0.01 | 0.05 |
| Hallucinogens | 0.27 | 0.28 | 0.26 | 0.24 | 0.17 | 0.15 |
| LSD | 0.14 | 0.12 | 0.12 | 0.12 | 0.05 | 0.07 |
| PCP | 0.10 | 0.07 | 0.08 | 0.07 | 0.02 | 0.01 |
| Ecstasy | 0.22 | 0.21 | 0.20 | 0.17 | 0.13 | 0.10 |
| Inhalants | 0.47 | 0.40 | 0.32 | 0.29 | 0.18 | 0.14 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.45 | 0.40 | 0.39 | 0.36 | 0.26 | 0.22 |
| Pain Relievers | 0.44 | 0.37 | 0.37 | 0.32 | 0.25 | 0.20 |
| OxyContin [®] | 0.18 | 0.14 | 0.16 | 0.11 | 0.09 | 0.05 |
| Tranquilizers | 0.23 | 0.23 | 0.21 | 0.19 | 0.12 | 0.09 |
| Stimulants ³ | 0.21 | 0.19 | 0.18 | 0.16 | 0.11 | 0.09 |
| Methamphetamine ³ | 0.15 | 0.11 | 0.13 | 0.10 | 0.08 | 0.07 |
| Sedatives | 0.15 | 0.12 | 0.14 | 0.10 | 0.08 | 0.05 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.59 | 0.55 | 0.49 | 0.47 | 0.33 | 0.29 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.3P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.4449 | | 0.7242 | | 0.8502 | |
| Marijuana and Hashish | 0.8537 | | 0.9132 | | 0.7065 | |
| Cocaine | 0.9215 | | 0.4943 | | 0.6210 | |
| Crack | 0.2617 | | 0.7404 | | 0.9502 | |
| Heroin | 0.1561 | | 0.0401 | | 0.2254 | |
| Hallucinogens | 0.2254 | | 0.0784 | | 0.2195 | |
| LSD | 0.4571 | | 0.9709 | | 0.9025 | |
| PCP | 0.1125 | | 0.3750 | | 0.3161 | |
| Ecstasy | 0.2885 | | 0.1025 | | 0.1351 | |
| Inhalants | 0.0644 | | 0.1636 | | 0.1186 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.0983 | | 0.4881 | | 0.3504 | |
| Pain Relievers | 0.0606 | | 0.3960 | | 0.4124 | |
| OxyContin [®] | 0.1419 | | 0.2917 | | 0.3584 | |
| Tranquilizers | 0.6784 | | 0.4424 | | 0.2229 | |
| Stimulants ³ | 0.3898 | | 0.3769 | | 0.4384 | |
| Methamphetamine ³ | 0.1828 | | 0.3340 | | 0.8912 | |
| Sedatives | 0.1888 | | 0.0834 | | 0.2605 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.0934 | | 0.3056 | | 0.2336 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.4C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 87 | 84 | 78 | 71 | 55 | 53 |
| Marijuana and Hashish | 76 | 76 | 70 | 67 | 51 | 50 |
| Cocaine | 23 | 19 | 19 | 15 | 10 | 9 |
| Crack | 8 | 7 | 4 | 6 | 1 | 1 |
| Heroin | 7 | 9 | 6 | 8 | 2 | 6 |
| Hallucinogens | 37 | 35 | 32 | 29 | 15 | 18 |
| LSD | 17 | 17 | 14 | 15 | 9 | 6 |
| PCP | 7 | 8 | 3 | 4 | 2 | 2 |
| Ecstasy | 30 | 31 | 27 | 26 | 10 | 13 |
| Inhalants | 36 | 36 | 21 | 22 | 10 | 10 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 54 | 51 | 46 | 40 | 26 | 25 |
| Pain Relievers | 51 | 48 | 41 | 38 | 23 | 21 |
| OxyContin [®] | 23 | 22 | 18 | 15 | 7 | 8 |
| Tranquilizers | 31 | 30 | 25 | 25 | 12 | 15 |
| Stimulants ³ | 24 | 22 | 18 | 16 | 11 | 9 |
| Methamphetamine ³ | 14 | 15 | 10 | 11 | 5 | 6 |
| Sedatives | 12 | 9 | 10 | 6 | 3 | 3 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 64 | 60 | 54 | 47 | 31 | 31 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.4D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.78 | 0.79 | 0.74 | 0.71 | 0.58 | 0.55 |
| Marijuana and Hashish | 0.72 | 0.75 | 0.69 | 0.67 | 0.54 | 0.53 |
| Cocaine | 0.27 | 0.21 | 0.22 | 0.18 | 0.11 | 0.11 |
| Crack | 0.10 | 0.08 | 0.04 | 0.07 | 0.01 | 0.01 |
| Heroin | 0.08 | 0.10 | 0.07 | 0.09 | 0.03 | 0.07 |
| Hallucinogens | 0.41 | 0.39 | 0.37 | 0.33 | 0.18 | 0.21 |
| LSD | 0.20 | 0.19 | 0.17 | 0.17 | 0.10 | 0.07 |
| PCP | 0.08 | 0.09 | 0.04 | 0.05 | 0.02 | 0.02 |
| Ecstasy | 0.35 | 0.35 | 0.32 | 0.30 | 0.12 | 0.15 |
| Inhalants | 0.40 | 0.41 | 0.25 | 0.26 | 0.12 | 0.11 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.57 | 0.53 | 0.49 | 0.44 | 0.30 | 0.28 |
| Pain Relievers | 0.54 | 0.50 | 0.45 | 0.41 | 0.26 | 0.25 |
| OxyContin [®] | 0.26 | 0.25 | 0.21 | 0.17 | 0.08 | 0.10 |
| Tranquilizers | 0.35 | 0.33 | 0.29 | 0.28 | 0.14 | 0.17 |
| Stimulants ³ | 0.27 | 0.26 | 0.21 | 0.19 | 0.13 | 0.10 |
| Methamphetamine ³ | 0.16 | 0.18 | 0.12 | 0.13 | 0.06 | 0.07 |
| Sedatives | 0.14 | 0.10 | 0.11 | 0.08 | 0.03 | 0.04 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.65 | 0.61 | 0.57 | 0.50 | 0.35 | 0.34 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.4P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.5160 | | 0.9842 | | 0.5081 | |
| Marijuana and Hashish | 0.5737 | | 0.8221 | | 0.3663 | |
| Cocaine | 0.1776 | | 0.6007 | | 0.9805 | |
| Crack | 0.5937 | | 0.2452 | | 0.8310 | |
| Heroin | 0.2826 | | 0.2445 | | 0.6690 | |
| Hallucinogens | 0.2898 | | 0.2585 | | 0.2860 | |
| LSD | 0.7961 | | 0.3558 | | 0.2642 | |
| PCP | 0.0991 | | 0.1140 | | 0.9748 | |
| Ecstasy | 0.8159 | | 0.6758 | | 0.4503 | |
| Inhalants | 0.7592 | | 0.3110 | | 0.3530 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.8049 | | 0.5974 | | 0.4699 | |
| Pain Relievers | 0.7507 | | 0.9108 | | 0.7585 | |
| OxyContin [®] | 0.6403 | | 0.8829 | | 0.4991 | |
| Tranquilizers | 0.3350 | | 0.7697 | | 0.0570 | |
| Stimulants ³ | 0.2932 | | 0.1895 | | 0.6179 | |
| Methamphetamine ³ | 0.3911 | | 0.6757 | | 0.3670 | |
| Sedatives | 0.2351 | | 0.3533 | | 0.8887 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.4817 | | 0.2124 | | 0.3757 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.5C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 173 | 213 | 151 | 177 | 110 | 133 |
| Marijuana and Hashish | 161 | 202 | 141 | 169 | 103 | 123 |
| Cocaine | 62 | 60 | 42 | 45 | 23 | 28 |
| Crack | 24 | 20 | 15 | 10 | 11 | 5 |
| Heroin | 26 | 18 | 20 | 16 | 12 | 12 |
| Hallucinogens | 79 | 80 | 60 | 63 | 31 | 26 |
| LSD | 39 | 40 | 26 | 30 | 12 | 9 |
| PCP | 19 | 17 | 8 | 11 | * | 3 |
| Ecstasy | 66 | 68 | 48 | 49 | 25 | 18 |
| Inhalants | 56 | 58 | 31 | 35 | 11 | 26 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 104 | 110 | 80 | 76 | 50 | 46 |
| Pain Relievers | 93 | 93 | 71 | 62 | 44 | 38 |
| OxyContin [®] | 43 | 45 | 31 | 26 | 18 | 10 |
| Tranquilizers | 64 | 61 | 45 | 43 | 22 | 27 |
| Stimulants ³ | 53 | 53 | 37 | 39 | 23 | 19 |
| Methamphetamine ³ | 30 | 29 | 18 | 17 | 13 | 8 |
| Sedatives | 19 | 19 | 13 | 11 | 9 | 6 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 119 | 137 | 93 | 101 | 58 | 62 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.5D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.74 | 0.86 | 0.78 | 0.81 | 0.67 | 0.72 |
| Marijuana and Hashish | 0.75 | 0.87 | 0.76 | 0.81 | 0.64 | 0.68 |
| Cocaine | 0.43 | 0.42 | 0.30 | 0.32 | 0.17 | 0.20 |
| Crack | 0.17 | 0.15 | 0.11 | 0.07 | 0.08 | 0.04 |
| Heroin | 0.19 | 0.13 | 0.14 | 0.12 | 0.09 | 0.09 |
| Hallucinogens | 0.53 | 0.51 | 0.41 | 0.41 | 0.22 | 0.19 |
| LSD | 0.28 | 0.28 | 0.19 | 0.21 | 0.09 | 0.07 |
| PCP | 0.14 | 0.12 | 0.06 | 0.08 | * | 0.02 |
| Ecstasy | 0.46 | 0.44 | 0.34 | 0.33 | 0.18 | 0.13 |
| Inhalants | 0.39 | 0.40 | 0.22 | 0.25 | 0.08 | 0.19 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.64 | 0.62 | 0.53 | 0.49 | 0.35 | 0.32 |
| Pain Relievers | 0.60 | 0.54 | 0.49 | 0.43 | 0.31 | 0.27 |
| OxyContin [®] | 0.31 | 0.31 | 0.23 | 0.19 | 0.13 | 0.07 |
| Tranquilizers | 0.44 | 0.42 | 0.32 | 0.31 | 0.16 | 0.20 |
| Stimulants ³ | 0.35 | 0.36 | 0.26 | 0.27 | 0.16 | 0.13 |
| Methamphetamine ³ | 0.22 | 0.22 | 0.13 | 0.13 | 0.09 | 0.06 |
| Sedatives | 0.14 | 0.14 | 0.10 | 0.08 | 0.07 | 0.04 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.68 | 0.70 | 0.60 | 0.59 | 0.40 | 0.41 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.5P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.4235 | | 0.4961 | | 0.5477 | |
| Marijuana and Hashish | 0.4584 | | 0.0859 | | 0.1552 | |
| Cocaine | 0.6146 | | 0.4186 | | 0.1733 | |
| Crack | 0.1868 | | 0.2068 | | 0.3584 | |
| Heroin | 0.0096 | | 0.8607 | | 0.8985 | |
| Hallucinogens | 0.5488 | | 0.9417 | | 0.1245 | |
| LSD | 0.4631 | | 0.8880 | | 0.6175 | |
| PCP | 0.2700 | | 0.6863 | | * | |
| Ecstasy | 0.8174 | | 0.5399 | | 0.1007 | |
| Inhalants | 0.0547 | | 0.5096 | | 0.3770 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.0293 | | 0.0395 | | 0.0830 | |
| Pain Relievers | 0.0071 | | 0.0339 | | 0.0408 | |
| OxyContin [®] | 0.4859 | | 0.0682 | | 0.0016 | |
| Tranquilizers | 0.2030 | | 0.5493 | | 0.3127 | |
| Stimulants ³ | 0.0032 | | 0.1675 | | 0.1433 | |
| Methamphetamine ³ | 0.1320 | | 0.6215 | | 0.3065 | |
| Sedatives | 0.4804 | | 0.3954 | | 0.3018 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.0524 | | 0.2975 | | 0.2584 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.6C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 231 | 246 | 168 | 163 | 121 | 119 |
| Marijuana and Hashish | 218 | 235 | 153 | 150 | 110 | 109 |
| Cocaine | 105 | 97 | 60 | 57 | 32 | 28 |
| Crack | 41 | 37 | 16 | 12 | 11 | 4 |
| Heroin | 30 | 36 | 16 | 19 | 12 | 14 |
| Hallucinogens | 125 | 110 | 71 | 61 | 36 | 30 |
| LSD | 72 | 65 | 37 | 32 | 12 | 10 |
| PCP | 28 | 28 | 6 | 12 | 3 | 5 |
| Ecstasy | 102 | 93 | 54 | 49 | 28 | 25 |
| Inhalants | 81 | 82 | 30 | 28 | 18 | 16 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 153 | 145 | 103 | 92 | 58 | 55 |
| Pain Relievers | 134 | 127 | 86 | 80 | 49 | 46 |
| OxyContin [®] | 65 | 59 | 32 | 32 | 18 | 19 |
| Tranquilizers | 101 | 92 | 59 | 50 | 33 | 31 |
| Stimulants ³ | 90 | 83 | 47 | 41 | 27 | 26 |
| Methamphetamine ³ | 55 | 49 | 20 | 18 | 11 | 12 |
| Sedatives | 29 | 31 | 15 | 14 | 8 | 9 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 177 | 168 | 123 | 107 | 72 | 67 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.6D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.60 | 0.57 | 0.59 | 0.56 | 0.49 | 0.47 |
| Marijuana and Hashish | 0.60 | 0.58 | 0.59 | 0.54 | 0.47 | 0.45 |
| Cocaine | 0.46 | 0.42 | 0.27 | 0.26 | 0.15 | 0.13 |
| Crack | 0.20 | 0.18 | 0.08 | 0.06 | 0.06 | 0.02 |
| Heroin | 0.15 | 0.17 | 0.08 | 0.09 | 0.06 | 0.07 |
| Hallucinogens | 0.52 | 0.46 | 0.33 | 0.28 | 0.17 | 0.14 |
| LSD | 0.33 | 0.30 | 0.18 | 0.15 | 0.06 | 0.05 |
| PCP | 0.14 | 0.14 | 0.03 | 0.06 | 0.02 | 0.03 |
| Ecstasy | 0.44 | 0.40 | 0.25 | 0.23 | 0.13 | 0.12 |
| Inhalants | 0.35 | 0.36 | 0.14 | 0.13 | 0.09 | 0.08 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.55 | 0.55 | 0.43 | 0.40 | 0.27 | 0.25 |
| Pain Relievers | 0.51 | 0.50 | 0.37 | 0.35 | 0.23 | 0.21 |
| OxyContin [®] | 0.30 | 0.28 | 0.16 | 0.15 | 0.09 | 0.09 |
| Tranquilizers | 0.43 | 0.40 | 0.27 | 0.23 | 0.16 | 0.15 |
| Stimulants ³ | 0.39 | 0.38 | 0.22 | 0.19 | 0.13 | 0.12 |
| Methamphetamine ³ | 0.27 | 0.23 | 0.10 | 0.09 | 0.05 | 0.06 |
| Sedatives | 0.14 | 0.15 | 0.08 | 0.06 | 0.04 | 0.04 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.60 | 0.58 | 0.50 | 0.45 | 0.32 | 0.30 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.6P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.8530 | | 0.5923 | | 0.4094 | |
| Marijuana and Hashish | 0.8904 | | 0.8211 | | 0.9782 | |
| Cocaine | 0.0078 | | 0.2536 | | 0.0344 | |
| Crack | 0.0095 | | 0.0126 | | 0.0007 | |
| Heroin | 0.2045 | | 0.3624 | | 0.4340 | |
| Hallucinogens | 0.0640 | | 0.1286 | | 0.1977 | |
| LSD | 0.1794 | | 0.5270 | | 0.7118 | |
| PCP | 0.0638 | | 0.1386 | | 0.6738 | |
| Ecstasy | 0.8368 | | 0.4277 | | 0.2780 | |
| Inhalants | 0.0835 | | 0.0891 | | 0.5084 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.0702 | | 0.0033 | | 0.0097 | |
| Pain Relievers | 0.0271 | | 0.0119 | | 0.0038 | |
| OxyContin [®] | 0.5736 | | 0.2248 | | 0.4926 | |
| Tranquilizers | 0.0857 | | 0.0016 | | 0.1403 | |
| Stimulants ³ | 0.0757 | | 0.0562 | | 0.4093 | |
| Methamphetamine ³ | 0.0211 | | 0.2339 | | 0.4955 | |
| Sedatives | 0.6684 | | 0.2183 | | 0.5948 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.0183 | | 0.0129 | | 0.0038 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.7C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 312 | 326 | 271 | 268 | 227 | 214 |
| Marijuana and Hashish | 318 | 330 | 247 | 244 | 199 | 195 |
| Cocaine | 242 | 248 | 121 | 90 | 66 | 47 |
| Crack | 132 | 113 | 35 | 20 | 9 | 11 |
| Heroin | 96 | 87 | 44 | 39 | 15 | 30 |
| Hallucinogens | 257 | 257 | 102 | 75 | 52 | 32 |
| LSD | 212 | 203 | 44 | 21 | 3 | 6 |
| PCP | 94 | 77 | 3 | 9 | 3 | 7 |
| Ecstasy | 233 | 222 | 81 | 63 | 41 | 26 |
| Inhalants | 203 | 184 | 46 | 45 | 24 | 29 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 281 | 287 | 178 | 176 | 123 | 120 |
| Pain Relievers | 259 | 266 | 173 | 150 | 113 | 104 |
| OxyContin [®] | 126 | 120 | 67 | 59 | 32 | 28 |
| Tranquilizers | 204 | 217 | 107 | 117 | 63 | 65 |
| Stimulants ³ | 203 | 196 | 69 | 70 | 49 | 47 |
| Methamphetamine ³ | 164 | 155 | 39 | 51 | 24 | 35 |
| Sedatives | 86 | 69 | 39 | 24 | 27 | 8 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 310 | 305 | 201 | 194 | 144 | 128 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.7D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.85 | 0.90 | 0.74 | 0.74 | 0.62 | 0.59 |
| Marijuana and Hashish | 0.87 | 0.91 | 0.68 | 0.67 | 0.54 | 0.54 |
| Cocaine | 0.66 | 0.68 | 0.33 | 0.25 | 0.18 | 0.13 |
| Crack | 0.36 | 0.31 | 0.10 | 0.06 | 0.03 | 0.03 |
| Heroin | 0.26 | 0.24 | 0.12 | 0.11 | 0.04 | 0.08 |
| Hallucinogens | 0.70 | 0.71 | 0.28 | 0.21 | 0.14 | 0.09 |
| LSD | 0.58 | 0.56 | 0.12 | 0.06 | 0.01 | 0.02 |
| PCP | 0.26 | 0.21 | 0.01 | 0.02 | 0.01 | 0.02 |
| Ecstasy | 0.64 | 0.61 | 0.22 | 0.17 | 0.11 | 0.07 |
| Inhalants | 0.56 | 0.51 | 0.13 | 0.12 | 0.07 | 0.08 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.77 | 0.79 | 0.49 | 0.48 | 0.34 | 0.33 |
| Pain Relievers | 0.71 | 0.73 | 0.47 | 0.41 | 0.31 | 0.29 |
| OxyContin [®] | 0.34 | 0.33 | 0.18 | 0.16 | 0.09 | 0.08 |
| Tranquilizers | 0.56 | 0.60 | 0.29 | 0.32 | 0.17 | 0.18 |
| Stimulants ³ | 0.56 | 0.54 | 0.19 | 0.19 | 0.14 | 0.13 |
| Methamphetamine ³ | 0.45 | 0.43 | 0.11 | 0.14 | 0.06 | 0.10 |
| Sedatives | 0.24 | 0.19 | 0.11 | 0.07 | 0.07 | 0.02 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.85 | 0.84 | 0.55 | 0.53 | 0.39 | 0.35 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.7P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.8271 | | 0.3230 | | 0.2752 | |
| Marijuana and Hashish | 0.7087 | | 0.2697 | | 0.6480 | |
| Cocaine | 0.0798 | | 0.0034 | | 0.1361 | |
| Crack | 0.0344 | | 0.1945 | | 0.8972 | |
| Heroin | 0.8423 | | 0.8446 | | 0.0928 | |
| Hallucinogens | 0.0309 | | 0.0190 | | 0.1120 | |
| LSD | 0.0107 | | 0.0849 | | 0.4038 | |
| PCP | 0.1034 | | 0.3028 | | 0.5923 | |
| Ecstasy | 0.0108 | | 0.1265 | | 0.3213 | |
| Inhalants | 0.0024 | | 0.8414 | | 0.8233 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.6996 | | 0.4498 | | 0.4092 | |
| Pain Relievers | 0.4425 | | 0.2075 | | 0.6038 | |
| OxyContin [®] | 0.2233 | | 0.4658 | | 0.3750 | |
| Tranquilizers | 0.9513 | | 0.8205 | | 0.5701 | |
| Stimulants ³ | 0.2312 | | 0.7442 | | 0.5782 | |
| Methamphetamine ³ | 0.1302 | | 0.2648 | | 0.4571 | |
| Sedatives | 0.0145 | | 0.0115 | | 0.0135 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.0902 | | 0.1545 | | 0.1770 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.8C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 904 | 936 | 482 | 453 | 368 | 348 |
| Marijuana and Hashish | 906 | 918 | 399 | 367 | 301 | 300 |
| Cocaine | 621 | 642 | 136 | 123 | 75 | 84 |
| Crack | 340 | 302 | 79 | 64 | 57 | 40 |
| Heroin | 222 | 203 | 55 | 45 | 37 | 26 |
| Hallucinogens | 574 | 553 | 75 | 75 | 33 | 21 |
| LSD | 488 | 471 | 17 | 28 | * | 15 |
| PCP | 300 | 293 | 20 | * | 20 | * |
| Ecstasy | 296 | 243 | 38 | 47 | 18 | 13 |
| Inhalants | 409 | 436 | 89 | 75 | 74 | 46 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 674 | 670 | 316 | 285 | 217 | 177 |
| Pain Relievers | 487 | 506 | 251 | 232 | 160 | 142 |
| OxyContin [®] | 184 | 149 | 80 | 68 | 54 | 38 |
| Tranquilizers | 422 | 414 | 164 | 161 | 117 | 101 |
| Stimulants ³ | 492 | 448 | 113 | 111 | 60 | 66 |
| Methamphetamine ³ | 404 | 365 | 85 | 96 | 43 | 60 |
| Sedatives | 340 | 343 | 93 | 54 | 66 | 47 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 780 | 807 | 335 | 312 | 234 | 197 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.8D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.57 | 0.58 | 0.30 | 0.28 | 0.23 | 0.21 |
| Marijuana and Hashish | 0.57 | 0.57 | 0.25 | 0.23 | 0.19 | 0.19 |
| Cocaine | 0.39 | 0.40 | 0.09 | 0.08 | 0.05 | 0.05 |
| Crack | 0.21 | 0.19 | 0.05 | 0.04 | 0.04 | 0.02 |
| Heroin | 0.14 | 0.13 | 0.03 | 0.03 | 0.02 | 0.02 |
| Hallucinogens | 0.36 | 0.34 | 0.05 | 0.05 | 0.02 | 0.01 |
| LSD | 0.31 | 0.29 | 0.01 | 0.02 | * | 0.01 |
| PCP | 0.19 | 0.18 | 0.01 | * | 0.01 | * |
| Ecstasy | 0.19 | 0.15 | 0.02 | 0.03 | 0.01 | 0.01 |
| Inhalants | 0.26 | 0.27 | 0.06 | 0.05 | 0.05 | 0.03 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.42 | 0.41 | 0.20 | 0.18 | 0.14 | 0.11 |
| Pain Relievers | 0.31 | 0.31 | 0.16 | 0.14 | 0.10 | 0.09 |
| OxyContin [®] | 0.12 | 0.09 | 0.05 | 0.04 | 0.03 | 0.02 |
| Tranquilizers | 0.27 | 0.26 | 0.10 | 0.10 | 0.07 | 0.06 |
| Stimulants ³ | 0.31 | 0.28 | 0.07 | 0.07 | 0.04 | 0.04 |
| Methamphetamine ³ | 0.25 | 0.23 | 0.05 | 0.06 | 0.03 | 0.04 |
| Sedatives | 0.21 | 0.21 | 0.06 | 0.03 | 0.04 | 0.03 |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.49 | 0.50 | 0.21 | 0.19 | 0.15 | 0.12 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.8P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.8468 | | 0.3942 | | 0.8987 | |
| Marijuana and Hashish | 0.7812 | | 0.7801 | | 0.4681 | |
| Cocaine | 0.9905 | | 0.2801 | | 0.7295 | |
| Crack | 0.1861 | | 0.1438 | | 0.1409 | |
| Heroin | 0.8434 | | 0.5937 | | 0.3396 | |
| Hallucinogens | 0.5283 | | 0.7062 | | 0.5307 | |
| LSD | 0.8009 | | 0.4902 | | * | |
| PCP | 0.9214 | | * | | * | |
| Ecstasy | 0.1002 | | 0.3442 | | 0.9022 | |
| Inhalants | 0.7092 | | 0.8726 | | 0.5822 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.6355 | | 0.0925 | | 0.1389 | |
| Pain Relievers | 0.9546 | | 0.2321 | | 0.2638 | |
| OxyContin [®] | 1.0000 | | 0.5614 | | 0.7632 | |
| Tranquilizers | 0.4395 | | 0.3134 | | 0.0613 | |
| Stimulants ³ | 0.1765 | | 0.9078 | | 0.9916 | |
| Methamphetamine ³ | 0.1367 | | 0.8030 | | 0.5509 | |
| Sedatives | 0.9307 | | 0.0568 | | 0.5285 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.6837 | | 0.1032 | | 0.2065 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.9C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 704 | 664 | 495 | 430 | 386 | 340 |
| Marijuana and Hashish | 697 | 667 | 429 | 386 | 329 | 319 |
| Cocaine | 527 | 522 | 173 | 142 | 91 | 88 |
| Crack | 299 | 281 | 65 | 61 | 44 | 34 |
| Heroin | 223 | 179 | 63 | 59 | 37 | 29 |
| Hallucinogens | 531 | 462 | 143 | 117 | 67 | 47 |
| LSD | 451 | 402 | 56 | 47 | 19 | 15 |
| PCP | 274 | 253 | 23 | 17 | 20 | 9 |
| Ecstasy | 351 | 260 | 101 | 83 | 49 | 35 |
| Inhalants | 385 | 379 | 103 | 91 | 79 | 57 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 575 | 518 | 312 | 250 | 200 | 165 |
| Pain Relievers | 446 | 436 | 272 | 218 | 171 | 145 |
| OxyContin [®] | 199 | 164 | 84 | 86 | 43 | 44 |
| Tranquilizers | 374 | 341 | 162 | 133 | 92 | 79 |
| Stimulants ³ | 425 | 364 | 116 | 117 | 64 | 74 |
| Methamphetamine ³ | 362 | 318 | 89 | 104 | 42 | 60 |
| Sedatives | 281 | 243 | 87 | 34 | 68 | 20 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 664 | 609 | 349 | 289 | 233 | 187 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.9D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.57 | 0.53 | 0.40 | 0.34 | 0.31 | 0.27 |
| Marijuana and Hashish | 0.56 | 0.53 | 0.35 | 0.31 | 0.27 | 0.26 |
| Cocaine | 0.43 | 0.42 | 0.14 | 0.11 | 0.07 | 0.07 |
| Crack | 0.24 | 0.23 | 0.05 | 0.05 | 0.04 | 0.03 |
| Heroin | 0.18 | 0.14 | 0.05 | 0.05 | 0.03 | 0.02 |
| Hallucinogens | 0.43 | 0.37 | 0.12 | 0.09 | 0.05 | 0.04 |
| LSD | 0.37 | 0.32 | 0.05 | 0.04 | 0.02 | 0.01 |
| PCP | 0.22 | 0.20 | 0.02 | 0.01 | 0.02 | 0.01 |
| Ecstasy | 0.28 | 0.21 | 0.08 | 0.07 | 0.04 | 0.03 |
| Inhalants | 0.31 | 0.30 | 0.08 | 0.07 | 0.06 | 0.05 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.47 | 0.42 | 0.25 | 0.20 | 0.16 | 0.13 |
| Pain Relievers | 0.36 | 0.35 | 0.22 | 0.18 | 0.14 | 0.12 |
| OxyContin [®] | 0.16 | 0.13 | 0.07 | 0.07 | 0.04 | 0.04 |
| Tranquilizers | 0.30 | 0.27 | 0.13 | 0.11 | 0.07 | 0.06 |
| Stimulants ³ | 0.34 | 0.29 | 0.09 | 0.09 | 0.05 | 0.06 |
| Methamphetamine ³ | 0.29 | 0.25 | 0.07 | 0.08 | 0.03 | 0.05 |
| Sedatives | 0.23 | 0.19 | 0.07 | 0.03 | 0.06 | 0.02 |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.54 | 0.49 | 0.28 | 0.23 | 0.19 | 0.15 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.9P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.1994 | | 0.2385 | | 0.7405 | |
| Marijuana and Hashish | 0.3014 | | 0.8277 | | 0.5828 | |
| Cocaine | 0.5407 | | 0.0124 | | 0.3568 | |
| Crack | 0.3834 | | 0.3809 | | 0.2035 | |
| Heroin | 0.4263 | | 0.7569 | | 0.8195 | |
| Hallucinogens | 0.1022 | | 0.0345 | | 0.0716 | |
| LSD | 0.5290 | | 0.6415 | | 0.4922 | |
| PCP | 0.5669 | | 0.5817 | | 0.7453 | |
| Ecstasy | 0.0004 | | 0.1160 | | 0.1233 | |
| Inhalants | 0.0282 | | 0.6595 | | 0.7106 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.2818 | | 0.0126 | | 0.0685 | |
| Pain Relievers | 0.6130 | | 0.0061 | | 0.1020 | |
| OxyContin [®] | 0.3795 | | 0.5601 | | 0.9966 | |
| Tranquilizers | 0.1135 | | 0.0885 | | 0.1405 | |
| Stimulants ³ | 0.2024 | | 0.7018 | | 0.7500 | |
| Methamphetamine ³ | 0.1827 | | 0.6035 | | 0.4947 | |
| Sedatives | 0.3791 | | 0.0001 | | 0.0058 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.1569 | | 0.0072 | | 0.0670 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.10C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 662 | 687 | 367 | 368 | 261 | 260 |
| Marijuana and Hashish | 648 | 658 | 303 | 305 | 206 | 215 |
| Cocaine | 397 | 424 | 101 | 87 | 59 | 54 |
| Crack | 192 | 166 | 64 | 36 | 42 | 24 |
| Heroin | 115 | 134 | 41 | 38 | 22 | 35 |
| Hallucinogens | 387 | 369 | 92 | 77 | 48 | 37 |
| LSD | 311 | 308 | 38 | 34 | 7 | 17 |
| PCP | 157 | 165 | 9 | 11 | 4 | * |
| Ecstasy | 228 | 238 | 71 | 67 | 34 | 30 |
| Inhalants | 257 | 265 | 54 | 54 | 28 | 30 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 502 | 496 | 242 | 242 | 167 | 152 |
| Pain Relievers | 383 | 360 | 199 | 201 | 129 | 120 |
| OxyContin [®] | 130 | 116 | 80 | 57 | 55 | 29 |
| Tranquilizers | 329 | 336 | 144 | 159 | 103 | 103 |
| Stimulants ³ | 339 | 317 | 92 | 87 | 58 | 48 |
| Methamphetamine ³ | 252 | 236 | 43 | 51 | 32 | 39 |
| Sedatives | 207 | 239 | 57 | 52 | 25 | 44 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 567 | 578 | 266 | 260 | 180 | 161 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.10D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2010 and 2011

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLICIT DRUGS¹ | 0.51 | 0.52 | 0.28 | 0.28 | 0.20 | 0.20 |
| Marijuana and Hashish | 0.50 | 0.49 | 0.23 | 0.23 | 0.16 | 0.16 |
| Cocaine | 0.31 | 0.32 | 0.08 | 0.07 | 0.04 | 0.04 |
| Crack | 0.15 | 0.12 | 0.05 | 0.03 | 0.03 | 0.02 |
| Heroin | 0.09 | 0.10 | 0.03 | 0.03 | 0.02 | 0.03 |
| Hallucinogens | 0.30 | 0.28 | 0.07 | 0.06 | 0.04 | 0.03 |
| LSD | 0.24 | 0.23 | 0.03 | 0.03 | 0.01 | 0.01 |
| PCP | 0.12 | 0.12 | 0.01 | 0.01 | 0.00 | * |
| Ecstasy | 0.18 | 0.18 | 0.05 | 0.05 | 0.03 | 0.02 |
| Inhalants | 0.20 | 0.20 | 0.04 | 0.04 | 0.02 | 0.02 |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.39 | 0.37 | 0.19 | 0.18 | 0.13 | 0.11 |
| Pain Relievers | 0.29 | 0.27 | 0.15 | 0.15 | 0.10 | 0.09 |
| OxyContin [®] | 0.10 | 0.09 | 0.06 | 0.04 | 0.04 | 0.02 |
| Tranquilizers | 0.25 | 0.25 | 0.11 | 0.12 | 0.08 | 0.08 |
| Stimulants ³ | 0.26 | 0.24 | 0.07 | 0.07 | 0.04 | 0.04 |
| Methamphetamine ³ | 0.19 | 0.18 | 0.03 | 0.04 | 0.02 | 0.03 |
| Sedatives | 0.16 | 0.18 | 0.04 | 0.04 | 0.02 | 0.03 |
| ILLICIT DRUGS OTHER THAN MARIJUANA¹ | 0.44 | 0.43 | 0.20 | 0.20 | 0.14 | 0.12 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.10P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Lifetime (2010) | Lifetime (2011) | Past Year (2010) | Past Year (2011) | Past Month (2010) | Past Month (2011) |
|--|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| ILLCIT DRUGS¹ | 0.5664 | | 0.4391 | | 0.3482 | |
| Marijuana and Hashish | 0.5197 | | 0.7324 | | 0.5687 | |
| Cocaine | 0.3597 | | 0.2528 | | 0.9485 | |
| Crack | 0.0027 | | 0.0112 | | 0.0973 | |
| Heroin | 0.2823 | | 0.7474 | | 0.2329 | |
| Hallucinogens | 0.2880 | | 0.2844 | | 0.0954 | |
| LSD | 0.4542 | | 0.6827 | | 0.5210 | |
| PCP | 0.9370 | | 0.6814 | | * | |
| Ecstasy | 0.9046 | | 0.5668 | | 0.1821 | |
| Inhalants | 0.3105 | | 0.0450 | | 0.3391 | |
| Nonmedical Use of Psychotherapeutics ^{2,3} | 0.4491 | | 0.0795 | | 0.0432 | |
| Pain Relievers | 0.1082 | | 0.3046 | | 0.0700 | |
| OxyContin [®] | 0.5564 | | 0.0329 | | 0.0252 | |
| Tranquilizers | 0.8291 | | 0.3650 | | 0.2477 | |
| Stimulants ³ | 0.0342 | | 0.1545 | | 0.2179 | |
| Methamphetamine ³ | 0.0420 | | 0.8719 | | 0.5757 | |
| Sedatives | 0.8013 | | 0.6530 | | 0.3450 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.2725 | | 0.0612 | | 0.0206 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.