

Table 1.49C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	784	752	397	363	268	231
AGE						
12-17	68	64	58	53	38	35
18-25	143	145	114	110	72	69
26 or Older	752	740	363	338	252	215
GENDER						
Male	575	518	312	250	200	165
Female	502	496	242	242	167	152
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	730	702	365	336	254	207
White	651	634	336	307	234	186
Black or African American	210	208	107	106	75	52
American Indian or Alaska Native	41	51	26	35	18	25
Native Hawaiian or Other Pacific Islander	19	*	12	16	11	15
Asian	144	111	107	53	30	38
Two or More Races	104	107	29	33	17	25
Hispanic or Latino	250	258	150	135	89	96

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.49D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.31	0.29	0.16	0.14	0.11	0.09
AGE						
12-17	0.28	0.26	0.24	0.21	0.16	0.14
18-25	0.42	0.42	0.33	0.32	0.21	0.20
26 or Older	0.39	0.37	0.19	0.17	0.13	0.11
GENDER						
Male	0.47	0.42	0.25	0.20	0.16	0.13
Female	0.39	0.37	0.19	0.18	0.13	0.11
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.34	0.32	0.17	0.15	0.12	0.09
White	0.38	0.37	0.20	0.18	0.14	0.11
Black or African American	0.70	0.69	0.35	0.35	0.25	0.17
American Indian or Alaska Native	3.05	3.19	2.12	2.34	1.48	1.63
Native Hawaiian or Other Pacific Islander	2.57	*	1.70	1.75	1.53	1.65
Asian	1.19	0.87	0.88	0.42	0.27	0.30
Two or More Races	2.71	2.31	0.97	0.84	0.54	0.63
Hispanic or Latino	0.68	0.65	0.41	0.34	0.24	0.24

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.49P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.2064		0.0026		0.0078	
AGE						
12-17	0.1482		0.1571		0.3216	
18-25	0.0066		0.0004		0.0024	
26 or Older	0.5191		0.0522		0.0776	
GENDER						
Male	0.2818		0.0126		0.0685	
Female	0.4491		0.0795		0.0432	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3076		0.0037		0.0133	
White	0.3784		0.0137		0.0161	
Black or African American	0.5762		0.6726		0.2551	
American Indian or Alaska Native	0.7365		0.8577		0.4623	
Native Hawaiian or Other Pacific Islander	*		0.6775		0.5746	
Asian	0.1115		0.0589		0.8499	
Two or More Races	0.3463		0.2711		0.9227	
Hispanic or Latino	0.5336		0.3511		0.2976	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.50C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	68	64	58	53	38	35
GENDER						
Male	48	46	41	37	24	24
Female	48	46	42	38	30	24
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	58	54	50	45	33	30
White	58	50	49	42	29	24
Black or African American	28	28	20	21	13	13
American Indian or Alaska Native	5	6	4	6	2	4
Native Hawaiian or Other Pacific Islander	2	*	*	*	1	*
Asian	13	13	10	10	8	6
Two or More Races	12	14	11	11	8	8
Hispanic or Latino	36	35	32	28	20	16
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	36	34	31	28	19	17
Female, White, Not Hispanic	41	34	35	30	22	18
Male, Black, Not Hispanic	20	18	13	15	7	8
Female, Black, Not Hispanic	19	20	16	13	11	10
Male, Hispanic	29	25	25	19	14	11
Female, Hispanic	25	27	20	21	15	12

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.50D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.28	0.26	0.24	0.21	0.16	0.14
GENDER						
Male	0.39	0.36	0.33	0.29	0.19	0.18
Female	0.40	0.37	0.35	0.31	0.25	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.30	0.28	0.26	0.23	0.17	0.16
White	0.35	0.32	0.31	0.28	0.19	0.17
Black or African American	0.70	0.73	0.51	0.55	0.35	0.37
American Indian or Alaska Native	3.39	3.46	2.91	3.40	1.29	2.83
Native Hawaiian or Other Pacific Islander	3.53	*	*	*	1.02	*
Asian	1.13	1.02	0.88	0.80	0.76	0.48
Two or More Races	1.97	1.81	1.84	1.52	1.52	1.07
Hispanic or Latino	0.74	0.65	0.65	0.51	0.41	0.30
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.45	0.45	0.41	0.37	0.25	0.23
Female, White, Not Hispanic	0.53	0.46	0.47	0.42	0.32	0.26
Male, Black, Not Hispanic	1.03	0.98	0.67	0.83	0.40	0.46
Female, Black, Not Hispanic	1.00	1.05	0.85	0.70	0.59	0.53
Male, Hispanic	1.01	0.85	0.92	0.67	0.54	0.39
Female, Hispanic	0.97	0.93	0.80	0.76	0.62	0.45

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.50P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.1482		0.1571		0.3216	
GENDER						
Male	0.8553		0.9580		0.6934	
Female	0.0651		0.0505		0.0959	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4573		0.5984		0.9247	
White	0.3472		0.4525		0.9125	
Black or African American	0.6263		0.4732		0.8954	
American Indian or Alaska Native	1.0000		0.5431		0.1768	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6739		0.6206		0.4535	
Two or More Races	0.4837		0.6107		0.5954	
Hispanic or Latino	0.1336		0.0868		0.0818	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.3624		0.4451		0.2628	
Female, White, Not Hispanic	0.0270		0.0882		0.4067	
Male, Black, Not Hispanic	0.3787		0.1143		0.8197	
Female, Black, Not Hispanic	0.8466		0.5694		0.9972	
Male, Hispanic	0.0409		0.0814		0.2463	
Female, Hispanic	0.8659		0.4393		0.1754	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.51C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	143	145	114	110	72	69
GENDER						
Male	105	104	88	79	56	50
Female	95	99	71	71	47	45
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	130	129	104	98	64	63
White	165	151	117	101	64	61
Black or African American	50	51	34	30	23	17
American Indian or Alaska Native	10	14	7	8	7	4
Native Hawaiian or Other Pacific Islander	*	*	5	*	*	*
Asian	25	31	18	18	10	11
Two or More Races	21	22	13	16	5	12
Hispanic or Latino	63	69	45	46	28	27
EDUCATION						
< High School	67	67	46	46	33	31
High School Graduate	101	99	67	68	45	41
Some College	115	121	82	78	44	46
College Graduate	70	68	52	37	26	23
CURRENT EMPLOYMENT						
Full-Time	112	112	81	70	52	41
Part-Time	96	91	64	61	35	40
Unemployed	61	59	42	44	32	30
Other ¹	80	79	56	55	33	34

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.51D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.42	0.42	0.33	0.32	0.21	0.20
GENDER						
Male	0.61	0.61	0.51	0.46	0.32	0.29
Female	0.56	0.58	0.43	0.42	0.28	0.26
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.47	0.38	0.36	0.23	0.23
White	0.55	0.56	0.45	0.43	0.29	0.28
Black or African American	0.91	0.92	0.68	0.59	0.48	0.36
American Indian or Alaska Native	4.35	4.43	3.58	3.36	3.35	1.78
Native Hawaiian or Other Pacific Islander	*	*	3.13	*	*	*
Asian	1.64	1.55	1.18	0.93	0.72	0.57
Two or More Races	3.29	2.74	2.29	2.12	0.99	1.63
Hispanic or Latino	0.96	0.98	0.68	0.65	0.43	0.38
EDUCATION						
< High School	0.95	0.94	0.73	0.73	0.53	0.52
High School Graduate	0.70	0.69	0.51	0.52	0.37	0.33
Some College	0.72	0.70	0.58	0.53	0.35	0.35
College Graduate	1.11	1.13	0.97	0.69	0.51	0.45
CURRENT EMPLOYMENT						
Full-Time	0.68	0.72	0.57	0.50	0.39	0.30
Part-Time	0.81	0.74	0.60	0.57	0.36	0.41
Unemployed	1.15	1.15	0.87	0.89	0.67	0.64
Other ¹	0.83	0.78	0.63	0.61	0.41	0.40

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.51P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0066		0.0004		0.0024	
GENDER						
Male	0.0776		0.0399		0.0441	
Female	0.0402		0.0020		0.0199	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0104		0.0002		0.0072	
White	0.0028		0.0001		0.0044	
Black or African American	0.0873		0.2619		0.7519	
American Indian or Alaska Native	0.2240		0.9269		0.1085	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7098		0.2892		0.8342	
Two or More Races	0.1736		0.6774		0.1978	
Hispanic or Latino	0.6494		0.8648		0.2263	
EDUCATION						
< High School	0.6124		0.1929		0.2118	
High School Graduate	0.2361		0.4664		0.0746	
Some College	0.2612		0.0404		0.0943	
College Graduate	0.0029		0.0001		0.1792	
CURRENT EMPLOYMENT						
Full-Time	0.2229		0.0166		0.0003	
Part-Time	0.1229		0.0257		0.7027	
Unemployed	0.8836		0.9136		0.4565	
Other ¹	0.0131		0.0394		0.2134	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.52C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	752	740	363	338	252	215
GENDER						
Male	559	514	288	234	189	155
Female	480	484	221	225	155	141
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	704	687	336	309	240	192
White	752	743	318	293	223	178
Black or African American	222	223	102	101	70	48
American Indian or Alaska Native	38	48	24	33	16	23
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	138	103	105	48	27	36
Two or More Races	103	103	23	25	13	18
Hispanic or Latino	231	245	135	126	81	91
EDUCATION						
< High School	265	241	135	121	84	88
High School Graduate	419	392	188	167	115	117
Some College	443	452	193	183	133	111
College Graduate	466	496	212	192	146	104
CURRENT EMPLOYMENT						
Full-Time	623	635	270	247	188	161
Part-Time	274	266	114	141	73	78
Unemployed	205	178	131	96	69	67
Other ¹	391	384	183	153	110	101

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.52D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.39	0.37	0.19	0.17	0.13	0.11
GENDER						
Male	0.60	0.54	0.31	0.25	0.20	0.16
Female	0.47	0.47	0.22	0.22	0.15	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.40	0.20	0.18	0.14	0.11
White	0.47	0.46	0.23	0.21	0.16	0.13
Black or African American	0.94	0.93	0.46	0.44	0.32	0.22
American Indian or Alaska Native	3.96	4.07	2.64	2.99	1.83	2.11
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.47	1.09	1.11	0.51	0.31	0.38
Two or More Races	3.97	3.44	1.12	0.97	0.63	0.72
Hispanic or Latino	0.92	0.91	0.53	0.47	0.32	0.34
EDUCATION						
< High School	0.89	0.82	0.47	0.43	0.29	0.32
High School Graduate	0.64	0.62	0.32	0.28	0.20	0.20
Some College	0.82	0.78	0.39	0.36	0.28	0.22
College Graduate	0.67	0.69	0.34	0.30	0.24	0.16
CURRENT EMPLOYMENT						
Full-Time	0.52	0.50	0.26	0.23	0.18	0.15
Part-Time	1.03	1.01	0.48	0.59	0.31	0.33
Unemployed	1.65	1.69	1.18	1.02	0.65	0.71
Other ¹	0.62	0.58	0.30	0.24	0.18	0.16

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.52P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.5191		0.0522		0.0776	
GENDER						
Male	0.4854		0.0434		0.1718	
Female	0.8434		0.5472		0.2548	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6112		0.0587		0.0757	
White	0.7983		0.2201		0.0985	
Black or African American	0.8740		0.3916		0.2964	
American Indian or Alaska Native	0.5310		0.9339		0.2802	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0969		0.0790		0.8443	
Two or More Races	0.5957		0.2524		0.6761	
Hispanic or Latino	0.7607		0.5626		0.7301	
EDUCATION						
< High School	1.0000		0.8850		0.6180	
High School Graduate	0.5577		0.8189		0.9963	
Some College	0.4849		0.0800		0.2229	
College Graduate	0.7938		0.0958		0.0206	
CURRENT EMPLOYMENT						
Full-Time	0.5266		0.0537		0.0987	
Part-Time	0.6047		0.3639		0.8755	
Unemployed	0.9959		0.9817		0.7238	
Other ¹	0.6604		0.1661		0.4242	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.53C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	777	752	389	357	265	228
GENDER						
Male	571	518	307	246	199	164
Female	496	496	236	238	163	150
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	726	702	359	330	251	204
White	790	771	346	314	236	187
Black or African American	227	230	106	107	73	51
American Indian or Alaska Native	40	50	25	35	18	24
Native Hawaiian or Other Pacific Islander	19	*	12	16	*	15
Asian	144	109	107	52	29	37
Two or More Races	105	105	26	31	14	23
Hispanic or Latino	246	254	145	132	87	95
EDUCATION						
< High School	274	252	143	130	90	94
High School Graduate	439	407	200	182	125	124
Some College	462	478	212	202	142	124
College Graduate	479	506	219	196	148	106
CURRENT EMPLOYMENT						
Full-Time	641	646	283	254	197	165
Part-Time	287	278	125	151	79	87
Unemployed	220	188	141	106	77	73
Other ¹	402	393	192	164	116	108

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.53D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.34	0.32	0.17	0.15	0.12	0.10
GENDER						
Male	0.51	0.46	0.28	0.22	0.18	0.15
Female	0.42	0.41	0.20	0.20	0.14	0.12
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.35	0.18	0.17	0.13	0.10
White	0.42	0.41	0.21	0.19	0.15	0.12
Black or African American	0.78	0.77	0.40	0.38	0.28	0.19
American Indian or Alaska Native	3.42	3.52	2.33	2.56	1.66	1.76
Native Hawaiian or Other Pacific Islander	2.82	*	1.86	1.86	*	1.76
Asian	1.31	0.95	0.97	0.46	0.28	0.33
Two or More Races	3.23	2.76	1.07	0.94	0.55	0.71
Hispanic or Latino	0.77	0.75	0.46	0.39	0.27	0.28
EDUCATION						
< High School	0.77	0.71	0.42	0.38	0.26	0.28
High School Graduate	0.55	0.53	0.28	0.26	0.18	0.18
Some College	0.68	0.65	0.34	0.31	0.24	0.20
College Graduate	0.63	0.65	0.33	0.28	0.22	0.15
CURRENT EMPLOYMENT						
Full-Time	0.47	0.45	0.24	0.21	0.17	0.14
Part-Time	0.76	0.74	0.38	0.45	0.24	0.27
Unemployed	1.21	1.19	0.87	0.75	0.50	0.52
Other ¹	0.57	0.53	0.28	0.23	0.17	0.15

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.53P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.2604		0.0039		0.0105	
GENDER						
Male	0.3019		0.0107		0.0586	
Female	0.5612		0.1355		0.0755	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3331		0.0036		0.0127	
White	0.4046		0.0157		0.0145	
Black or African American	0.6180		0.5773		0.2564	
American Indian or Alaska Native	0.7163		0.9234		0.6040	
Native Hawaiian or Other Pacific Islander	*		0.7123		*	
Asian	0.1187		0.0643		0.9686	
Two or More Races	0.4212		0.2850		0.7684	
Hispanic or Latino	0.6920		0.5577		0.4820	
EDUCATION						
< High School	0.9558		0.8456		0.9072	
High School Graduate	0.4147		0.6470		0.5637	
Some College	0.3580		0.0212		0.0922	
College Graduate	0.5202		0.0169		0.0119	
CURRENT EMPLOYMENT						
Full-Time	0.4061		0.0133		0.0124	
Part-Time	0.3487		0.9854		0.7430	
Unemployed	0.9393		0.9492		0.6042	
Other ¹	0.9774		0.0638		0.2647	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.