

Table 1.76C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	473	491	115	107	83	68
GEOGRAPHIC DIVISION						
Northeast	205	257	55	41	34	23
New England	114	111	18	20	12	12
Middle Atlantic	169	232	52	35	32	19
Midwest	157	165	33	40	20	30
East North Central	134	140	30	23	19	10
West North Central	90	93	15	33	5	28
South	273	287	77	45	69	34
South Atlantic	213	220	73	38	68	29
East South Central	97	86	17	12	8	7
West South Central	140	152	24	22	10	16
West	318	279	55	76	26	46
Mountain	145	149	20	43	13	23
Pacific	285	236	52	63	22	39
COUNTY TYPE						
Large Metro	414	417	103	75	77	43
Small Metro	298	278	47	75	30	51
250K – 1 Mil. Pop.	250	253	41	65	26	47
< 250K Pop.	165	125	24	36	16	21
Nonmetro	195	142	23	17	10	9
Urbanized	133	101	13	15	7	10
Less Urbanized	146	97	16	12	7	7
Completely Rural	42	28	9	4	3	1

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.76D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.19	0.19	0.05	0.04	0.03	0.03
GEOGRAPHIC DIVISION						
Northeast	0.44	0.55	0.12	0.09	0.07	0.05
New England	0.93	0.90	0.15	0.16	0.10	0.10
Middle Atlantic	0.49	0.67	0.15	0.10	0.09	0.06
Midwest	0.28	0.30	0.06	0.07	0.04	0.05
East North Central	0.35	0.36	0.08	0.06	0.05	0.03
West North Central	0.53	0.55	0.09	0.19	0.03	0.17
South	0.29	0.30	0.08	0.05	0.07	0.04
South Atlantic	0.43	0.44	0.15	0.08	0.14	0.06
East South Central	0.65	0.56	0.11	0.08	0.05	0.05
West South Central	0.49	0.51	0.08	0.07	0.03	0.06
West	0.54	0.47	0.09	0.13	0.04	0.08
Mountain	0.81	0.82	0.11	0.24	0.07	0.13
Pacific	0.70	0.57	0.13	0.15	0.05	0.09
COUNTY TYPE						
Large Metro	0.27	0.29	0.08	0.05	0.06	0.03
Small Metro	0.33	0.31	0.06	0.09	0.04	0.06
250K – 1 Mil. Pop.	0.43	0.40	0.08	0.12	0.05	0.09
< 250K Pop.	0.52	0.47	0.09	0.14	0.06	0.08
Nonmetro	0.44	0.33	0.06	0.04	0.03	0.02
Urbanized	0.66	0.55	0.07	0.09	0.04	0.06
Less Urbanized	0.66	0.45	0.08	0.06	0.04	0.03
Completely Rural	1.05	0.65	0.25	0.09	0.07	0.02

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.76P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0190		0.1784		0.4747	
GEOGRAPHIC DIVISION						
Northeast	0.5285		0.1307		0.3552	
New England	0.7740		0.9712		0.8377	
Middle Atlantic	0.5744		0.1010		0.3546	
Midwest	0.2250		0.3549		0.9513	
East North Central	0.2500		0.0274		0.0234	
West North Central	0.6921		0.3425		0.0957	
South	0.0566		0.0981		0.2901	
South Atlantic	0.0399		0.2584		0.2251	
East South Central	0.9128		0.0449		0.7461	
West South Central	0.5610		0.6841		0.5592	
West	0.0761		0.4154		0.3745	
Mountain	0.2171		0.1543		0.9432	
Pacific	0.1740		0.8569		0.2860	
COUNTY TYPE						
Large Metro	0.0997		0.0690		0.0834	
Small Metro	0.2198		0.4605		0.2504	
250K – 1 Mil. Pop.	0.5390		0.3856		0.2272	
< 250K Pop.	0.1436		0.8966		0.9340	
Nonmetro	0.2636		0.0553		0.6434	
Urbanized	0.3463		0.7679		0.2284	
Less Urbanized	0.5967		0.1120		0.7419	
Completely Rural	0.6852		0.0788		0.0591	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.77C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	61	58	42	40	23	20
GEOGRAPHIC DIVISION						
Northeast	23	22	16	18	9	8
New England	11	9	8	6	5	3
Middle Atlantic	20	20	13	17	7	8
Midwest	23	23	16	15	9	7
East North Central	20	17	14	10	9	5
West North Central	12	14	8	10	3	4
South	36	32	25	20	14	11
South Atlantic	26	22	18	14	10	7
East South Central	13	11	10	8	6	6
West South Central	19	20	12	12	8	6
West	38	37	26	27	13	13
Mountain	16	15	10	12	4	3
Pacific	35	34	24	23	13	12
COUNTY TYPE						
Large Metro	52	52	35	34	19	15
Small Metro	37	31	22	20	13	11
250K – 1 Mil. Pop.	31	25	18	18	10	10
< 250K Pop.	21	19	13	10	9	4
Nonmetro	21	23	15	13	6	8
Urbanized	13	15	10	9	3	5
Less Urbanized	15	18	11	9	5	6
Completely Rural	7	6	5	3	2	1

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.77D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.25	0.23	0.17	0.16	0.10	0.08
GEOGRAPHIC DIVISION						
Northeast	0.55	0.52	0.37	0.43	0.21	0.19
New England	1.03	0.79	0.74	0.58	0.43	0.28
Middle Atlantic	0.64	0.64	0.43	0.54	0.24	0.24
Midwest	0.42	0.42	0.29	0.27	0.17	0.12
East North Central	0.53	0.46	0.36	0.27	0.24	0.13
West North Central	0.74	0.87	0.50	0.60	0.18	0.27
South	0.40	0.34	0.27	0.21	0.16	0.12
South Atlantic	0.58	0.48	0.40	0.29	0.23	0.16
East South Central	0.93	0.73	0.70	0.56	0.43	0.38
West South Central	0.62	0.65	0.41	0.39	0.25	0.20
West	0.66	0.62	0.44	0.44	0.23	0.21
Mountain	0.90	0.81	0.58	0.67	0.24	0.17
Pacific	0.87	0.82	0.59	0.56	0.31	0.29
COUNTY TYPE						
Large Metro	0.37	0.35	0.26	0.25	0.14	0.11
Small Metro	0.44	0.36	0.28	0.26	0.17	0.14
250K – 1 Mil. Pop.	0.54	0.42	0.33	0.32	0.19	0.19
< 250K Pop.	0.70	0.64	0.52	0.39	0.34	0.16
Nonmetro	0.50	0.54	0.38	0.32	0.15	0.21
Urbanized	0.75	0.79	0.58	0.50	0.19	0.32
Less Urbanized	0.72	0.88	0.54	0.49	0.23	0.34
Completely Rural	1.98	1.63	1.61	0.75	0.64	0.26

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.77P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0271		0.1489		0.1453	
GEOGRAPHIC DIVISION						
Northeast	0.1691		0.7964		0.5538	
New England	0.0351		0.4052		0.0903	
Middle Atlantic	0.5832		0.9240		0.8196	
Midwest	0.1235		0.1426		0.1350	
East North Central	0.0255		0.0329		0.0185	
West North Central	0.6647		0.7402		0.1524	
South	0.3504		0.0426		0.1152	
South Atlantic	0.1569		0.0514		0.1868	
East South Central	0.0281		0.0137		0.6469	
West South Central	0.0970		0.4793		0.4217	
West	0.3800		0.6984		0.7426	
Mountain	0.0810		0.7578		0.5162	
Pacific	0.8083		0.7754		0.5875	
COUNTY TYPE						
Large Metro	0.1768		0.5203		0.0559	
Small Metro	0.0107		0.4711		0.7758	
250K – 1 Mil. Pop.	0.0049		0.8266		0.5628	
< 250K Pop.	0.5304		0.0914		0.1210	
Nonmetro	0.6727		0.0762		0.5385	
Urbanized	0.7632		0.3185		0.2050	
Less Urbanized	0.2695		0.4438		0.7221	
Completely Rural	0.5551		0.0483		0.1077	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.