

**Table 1.78C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	784	752	397	363	268	231
<b>GEOGRAPHIC DIVISION</b>						
Northeast	285	321	149	139	117	91
New England	146	147	61	66	49	42
Middle Atlantic	245	284	135	122	106	81
Midwest	275	273	133	133	94	95
East North Central	231	223	118	115	88	83
West North Central	149	158	63	70	32	49
South	463	447	251	234	167	130
South Atlantic	349	354	197	183	136	95
East South Central	188	164	86	87	54	56
West South Central	223	231	122	114	76	76
West	488	437	240	202	135	135
Mountain	228	195	112	93	69	53
Pacific	430	391	216	181	117	124
<b>COUNTY TYPE</b>						
Large Metro	740	702	322	290	206	174
Small Metro	508	494	224	209	129	132
250K – 1 Mil. Pop.	415	414	191	164	111	108
< 250K Pop.	300	300	123	130	72	76
Nonmetro	329	334	158	146	110	93
Urbanized	241	252	99	117	70	69
Less Urbanized	268	205	129	82	84	59
Completely Rural	75	110	27	35	14	25

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.78D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.31	0.29	0.16	0.14	0.11	0.09
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.61	0.69	0.32	0.30	0.25	0.19
New England	1.19	1.20	0.50	0.54	0.40	0.34
Middle Atlantic	0.71	0.82	0.39	0.35	0.31	0.24
Midwest	0.50	0.49	0.24	0.24	0.17	0.17
East North Central	0.60	0.58	0.31	0.30	0.23	0.21
West North Central	0.89	0.93	0.38	0.41	0.19	0.29
South	0.50	0.47	0.27	0.25	0.18	0.14
South Atlantic	0.71	0.71	0.40	0.36	0.28	0.19
East South Central	1.25	1.07	0.57	0.57	0.36	0.37
West South Central	0.77	0.78	0.42	0.39	0.26	0.25
West	0.83	0.73	0.41	0.34	0.23	0.22
Mountain	1.27	1.07	0.62	0.51	0.38	0.29
Pacific	1.06	0.94	0.53	0.43	0.29	0.30
<b>COUNTY TYPE</b>						
Large Metro	0.45	0.44	0.23	0.21	0.15	0.13
Small Metro	0.52	0.46	0.27	0.24	0.16	0.16
250K – 1 Mil. Pop.	0.62	0.57	0.34	0.28	0.21	0.20
< 250K Pop.	0.92	0.81	0.43	0.47	0.27	0.28
Nonmetro	0.69	0.66	0.36	0.32	0.26	0.22
Urbanized	1.06	1.08	0.52	0.58	0.39	0.36
Less Urbanized	1.06	0.80	0.58	0.40	0.40	0.30
Completely Rural	1.60	1.96	0.71	0.78	0.36	0.57

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.78P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.2064		0.0026		0.0078	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.8015		0.2385		0.0924	
New England	0.9600		0.4463		0.0495	
Middle Atlantic	0.7955		0.3467		0.3662	
Midwest	0.8791		0.4215		0.9937	
East North Central	0.5838		0.2914		0.6679	
West North Central	0.5716		0.8577		0.4150	
South	0.5786		0.2317		0.0351	
South Atlantic	0.4044		0.1466		0.0070	
East South Central	0.4152		0.7367		0.8728	
West South Central	0.4488		0.6762		0.8998	
West	0.1861		0.0094		0.2129	
Mountain	0.6174		0.1332		0.1144	
Pacific	0.2178		0.0345		0.5459	
<b>COUNTY TYPE</b>						
Large Metro	0.0949		0.0109		0.0161	
Small Metro	0.9804		0.1116		0.3322	
250K – 1 Mil. Pop.	0.4835		0.0470		0.2005	
< 250K Pop.	0.2796		0.9584		0.8069	
Nonmetro	0.8269		0.6481		0.3983	
Urbanized	0.8822		0.6356		0.6456	
Less Urbanized	0.5682		0.2039		0.3438	
Completely Rural	0.0624		0.4294		0.3492	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.79C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	68	64	58	53	38	35
<b>GEOGRAPHIC DIVISION</b>						
Northeast	20	24	18	19	11	14
New England	11	10	10	9	6	6
Middle Atlantic	17	22	14	17	10	12
Midwest	24	26	21	21	14	15
East North Central	20	20	17	17	11	12
West North Central	13	16	12	13	8	8
South	47	39	39	33	24	21
South Atlantic	36	26	30	22	18	15
East South Central	15	15	13	13	9	7
West South Central	25	24	21	20	14	13
West	41	37	35	32	25	19
Mountain	21	18	20	13	17	9
Pacific	34	33	29	29	17	17
<b>COUNTY TYPE</b>						
Large Metro	58	56	48	43	31	26
Small Metro	41	41	33	35	20	21
250K – 1 Mil. Pop.	34	31	28	27	17	16
< 250K Pop.	24	26	17	20	11	14
Nonmetro	34	27	29	22	16	13
Urbanized	27	20	22	17	14	9
Less Urbanized	21	21	17	15	8	9
Completely Rural	6	8	4	6	3	3

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.79D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.28	0.26	0.24	0.21	0.16	0.14
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.48	0.57	0.42	0.45	0.27	0.32
New England	0.98	0.94	0.94	0.77	0.54	0.57
Middle Atlantic	0.55	0.69	0.46	0.54	0.31	0.38
Midwest	0.45	0.47	0.39	0.38	0.26	0.27
East North Central	0.54	0.53	0.46	0.44	0.29	0.32
West North Central	0.84	0.99	0.75	0.78	0.52	0.51
South	0.52	0.42	0.44	0.35	0.27	0.23
South Atlantic	0.80	0.56	0.67	0.48	0.39	0.33
East South Central	1.06	1.05	0.93	0.90	0.62	0.49
West South Central	0.83	0.77	0.71	0.63	0.46	0.40
West	0.70	0.62	0.61	0.53	0.42	0.32
Mountain	1.18	0.97	1.09	0.72	0.97	0.47
Pacific	0.85	0.79	0.72	0.70	0.42	0.42
<b>COUNTY TYPE</b>						
Large Metro	0.40	0.37	0.34	0.29	0.23	0.19
Small Metro	0.47	0.47	0.39	0.40	0.26	0.26
250K – 1 Mil. Pop.	0.58	0.54	0.49	0.47	0.33	0.29
< 250K Pop.	0.80	0.86	0.62	0.71	0.44	0.53
Nonmetro	0.74	0.60	0.65	0.52	0.39	0.33
Urbanized	1.25	0.94	1.10	0.82	0.76	0.50
Less Urbanized	0.95	0.94	0.83	0.73	0.42	0.48
Completely Rural	1.76	1.99	1.22	1.48	0.95	0.96

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.79P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.1482		0.1571		0.3216	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.2603		0.5671		0.8407	
New England	0.9804		0.6410		0.7909	
Middle Atlantic	0.1884		0.3392		0.7128	
Midwest	0.2889		0.1884		0.8144	
East North Central	0.3061		0.3511		0.8619	
West North Central	0.6945		0.3506		0.8693	
South	0.2592		0.6648		0.8500	
South Atlantic	0.0192		0.1033		0.6239	
East South Central	0.6921		0.4503		0.4723	
West South Central	0.4689		0.4357		0.8750	
West	0.2244		0.1830		0.0862	
Mountain	0.0851		0.0302		0.0740	
Pacific	0.7094		0.8111		0.4313	
<b>COUNTY TYPE</b>						
Large Metro	0.4357		0.2992		0.1828	
Small Metro	0.0980		0.3499		0.9295	
250K – 1 Mil. Pop.	0.1035		0.3376		0.8223	
< 250K Pop.	0.5411		0.7782		0.8690	
Nonmetro	0.8624		0.8089		0.8629	
Urbanized	0.5592		0.2488		0.7304	
Less Urbanized	0.5539		0.7243		0.5517	
Completely Rural	0.4229		0.2185		0.8469	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.80C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	143	145	114	110	72	69
<b>GEOGRAPHIC DIVISION</b>						
Northeast	59	53	45	39	27	26
New England	27	25	20	20	14	13
Middle Atlantic	53	47	40	34	23	22
Midwest	62	58	45	41	28	27
East North Central	51	49	38	35	24	23
West North Central	33	31	25	24	16	14
South	93	93	71	68	46	44
South Atlantic	71	65	56	50	34	32
East South Central	34	36	27	25	18	21
West South Central	49	53	34	38	25	23
West	80	81	60	60	39	39
Mountain	34	41	25	29	16	20
Pacific	72	70	55	52	36	33
<b>COUNTY TYPE</b>						
Large Metro	153	141	99	93	56	53
Small Metro	127	129	85	77	46	49
250K – 1 Mil. Pop.	97	93	60	59	36	36
< 250K Pop.	80	91	57	50	31	32
Nonmetro	69	72	48	49	27	25
Urbanized	52	61	37	41	21	18
Less Urbanized	43	40	28	27	17	17
Completely Rural	20	21	13	8	2	4

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.80D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.42	0.42	0.33	0.32	0.21	0.20
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.98	0.86	0.74	0.64	0.44	0.42
New England	1.65	1.53	1.24	1.26	0.88	0.80
Middle Atlantic	1.19	1.03	0.91	0.75	0.52	0.49
Midwest	0.83	0.78	0.60	0.57	0.38	0.36
East North Central	1.00	0.97	0.74	0.69	0.47	0.46
West North Central	1.40	1.35	1.05	1.05	0.67	0.62
South	0.75	0.73	0.57	0.54	0.37	0.35
South Atlantic	1.13	1.01	0.89	0.78	0.54	0.50
East South Central	1.69	1.79	1.37	1.23	0.93	1.02
West South Central	1.18	1.27	0.83	0.92	0.61	0.56
West	0.99	0.99	0.74	0.73	0.49	0.47
Mountain	1.37	1.68	0.99	1.16	0.63	0.81
Pacific	1.29	1.22	0.97	0.89	0.64	0.58
<b>COUNTY TYPE</b>						
Large Metro	0.63	0.59	0.46	0.44	0.29	0.27
Small Metro	0.72	0.77	0.58	0.55	0.37	0.38
250K – 1 Mil. Pop.	0.85	0.98	0.68	0.73	0.47	0.47
< 250K Pop.	1.27	1.16	1.01	0.84	0.62	0.63
Nonmetro	0.98	1.02	0.83	0.84	0.51	0.46
Urbanized	1.47	1.67	1.33	1.43	0.85	0.71
Less Urbanized	1.26	1.29	0.99	1.08	0.67	0.71
Completely Rural	4.05	3.53	3.22	1.61	0.65	0.89

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.80P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.0066		0.0004		0.0024	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0025		0.0095		0.0671	
New England	0.0212		0.2194		0.1879	
Middle Atlantic	0.0269		0.0245		0.1933	
Midwest	0.6650		0.0875		0.0911	
East North Central	0.5580		0.1156		0.0955	
West North Central	0.9154		0.4782		0.6175	
South	0.0444		0.0134		0.0834	
South Atlantic	0.2482		0.0581		0.1635	
East South Central	0.9483		0.8460		0.8984	
West South Central	0.0414		0.0408		0.1548	
West	0.9661		0.4909		0.3121	
Mountain	0.1211		0.2417		0.3632	
Pacific	0.4083		0.1601		0.5061	
<b>COUNTY TYPE</b>						
Large Metro	0.0207		0.0045		0.1315	
Small Metro	0.1848		0.0300		0.0134	
250K – 1 Mil. Pop.	0.2809		0.0853		0.0022	
< 250K Pop.	0.4306		0.1830		0.9746	
Nonmetro	0.8443		0.6316		0.1518	
Urbanized	0.9540		0.7808		0.1756	
Less Urbanized	0.9579		0.9904		0.5380	
Completely Rural	0.4194		0.2775		0.6960	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.81C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	752	740	363	338	252	215
<b>GEOGRAPHIC DIVISION</b>						
Northeast	280	316	142	129	114	85
New England	139	143	57	61	46	38
Middle Atlantic	242	281	129	114	104	76
Midwest	263	261	122	124	87	89
East North Central	222	211	108	108	82	77
West North Central	142	154	56	65	28	47
South	441	437	225	214	158	119
South Atlantic	333	347	180	169	131	85
East South Central	180	156	77	78	48	49
West South Central	210	221	105	106	66	69
West	467	434	224	187	121	128
Mountain	219	188	100	84	62	48
Pacific	410	390	202	168	104	119
<b>COUNTY TYPE</b>						
Large Metro	688	671	294	266	194	160
Small Metro	462	457	199	181	118	118
250K – 1 Mil. Pop.	378	392	173	144	103	102
< 250K Pop.	270	264	101	109	62	60
Nonmetro	299	307	137	131	98	89
Urbanized	208	236	80	106	61	67
Less Urbanized	249	179	114	71	76	53
Completely Rural	66	96	22	30	13	24

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.81D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.39	0.37	0.19	0.17	0.13	0.11
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.77	0.87	0.39	0.35	0.31	0.23
New England	1.46	1.49	0.60	0.64	0.48	0.40
Middle Atlantic	0.91	1.04	0.48	0.43	0.39	0.28
Midwest	0.62	0.61	0.29	0.29	0.20	0.21
East North Central	0.75	0.71	0.36	0.36	0.28	0.26
West North Central	1.11	1.18	0.43	0.50	0.22	0.36
South	0.62	0.60	0.31	0.29	0.22	0.16
South Atlantic	0.87	0.89	0.47	0.43	0.34	0.22
East South Central	1.55	1.32	0.66	0.66	0.41	0.41
West South Central	0.96	0.99	0.48	0.47	0.30	0.31
West	1.04	0.95	0.50	0.41	0.27	0.28
Mountain	1.59	1.35	0.73	0.61	0.45	0.35
Pacific	1.32	1.23	0.65	0.53	0.33	0.37
<b>COUNTY TYPE</b>						
Large Metro	0.57	0.56	0.27	0.25	0.18	0.15
Small Metro	0.65	0.59	0.32	0.28	0.20	0.19
250K – 1 Mil. Pop.	0.77	0.72	0.42	0.33	0.26	0.24
< 250K Pop.	1.16	1.04	0.50	0.55	0.32	0.31
Nonmetro	0.83	0.80	0.41	0.38	0.30	0.27
Urbanized	1.26	1.39	0.59	0.73	0.46	0.47
Less Urbanized	1.32	0.95	0.68	0.44	0.47	0.35
Completely Rural	1.76	2.18	0.72	0.85	0.42	0.69

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.81P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.5191		0.0522		0.0776	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.8255		0.5637		0.1835	
New England	0.7100		0.7375		0.1021	
Middle Atlantic	0.9562		0.6307		0.4986	
Midwest	0.9232		0.9530		0.5554	
East North Central	0.7537		0.7038		0.9399	
West North Central	0.5094		0.4530		0.2505	
South	0.9678		0.6524		0.1064	
South Atlantic	0.6739		0.4648		0.0220	
East South Central	0.3855		0.8555		1.0000	
West South Central	0.2223		0.9092		0.6696	
West	0.2159		0.0158		0.4483	
Mountain	0.5501		0.0993		0.3085	
Pacific	0.2746		0.0638		0.7284	
<b>COUNTY TYPE</b>						
Large Metro	0.2194		0.0714		0.0499	
Small Metro	0.6213		0.4508		0.9049	
250K – 1 Mil. Pop.	0.2285		0.1856		0.7577	
< 250K Pop.	0.3660		0.5304		0.7648	
Nonmetro	0.7962		0.7773		0.6201	
Urbanized	0.8340		0.5555		0.9518	
Less Urbanized	0.5422		0.1971		0.3867	
Completely Rural	0.0328		0.3053		0.4077	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.82C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	777	752	389	357	265	228
<b>GEOGRAPHIC DIVISION</b>						
Northeast	283	322	147	136	117	89
New England	144	147	60	66	48	42
Middle Atlantic	243	286	134	120	105	80
Midwest	273	270	131	131	93	94
East North Central	230	221	116	113	87	82
West North Central	149	154	62	69	33	48
South	462	444	243	228	165	128
South Atlantic	349	351	192	179	135	93
East South Central	187	163	84	85	52	56
West South Central	221	230	117	111	73	73
West	481	439	237	198	132	133
Mountain	224	192	108	90	66	52
Pacific	425	393	214	178	115	123
<b>COUNTY TYPE</b>						
Large Metro	731	695	315	285	203	170
Small Metro	497	487	217	201	127	130
250K – 1 Mil. Pop.	407	411	185	158	109	107
< 250K Pop.	292	290	118	124	71	74
Nonmetro	324	325	153	138	107	91
Urbanized	232	245	92	110	66	68
Less Urbanized	266	197	126	78	83	58
Completely Rural	73	107	26	34	13	24

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.82D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.34	0.32	0.17	0.15	0.12	0.10
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.67	0.76	0.35	0.32	0.28	0.21
New England	1.29	1.31	0.54	0.58	0.43	0.37
Middle Atlantic	0.78	0.91	0.43	0.38	0.34	0.25
Midwest	0.55	0.54	0.26	0.26	0.19	0.19
East North Central	0.66	0.63	0.33	0.32	0.25	0.24
West North Central	0.98	1.00	0.40	0.45	0.22	0.31
South	0.55	0.52	0.29	0.27	0.20	0.15
South Atlantic	0.78	0.77	0.43	0.39	0.30	0.20
East South Central	1.37	1.18	0.62	0.62	0.38	0.41
West South Central	0.85	0.87	0.45	0.42	0.28	0.28
West	0.91	0.81	0.45	0.37	0.25	0.25
Mountain	1.38	1.18	0.66	0.55	0.41	0.32
Pacific	1.16	1.05	0.58	0.47	0.31	0.33
<b>COUNTY TYPE</b>						
Large Metro	0.50	0.48	0.25	0.23	0.16	0.14
Small Metro	0.57	0.51	0.29	0.26	0.18	0.18
250K – 1 Mil. Pop.	0.68	0.62	0.37	0.30	0.23	0.21
< 250K Pop.	1.00	0.89	0.47	0.50	0.30	0.30
Nonmetro	0.76	0.72	0.39	0.34	0.28	0.24
Urbanized	1.16	1.19	0.55	0.63	0.42	0.41
Less Urbanized	1.18	0.88	0.64	0.42	0.44	0.33
Completely Rural	1.72	2.10	0.76	0.84	0.39	0.61

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.82P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.2604		0.0039		0.0105	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.7387		0.2073		0.0846	
New England	0.9676		0.4891		0.0519	
Middle Atlantic	0.7207		0.2870		0.3389	
Midwest	0.9697		0.5448		0.9721	
East North Central	0.6572		0.3572		0.6442	
West North Central	0.5313		0.7292		0.4297	
South	0.6582		0.2436		0.0355	
South Atlantic	0.5299		0.2111		0.0084	
East South Central	0.3956		0.8186		0.9635	
West South Central	0.4852		0.5641		0.9166	
West	0.2236		0.0143		0.3183	
Mountain	0.7477		0.2277		0.2126	
Pacific	0.2288		0.0354		0.6139	
<b>COUNTY TYPE</b>						
Large Metro	0.1110		0.0140		0.0242	
Small Metro	0.8942		0.1356		0.3302	
250K – 1 Mil. Pop.	0.3865		0.0587		0.2054	
< 250K Pop.	0.2973		0.9196		0.8253	
Nonmetro	0.8295		0.6660		0.3785	
Urbanized	0.7863		0.5283		0.6891	
Less Urbanized	0.5082		0.1829		0.2994	
Completely Rural	0.0683		0.5454		0.3603	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.