

Table 1.83C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	610	598	336	302	215	189
GEOGRAPHIC DIVISION						
Northeast	225	236	120	123	85	78
New England	125	124	52	57	40	40
Middle Atlantic	187	200	107	109	75	67
Midwest	222	229	117	124	86	85
East North Central	184	194	103	106	81	76
West North Central	114	124	54	63	29	41
South	355	376	195	197	128	104
South Atlantic	265	291	148	153	103	75
East South Central	145	124	67	70	46	43
West South Central	173	211	103	104	61	63
West	372	339	217	155	125	106
Mountain	188	153	103	73	56	45
Pacific	319	305	193	135	112	97
COUNTY TYPE						
Large Metro	530	497	271	242	168	137
Small Metro	381	374	188	179	98	116
250K – 1 Mil. Pop.	294	310	160	144	86	98
< 250K Pop.	240	233	101	105	52	63
Nonmetro	226	257	128	108	101	75
Urbanized	161	194	83	81	61	55
Less Urbanized	176	157	99	65	79	46
Completely Rural	60	82	27	28	14	25

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.83D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.24	0.23	0.13	0.12	0.08	0.07
GEOGRAPHIC DIVISION						
Northeast	0.48	0.50	0.26	0.26	0.18	0.17
New England	1.02	1.01	0.43	0.46	0.33	0.32
Middle Atlantic	0.54	0.58	0.31	0.32	0.22	0.19
Midwest	0.40	0.41	0.21	0.22	0.16	0.15
East North Central	0.48	0.50	0.27	0.28	0.21	0.20
West North Central	0.68	0.73	0.32	0.37	0.17	0.24
South	0.38	0.39	0.21	0.21	0.14	0.11
South Atlantic	0.54	0.58	0.30	0.30	0.21	0.15
East South Central	0.96	0.81	0.45	0.46	0.31	0.28
West South Central	0.60	0.71	0.36	0.35	0.21	0.21
West	0.63	0.57	0.37	0.26	0.21	0.18
Mountain	1.04	0.84	0.57	0.40	0.31	0.25
Pacific	0.78	0.73	0.47	0.32	0.27	0.23
COUNTY TYPE						
Large Metro	0.34	0.33	0.20	0.17	0.12	0.10
Small Metro	0.42	0.40	0.23	0.20	0.13	0.14
250K – 1 Mil. Pop.	0.49	0.48	0.29	0.24	0.17	0.18
< 250K Pop.	0.80	0.72	0.37	0.38	0.20	0.23
Nonmetro	0.50	0.54	0.30	0.25	0.24	0.18
Urbanized	0.80	0.89	0.45	0.46	0.35	0.31
Less Urbanized	0.71	0.68	0.46	0.32	0.38	0.24
Completely Rural	1.42	1.69	0.69	0.64	0.36	0.57

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.83P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.1653		0.0051		0.0151	
GEOGRAPHIC DIVISION						
Northeast	0.3847		0.4226		0.2137	
New England	0.3646		0.3635		0.1782	
Middle Atlantic	0.6354		0.6513		0.5063	
Midwest	0.8169		0.4649		0.8382	
East North Central	0.8061		0.6477		0.9599	
West North Central	0.4238		0.5163		0.6057	
South	0.8104		0.4757		0.0825	
South Atlantic	0.6899		0.2823		0.0235	
East South Central	0.1853		0.8027		0.9439	
West South Central	0.3065		0.9803		0.9649	
West	0.0910		0.0025		0.1626	
Mountain	0.1203		0.0718		0.1537	
Pacific	0.2759		0.0140		0.3867	
COUNTY TYPE						
Large Metro	0.0807		0.0136		0.0296	
Small Metro	0.8040		0.4595		0.6606	
250K – 1 Mil. Pop.	0.7356		0.3533		0.3687	
< 250K Pop.	0.3677		0.9437		0.5101	
Nonmetro	0.8502		0.1442		0.1665	
Urbanized	0.7659		0.8464		0.4056	
Less Urbanized	0.7602		0.0479		0.1644	
Completely Rural	0.5042		0.9300		0.4485	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.84C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	64	59	54	48	35	31
GEOGRAPHIC DIVISION						
Northeast	19	23	17	18	11	12
New England	10	10	10	8	6	6
Middle Atlantic	16	21	13	16	9	11
Midwest	22	24	18	20	12	14
East North Central	18	19	15	16	10	11
West North Central	13	15	11	12	7	8
South	44	35	36	29	23	18
South Atlantic	34	23	27	20	17	13
East South Central	15	14	13	12	8	6
West South Central	24	22	20	17	13	10
West	37	36	32	30	21	17
Mountain	20	17	18	13	14	8
Pacific	31	31	25	27	16	15
COUNTY TYPE						
Large Metro	53	52	42	40	27	23
Small Metro	39	38	31	31	19	19
250K – 1 Mil. Pop.	32	30	26	26	17	14
< 250K Pop.	23	22	15	16	10	12
Nonmetro	33	22	27	18	16	12
Urbanized	26	17	22	12	14	9
Less Urbanized	20	18	16	13	8	8
Completely Rural	6	7	4	6	3	3

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.84D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.26	0.23	0.22	0.19	0.14	0.12
GEOGRAPHIC DIVISION						
Northeast	0.45	0.54	0.39	0.42	0.26	0.29
New England	0.94	0.88	0.91	0.74	0.53	0.55
Middle Atlantic	0.51	0.65	0.42	0.50	0.30	0.34
Midwest	0.42	0.44	0.35	0.36	0.23	0.25
East North Central	0.49	0.49	0.39	0.41	0.27	0.28
West North Central	0.81	0.93	0.70	0.73	0.43	0.51
South	0.49	0.37	0.40	0.31	0.25	0.20
South Atlantic	0.76	0.49	0.61	0.42	0.38	0.28
East South Central	1.02	0.93	0.87	0.82	0.58	0.41
West South Central	0.80	0.68	0.66	0.54	0.42	0.33
West	0.64	0.59	0.54	0.50	0.36	0.29
Mountain	1.14	0.94	1.03	0.72	0.78	0.44
Pacific	0.77	0.75	0.63	0.65	0.39	0.37
COUNTY TYPE						
Large Metro	0.37	0.34	0.31	0.27	0.20	0.17
Small Metro	0.45	0.43	0.37	0.36	0.25	0.24
250K – 1 Mil. Pop.	0.55	0.51	0.46	0.46	0.32	0.27
< 250K Pop.	0.77	0.77	0.58	0.57	0.40	0.45
Nonmetro	0.73	0.53	0.63	0.44	0.39	0.31
Urbanized	1.26	0.84	1.09	0.66	0.76	0.49
Less Urbanized	0.93	0.86	0.78	0.66	0.40	0.44
Completely Rural	1.75	1.68	1.19	1.38	0.91	0.84

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.84P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0770		0.2855		0.1993	
GEOGRAPHIC DIVISION						
Northeast	0.2632		0.6517		0.9289	
New England	0.9705		0.7031		0.7287	
Middle Atlantic	0.2000		0.4353		0.9290	
Midwest	0.2065		0.3438		0.6460	
East North Central	0.2837		0.6220		0.9926	
West North Central	0.5027		0.3637		0.4212	
South	0.0672		0.5968		0.4184	
South Atlantic	0.0091		0.1203		0.4365	
East South Central	0.9146		0.3332		0.5640	
West South Central	0.7538		0.6418		0.4393	
West	0.3757		0.3754		0.1107	
Mountain	0.1942		0.1161		0.0479	
Pacific	0.8262		0.9761		0.5624	
COUNTY TYPE						
Large Metro	0.5465		0.7475		0.1389	
Small Metro	0.0391		0.2872		0.6068	
250K – 1 Mil. Pop.	0.0802		0.3125		0.3699	
< 250K Pop.	0.2463		0.6592		0.6896	
Nonmetro	0.6918		0.5563		0.8599	
Urbanized	0.4030		0.1172		0.7398	
Less Urbanized	0.8364		0.6538		0.4639	
Completely Rural	0.7415		0.2801		0.8927	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.85C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	134	130	100	94	64	58
GEOGRAPHIC DIVISION						
Northeast	54	45	37	34	24	21
New England	26	25	17	17	11	11
Middle Atlantic	47	38	32	29	21	18
Midwest	57	53	42	36	28	23
East North Central	47	44	35	31	23	21
West North Central	30	31	23	20	16	11
South	83	87	60	58	36	34
South Atlantic	63	63	46	42	26	21
East South Central	31	35	24	24	15	19
West South Central	46	47	32	31	21	18
West	77	71	53	53	36	35
Mountain	31	32	22	28	14	19
Pacific	71	63	48	44	33	30
COUNTY TYPE						
Large Metro	134	119	84	80	50	46
Small Metro	107	110	70	62	38	38
250K – 1 Mil. Pop.	82	77	52	46	32	27
< 250K Pop.	70	79	47	41	23	26
Nonmetro	61	65	39	42	22	21
Urbanized	48	54	28	34	16	15
Less Urbanized	38	35	23	23	15	15
Completely Rural	18	19	13	7	2	4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.85D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.39	0.38	0.29	0.27	0.19	0.17
GEOGRAPHIC DIVISION						
Northeast	0.89	0.74	0.61	0.55	0.39	0.35
New England	1.60	1.54	1.07	1.06	0.69	0.71
Middle Atlantic	1.07	0.84	0.73	0.64	0.48	0.40
Midwest	0.77	0.72	0.56	0.49	0.37	0.32
East North Central	0.92	0.86	0.68	0.61	0.45	0.41
West North Central	1.26	1.38	0.96	0.87	0.66	0.47
South	0.67	0.69	0.49	0.46	0.29	0.27
South Atlantic	1.00	0.98	0.73	0.66	0.41	0.32
East South Central	1.58	1.75	1.20	1.21	0.75	0.97
West South Central	1.13	1.14	0.77	0.74	0.50	0.44
West	0.95	0.86	0.65	0.65	0.44	0.43
Mountain	1.24	1.31	0.88	1.12	0.55	0.76
Pacific	1.26	1.10	0.86	0.76	0.59	0.51
COUNTY TYPE						
Large Metro	0.58	0.53	0.41	0.39	0.27	0.24
Small Metro	0.67	0.67	0.51	0.46	0.31	0.31
250K – 1 Mil. Pop.	0.81	0.84	0.63	0.58	0.41	0.36
< 250K Pop.	1.22	1.07	0.85	0.74	0.48	0.55
Nonmetro	0.95	1.00	0.71	0.75	0.44	0.40
Urbanized	1.46	1.62	1.13	1.27	0.71	0.61
Less Urbanized	1.28	1.23	0.85	0.95	0.61	0.63
Completely Rural	3.89	3.47	3.20	1.33	0.62	0.88

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.85P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0007		0.0008		0.0004	
GEOGRAPHIC DIVISION						
Northeast	0.0007		0.0125		0.0518	
New England	0.2326		0.5594		0.3147	
Middle Atlantic	0.0014		0.0104		0.1009	
Midwest	0.8107		0.0134		0.0256	
East North Central	0.6524		0.0453		0.0851	
West North Central	0.8215		0.1433		0.1433	
South	0.0236		0.0396		0.0254	
South Atlantic	0.1518		0.1790		0.0235	
East South Central	0.9904		0.9099		0.3136	
West South Central	0.0307		0.0416		0.0319	
West	0.4449		0.7085		0.3134	
Mountain	0.3435		0.2702		0.3959	
Pacific	0.2044		0.2967		0.4886	
COUNTY TYPE						
Large Metro	0.0044		0.0241		0.0673	
Small Metro	0.0936		0.0136		0.0010	
250K – 1 Mil. Pop.	0.1524		0.0559		0.0002	
< 250K Pop.	0.3997		0.1169		0.7318	
Nonmetro	0.7113		0.5205		0.2981	
Urbanized	0.7617		0.5439		0.1283	
Less Urbanized	0.8212		0.7124		0.9178	
Completely Rural	0.2601		0.1375		0.7234	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.86C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	579	589	305	280	196	178
GEOGRAPHIC DIVISION						
Northeast	217	227	112	113	80	73
New England	118	116	50	50	38	37
Middle Atlantic	181	194	100	102	71	63
Midwest	209	216	106	114	78	80
East North Central	175	185	93	98	74	71
West North Central	106	117	50	58	25	39
South	339	369	173	179	121	95
South Atlantic	250	284	132	139	99	68
East South Central	138	118	59	63	42	34
West South Central	161	208	88	98	53	60
West	346	337	205	142	112	100
Mountain	176	149	94	65	52	41
Pacific	297	304	184	125	99	92
COUNTY TYPE						
Large Metro	491	473	249	219	156	125
Small Metro	338	342	162	157	86	107
250K – 1 Mil. Pop.	261	294	140	130	77	95
< 250K Pop.	211	199	84	88	46	51
Nonmetro	198	234	114	93	92	69
Urbanized	132	180	69	70	56	50
Less Urbanized	157	140	88	54	73	41
Completely Rural	51	69	22	26	13	24

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.86D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.30	0.30	0.16	0.14	0.10	0.09
GEOGRAPHIC DIVISION						
Northeast	0.60	0.62	0.31	0.31	0.22	0.20
New England	1.24	1.21	0.53	0.52	0.40	0.39
Middle Atlantic	0.68	0.72	0.38	0.38	0.26	0.23
Midwest	0.49	0.50	0.25	0.27	0.18	0.19
East North Central	0.59	0.62	0.31	0.33	0.25	0.24
West North Central	0.83	0.90	0.39	0.45	0.19	0.30
South	0.47	0.50	0.24	0.24	0.17	0.13
South Atlantic	0.65	0.73	0.35	0.36	0.26	0.17
East South Central	1.18	1.00	0.50	0.53	0.36	0.29
West South Central	0.74	0.93	0.41	0.44	0.24	0.27
West	0.77	0.74	0.46	0.31	0.25	0.22
Mountain	1.28	1.08	0.69	0.47	0.38	0.29
Pacific	0.95	0.96	0.59	0.39	0.32	0.29
COUNTY TYPE						
Large Metro	0.43	0.42	0.24	0.21	0.15	0.12
Small Metro	0.52	0.50	0.27	0.24	0.15	0.17
250K – 1 Mil. Pop.	0.60	0.62	0.34	0.29	0.19	0.22
< 250K Pop.	0.99	0.88	0.42	0.45	0.24	0.26
Nonmetro	0.59	0.65	0.35	0.28	0.28	0.21
Urbanized	0.94	1.12	0.51	0.51	0.42	0.38
Less Urbanized	0.86	0.81	0.54	0.34	0.45	0.26
Completely Rural	1.50	1.84	0.70	0.75	0.42	0.69

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.86P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.6238		0.0793		0.2108	
GEOGRAPHIC DIVISION						
Northeast	0.7880		0.8400		0.4391	
New England	0.4964		0.4862		0.2788	
Middle Atlantic	0.9356		0.9205		0.7785	
Midwest	0.5993		0.7427		0.6044	
East North Central	0.9921		0.7292		0.5112	
West North Central	0.3363		0.9266		0.8947	
South	0.6156		1.0000		0.3251	
South Atlantic	0.8293		0.6445		0.1027	
East South Central	0.1775		0.9731		0.6369	
West South Central	0.1147		0.5318		0.3796	
West	0.1369		0.0029		0.3564	
Mountain	0.1049		0.0424		0.4266	
Pacific	0.4027		0.0215		0.5206	
COUNTY TYPE						
Large Metro	0.2646		0.0525		0.1224	
Small Metro	0.6405		0.7136		0.4104	
250K – 1 Mil. Pop.	0.3185		0.9353		0.6502	
< 250K Pop.	0.5304		0.4196		0.4035	
Nonmetro	0.7381		0.2034		0.2179	
Urbanized	0.6907		0.8295		0.6503	
Less Urbanized	0.7797		0.0266		0.1155	
Completely Rural	0.3364		0.6780		0.4776	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.87C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	606	597	329	297	212	187
GEOGRAPHIC DIVISION						
Northeast	223	236	118	120	84	76
New England	123	124	51	55	40	40
Middle Atlantic	186	201	106	107	74	66
Midwest	220	227	116	122	86	85
East North Central	183	193	102	105	80	75
West North Central	114	122	54	62	30	41
South	355	377	189	192	127	103
South Atlantic	265	291	142	150	102	74
East South Central	145	124	65	69	45	42
West South Central	172	211	99	102	59	62
West	367	337	215	151	122	105
Mountain	182	149	99	69	54	44
Pacific	317	304	192	133	110	95
COUNTY TYPE						
Large Metro	523	491	264	236	166	133
Small Metro	370	368	181	172	95	114
250K – 1 Mil. Pop.	285	308	154	139	83	97
< 250K Pop.	233	225	99	100	52	61
Nonmetro	221	253	125	105	99	73
Urbanized	153	191	77	79	59	54
Less Urbanized	174	151	97	62	79	45
Completely Rural	58	80	26	27	13	24

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.87D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.26	0.26	0.14	0.13	0.09	0.08
GEOGRAPHIC DIVISION						
Northeast	0.53	0.55	0.28	0.28	0.20	0.18
New England	1.11	1.10	0.46	0.49	0.36	0.35
Middle Atlantic	0.60	0.64	0.34	0.34	0.24	0.21
Midwest	0.44	0.45	0.23	0.24	0.17	0.17
East North Central	0.52	0.55	0.29	0.30	0.23	0.22
West North Central	0.75	0.80	0.36	0.41	0.20	0.27
South	0.42	0.44	0.22	0.22	0.15	0.12
South Atlantic	0.59	0.64	0.32	0.33	0.23	0.16
East South Central	1.06	0.90	0.48	0.50	0.33	0.31
West South Central	0.67	0.80	0.38	0.38	0.23	0.23
West	0.69	0.63	0.41	0.28	0.23	0.19
Mountain	1.12	0.91	0.61	0.42	0.33	0.27
Pacific	0.86	0.81	0.52	0.36	0.30	0.25
COUNTY TYPE						
Large Metro	0.38	0.36	0.21	0.19	0.13	0.11
Small Metro	0.46	0.43	0.24	0.22	0.14	0.15
250K – 1 Mil. Pop.	0.54	0.54	0.31	0.26	0.18	0.19
< 250K Pop.	0.87	0.78	0.40	0.40	0.22	0.25
Nonmetro	0.56	0.60	0.32	0.27	0.26	0.19
Urbanized	0.89	0.99	0.49	0.50	0.38	0.34
Less Urbanized	0.79	0.74	0.50	0.34	0.42	0.26
Completely Rural	1.52	1.82	0.74	0.68	0.38	0.61

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.87P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.2286		0.0071		0.0236	
GEOGRAPHIC DIVISION						
Northeast	0.3323		0.3801		0.2119	
New England	0.3638		0.3963		0.1885	
Middle Atlantic	0.5540		0.5747		0.4932	
Midwest	0.7076		0.5536		0.7738	
East North Central	0.8977		0.6976		0.9600	
West North Central	0.3721		0.6187		0.4866	
South	0.9689		0.5179		0.1058	
South Atlantic	0.9059		0.3970		0.0297	
East South Central	0.1934		0.9423		0.9772	
West South Central	0.3232		0.9488		0.9210	
West	0.1076		0.0033		0.2450	
Mountain	0.1492		0.1083		0.2928	
Pacific	0.2848		0.0136		0.4278	
COUNTY TYPE						
Large Metro	0.0937		0.0132		0.0477	
Small Metro	0.9741		0.5629		0.7226	
250K – 1 Mil. Pop.	0.5971		0.4354		0.4468	
< 250K Pop.	0.4202		0.8864		0.5564	
Nonmetro	0.8087		0.1628		0.1486	
Urbanized	0.6631		0.9288		0.4234	
Less Urbanized	0.7301		0.0361		0.1302	
Completely Rural	0.5151		0.9264		0.4305	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.