

Table 2.11A Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	174,409	173,890	83,384	81,859	69,663	68,225
12	206	183	93	107	41	38
13	479	419	309	259	147	115
14	718	697	530	472	258	229
15	1,246	1,120	882	807	512	443
16	1,536	1,550	1,158	1,163	668	699
17	1,927	1,814	1,445	1,395	985	966
18	2,707	2,705	2,249	2,245	1,680	1,677
19	2,851	2,622	2,315	2,134	1,775	1,595
20	3,150	2,979	2,380	2,248	1,920	1,782
21	2,923	3,083	2,202	2,199	1,777	1,763
22	2,948	2,999	2,218	2,188	1,771	1,783
23	2,941	2,918	2,040	2,070	1,656	1,684
24	3,044	3,061	2,085	2,107	1,711	1,668
25	2,874	2,936	1,993	1,969	1,643	1,581
26-29	12,811	12,433	8,225	8,111	7,003	6,750
30-34	15,074	14,351	8,412	8,048	7,076	6,759
35-39	13,696	14,135	7,058	7,156	5,943	6,109
40-44	15,386	15,075	7,559	6,990	6,528	5,985
45-49	17,339	16,162	7,922	7,577	6,995	6,651
50-54	16,828	17,575	7,349	7,770	6,315	7,036
55-59	14,648	14,132	5,832	5,063	5,143	4,379
60-64	12,018	13,237	3,759	3,965	3,375	3,486
65 or Older	27,060	27,704	5,365	5,816	4,739	5,048

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.11B Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	68.8 ^b	67.5	32.9 ^a	31.8	27.5 ^a	26.5
12	5.4	4.9	2.4	2.9	1.1	1.0
13	11.7	10.1	7.6	6.2	3.6	2.8
14	18.5	16.5	13.7 ^b	11.1	6.7	5.4
15	30.0 ^b	26.4	21.3 ^a	19.0	12.3 ^a	10.4
16	36.9	35.5	27.8	26.6	16.1	16.0
17	45.4	43.0	34.0	33.0	23.2	22.9
18	57.3	57.1	47.6	47.4	35.6	35.4
19	65.4 ^b	61.0	53.1 ^a	49.6	40.7 ^a	37.1
20	69.3	66.4	52.4	50.1	42.2	39.7
21	68.8	70.6	51.9	50.4	41.9	40.4
22	71.4	69.7	53.7	50.8	42.9	41.4
23	72.8	73.2	50.5	51.9	41.0	42.2
24	74.6	73.4	51.1	50.5	41.9	40.0
25	72.8	74.3	50.5	49.8	41.6	40.0
26-29	76.9 ^a	73.1	49.4	47.7	42.0	39.7
30-34	75.9	74.0	42.4	41.5	35.6	34.9
35-39	70.9	71.7	36.5	36.3	30.8	31.0
40-44	75.1	74.1	36.9	34.4	31.8	29.4
45-49	78.0 ^a	75.0	35.6	35.1	31.5	30.8
50-54	76.5	76.2	33.4	33.7	28.7	30.5
55-59	76.0	74.7	30.3	26.7	26.7	23.1
60-64	73.1	73.2	22.9	21.9	20.5	19.3
65 or Older	69.5	68.9	13.8	14.5	12.2	12.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.12A Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	162,940	161,799	68,599	67,126	58,336	56,819
12	151	136	65	76	25	27
13	373	308	245 ^a	181	117	80
14	580	559	402	365	206	182
15	1,008	909	689	601	389	323
16	1,238	1,317	889	940	517	562
17	1,637	1,544	1,169	1,123	786	763
18	2,336	2,292	1,830	1,798	1,337	1,370
19	2,518	2,302	1,941	1,821	1,490	1,353
20	2,792	2,675	1,996	1,932	1,553	1,549
21	2,657	2,718	1,855	1,839	1,517	1,497
22	2,731	2,767	1,889	1,920	1,521	1,496
23	2,687	2,678	1,737	1,787	1,392	1,473
24	2,820	2,770	1,829	1,770	1,500	1,431
25	2,682	2,728	1,657	1,648	1,391	1,318
26-29	12,076	11,565	6,982	6,847	6,141	5,741
30-34	14,214	13,596	7,100	6,822	6,142	5,754
35-39	12,912	13,068	5,665	5,750	4,767	4,934
40-44	14,283	14,129	6,136	5,502	5,460	4,911
45-49	16,275	15,124	6,412	6,277	5,845	5,699
50-54	16,019	16,695	6,118	6,407	5,529	6,086
55-59	13,922	13,283	4,879	4,049	4,375 ^a	3,545
60-64	11,389	12,480	3,077	3,215	2,775	2,838
65 or Older	25,639	26,157	4,038	4,455	3,561	3,884

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.12B Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	64.2 ^b	62.8	27.0 ^a	26.1	23.0 ^a	22.1
12	3.9	3.6	1.7	2.0	0.7	0.7
13	9.1 ^a	7.4	6.0 ^a	4.4	2.9	1.9
14	14.9	13.2	10.3 ^a	8.6	5.3	4.3
15	24.3 ^a	21.4	16.6 ^a	14.1	9.4 ^a	7.6
16	29.7	30.2	21.4	21.5	12.4	12.9
17	38.5	36.6	27.5	26.6	18.5	18.1
18	49.4	48.4	38.7	37.9	28.3	28.9
19	57.7 ^a	53.5	44.5	42.4	34.2	31.5
20	61.4	59.6	43.9	43.1	34.2	34.5
21	62.6	62.2	43.7	42.1	35.7	34.3
22	66.1	64.3	45.7	44.6	36.8	34.8
23	66.6	67.2	43.0	44.8	34.5	37.0
24	69.1	66.4	44.8	42.4	36.8	34.3
25	67.9	69.0	42.0	41.7	35.2	33.4
26-29	72.5 ^b	68.0	41.9	40.2	36.9	33.7
30-34	71.6	70.1	35.7	35.2	30.9	29.7
35-39	66.9	66.3	29.3	29.2	24.7	25.0
40-44	69.7	69.4	29.9	27.0	26.6	24.1
45-49	73.2 ^a	70.1	28.8	29.1	26.3	26.4
50-54	72.9	72.4	27.8	27.8	25.1	26.4
55-59	72.3	70.2	25.3 ^a	21.4	22.7 ^a	18.7
60-64	69.2	69.0	18.7	17.8	16.9	15.7
65 or Older	65.8	65.1	10.4	11.1	9.1	9.7

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.13A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	45,036	46,269	12,383	11,746	8,879	8,243
12	53	52	26	31	9	9
13	124	103	82	58	39	27
14	194	182	150 ^a	104	56	38
15	348	319	251	233	110	106
16	469	465	319	331	153	155
17	594	567	386	368	202	181
18	853	920	624	573	369	319
19	886 ^a	739	533 ^a	407	299 ^b	189
20	971	891	492	450	330 ^a	240
21	901	968	426	403	228	231
22	911	979	420	422	266	301
23	902 ^a	771	369	317	235	184
24	851	847	359	324	228	191
25	887	887	347	322	227	212
26-29	3,762	3,524	1,182	1,119	862	688
30-34	4,628	4,793	1,231	1,145	898	882
35-39	5,106	4,510	1,320	1,118	1,165	967
40-44	5,154	5,181	917	1,017	781	892
45-49	4,656	4,831	906	756	729	671
50-54	3,661	4,013	491	617	392	401
55-59	2,614	3,013	501	473	457	373
60-64	2,013	2,390	345	418	238	345
65 or Older	4,498	5,325	706	739	604	641

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.13B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	17.8	18.0	4.9	4.6	3.5	3.2
12	1.4	1.4	0.7	0.8	0.2	0.2
13	3.0	2.5	2.0	1.4	1.0	0.7
14	5.0	4.3	3.9 ^b	2.5	1.5	0.9
15	8.4	7.5	6.0	5.5	2.6	2.5
16	11.3	10.6	7.7	7.6	3.7	3.6
17	14.0	13.4	9.1	8.7	4.8	4.3
18	18.0	19.4	13.2	12.1	7.8	6.7
19	20.3 ^a	17.2	12.2 ^a	9.5	6.9 ^b	4.4
20	21.4	19.9	10.8	10.0	7.3 ^a	5.4
21	21.2	22.2	10.0	9.2	5.4	5.3
22	22.1	22.8	10.2	9.8	6.5	7.0
23	22.3 ^a	19.3	9.1	7.9	5.8	4.6
24	20.9	20.3	8.8	7.8	5.6	4.6
25	22.5	22.5	8.8	8.2	5.7	5.4
26-29	22.6	20.7	7.1	6.6	5.2	4.0
30-34	23.3	24.7	6.2	5.9	4.5	4.5
35-39	26.4 ^b	22.9	6.8	5.7	6.0	4.9
40-44	25.1	25.5	4.5	5.0	3.8	4.4
45-49	20.9	22.4	4.1	3.5	3.3	3.1
50-54	16.7	17.4	2.2	2.7	1.8	1.7
55-59	13.6	15.9	2.6	2.5	2.4	2.0
60-64	12.2	13.2	2.1	2.3	1.4	1.9
65 or Older	11.6	13.2	1.8	1.8	1.6	1.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.14A Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	88,760	88,505	26,061	24,980	13,273	12,865
12	34 ^a	64	17 ^a	42	9	14
13	151	158	93	101	33	35
14	275	261	197	177	60	61
15	545	511	376	360	159	153
16	742	739	532	558	206	227
17	947	942	667	701	321	368
18	1,601	1,494	1,274	1,153	647	589
19	1,682 ^a	1,460	1,134	1,053	567	539
20	1,790	1,748	1,065	1,107	557	554
21	1,813	1,837	1,011	976	488	494
22	1,780	1,743	927	889	433	437
23	1,820	1,676	811	778	403	406
24	1,763	1,856	755	866	381	390
25	1,738	1,725	759	746	382	314
26-29	7,809	7,224	2,847	2,917	1,282	1,573
30-34	8,623	8,550	2,399	2,290	1,208	1,179
35-39	7,476	7,641	2,115 ^a	1,696	1,072	892
40-44	8,379	7,623	2,139	1,826	1,091	874
45-49	8,283	7,698	1,979	1,750	1,164 ^b	772
50-54	7,869	8,351	1,712	1,707	894	956
55-59	7,025	6,962	1,447	1,120	759	659
60-64	5,533	6,041	815	920	528	548
65 or Older	11,083	12,200	991	1,247	631	830

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.14B Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	35.0	34.4	10.3 ^a	9.7	5.2	5.0
12	0.9 ^a	1.7	0.4 ^a	1.1	0.2	0.4
13	3.7	3.8	2.3	2.4	0.8	0.8
14	7.1	6.2	5.1	4.2	1.6	1.4
15	13.1	12.0	9.1	8.5	3.8	3.6
16	17.8	16.9	12.8	12.8	4.9	5.2
17	22.3	22.3	15.7	16.6	7.6	8.7
18	33.9	31.5	27.0	24.3	13.7	12.4
19	38.6 ^b	34.0	26.0	24.5	13.0	12.5
20	39.4	39.0	23.4	24.7	12.3	12.4
21	42.7	42.1	23.8	22.4	11.5	11.3
22	43.1	40.5	22.4	20.7	10.5	10.2
23	45.1	42.1	20.1	19.5	10.0	10.2
24	43.2	44.5	18.5	20.8	9.3	9.4
25	44.0	43.7	19.2	18.9	9.7	7.9
26-29	46.9 ^a	42.4	17.1	17.1	7.7	9.2
30-34	43.4	44.1	12.1	11.8	6.1	6.1
35-39	38.7	38.8	10.9 ^a	8.6	5.5	4.5
40-44	40.9 ^a	37.5	10.4	9.0	5.3	4.3
45-49	37.3	35.7	8.9	8.1	5.2 ^a	3.6
50-54	35.8	36.2	7.8	7.4	4.1	4.1
55-59	36.5	36.8	7.5	5.9	3.9	3.5
60-64	33.6	33.4	5.0	5.1	3.2	3.0
65 or Older	28.5	30.3	2.5	3.1	1.6	2.1

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.15A Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	209,264 ^a	211,747	168,381	170,422	131,374	133,385
12	272	265	170	158	61	60
13	713	657	515	450	189 ^a	136
14	1,111	1,082	835	836	353	327
15	1,792	1,760	1,477	1,391	645	632
16	2,181	2,291	1,820	1,900	877	937
17	2,552	2,555	2,165	2,205	1,193	1,234
18	3,410	3,187	3,014	2,849	1,963	1,858
19	3,391	3,224	3,038	2,850	2,134	1,988
20	3,773	3,648	3,400	3,299	2,519	2,482
21	3,840	3,850	3,663	3,664	3,033	3,045
22	3,719	3,900	3,481	3,615	2,845	3,000
23	3,710	3,626	3,465	3,322	2,865	2,794
24	3,728	3,817	3,399	3,498	2,800	2,906
25	3,617	3,649	3,323	3,333	2,760	2,741
26-29	15,173	15,389	13,434	13,672	10,875	11,115
30-34	17,925	17,390	15,526	15,159	12,688	12,115
35-39	17,156	17,688	14,660	15,157	11,709	11,978
40-44	18,511	18,626	15,777	15,578	12,515	12,777
45-49	20,441	19,248	16,364	15,410	13,093 ^a	12,048
50-54	19,784	20,758	15,733	16,677	12,708	13,792
55-59	17,456	16,890	12,833	12,802	10,171	9,991
60-64	14,490	15,807	10,586	11,872	8,484	9,216
65 or Older	30,517 ^b	32,439	19,702	20,726	14,896	16,213

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.15B Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	82.5	82.2	66.4	66.2	51.8	51.8
12	7.1	7.1	4.4	4.2	1.6	1.6
13	17.5	15.8	12.6	10.8	4.6 ^a	3.3
14	28.6 ^a	25.6	21.5	19.7	9.1	7.7
15	43.2	41.4	35.6	32.8	15.5	14.9
16	52.4	52.5	43.7	43.5	21.1	21.5
17	60.1	60.5	51.0	52.2	28.1	29.2
18	72.2 ^b	67.2	63.8 ^a	60.1	41.5	39.2
19	77.8	75.0	69.7 ^a	66.3	48.9	46.2
20	83.0	81.3	74.8	73.6	55.4	55.3
21	90.4	88.2	86.2	83.9	71.4	69.7
22	90.1	90.6	84.3	84.0	68.9	69.7
23	91.9	91.0	85.8	83.3	70.9	70.1
24	91.4	91.5	83.3	83.9	68.6	69.7
25	91.6	92.4	84.2	84.4	69.9	69.4
26-29	91.1	90.4	80.6	80.3	65.3	65.3
30-34	90.2	89.7	78.2	78.2	63.9	62.5
35-39	88.8	89.7	75.9	76.9	60.6	60.8
40-44	90.3	91.6	77.0	76.6	61.1	62.8
45-49	92.0 ^b	89.3	73.6	71.5	58.9	55.9
50-54	90.0	90.0	71.6	72.3	57.8	59.8
55-59	90.6	89.2	66.6	67.6	52.8	52.8
60-64	88.1	87.4	64.4	65.6	51.6	50.9
65 or Older	78.4	80.7	50.6	51.5	38.3	40.3

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.16A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011

Age Category	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	131,374	133,385	58,535	58,341	16,899	15,865
12	61	60	14	29	2	2
13	189 ^a	136	67	54	8	2
14	353	327	162	144	25	16
15	645	632	380	338	71	64
16	877	937	517	525	114	120
17	1,193	1,234	774	760	188	174
18	1,963	1,858	1,346	1,294	387	366
19	2,134	1,988	1,483 ^a	1,290	526	426
20	2,519	2,482	1,682	1,642	619	527
21	3,033	3,045	2,023	1,989	739	675
22	2,845	3,000	1,879	1,981	670	629
23	2,865	2,794	1,841	1,860	586	573
24	2,800	2,906	1,816	1,852	535	542
25	2,760	2,741	1,726	1,744	554 ^a	427
26-29	10,875	11,115	6,417	6,509	1,896	2,014
30-34	12,688	12,115	6,880	6,495	1,843	1,810
35-39	11,709	11,978	5,533	5,299	1,508	1,226
40-44	12,515	12,777	5,441	5,366	1,531	1,255
45-49	13,093 ^a	12,048	5,555	5,272	1,610	1,369
50-54	12,708	13,792	4,536	4,519	1,314	1,326
55-59	10,171	9,991	3,258	3,407	999	854
60-64	8,484	9,216	2,265	2,619	535	794
65 or Older	14,896	16,213	2,943	3,352	637	673

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.16B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2010 and 2011

Age Category	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	51.8	51.8	23.1	22.6	6.7 ^a	6.2
12	1.6	1.6	0.4	0.8	0.0	0.1
13	4.6 ^a	3.3	1.6	1.3	0.2	0.1
14	9.1	7.7	4.2	3.4	0.7	0.4
15	15.5	14.9	9.1	7.9	1.7	1.5
16	21.1	21.5	12.4	12.0	2.8	2.7
17	28.1	29.2	18.2	18.0	4.4	4.1
18	41.5	39.2	28.5	27.3	8.2	7.7
19	48.9	46.2	34.0 ^a	30.0	12.1	9.9
20	55.4	55.3	37.0	36.6	13.6	11.8
21	71.4	69.7	47.6	45.6	17.4	15.5
22	68.9	69.7	45.5	46.0	16.2	14.6
23	70.9	70.1	45.6	46.7	14.5	14.4
24	68.6	69.7	44.5	44.4	13.1	13.0
25	69.9	69.4	43.7	44.2	14.0 ^a	10.8
26-29	65.3	65.3	38.5	38.2	11.4	11.8
30-34	63.9	62.5	34.6	33.5	9.3	9.3
35-39	60.6	60.8	28.6	26.9	7.8	6.2
40-44	61.1	62.8	26.5	26.4	7.5	6.2
45-49	58.9	55.9	25.0	24.5	7.2	6.4
50-54	57.8	59.8	20.6	19.6	6.0	5.7
55-59	52.8	52.8	16.9	18.0	5.2	4.5
60-64	51.6	50.9	13.8	14.5	3.3	4.4
65 or Older	38.3	40.3	7.6	8.3	1.6	1.7

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.