

Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	174,409	173,890	83,384	81,859	69,663	68,225
Cigarettes	162,940	161,799	68,599	67,126	58,336	56,819
Smokeless Tobacco	45,036	46,269	12,383	11,746	8,879	8,243
Cigars	88,760	88,505	26,061	24,980	13,273	12,865
Pipe Tobacco ²	34,380	34,532	--	--	2,155	2,137
ALCOHOL	209,264 ^a	211,747	168,381	170,422	131,374	133,385
Binge Alcohol Use ³	--	--	--	--	58,535	58,341
Heavy Alcohol Use ³	--	--	--	--	16,899	15,865

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	68.8 ^b	67.5	32.9 ^a	31.8	27.5 ^a	26.5
Cigarettes	64.2 ^b	62.8	27.0 ^a	26.1	23.0 ^a	22.1
Smokeless Tobacco	17.8	18.0	4.9	4.6	3.5	3.2
Cigars	35.0	34.4	10.3 ^a	9.7	5.2	5.0
Pipe Tobacco ²	13.6	13.4	--	--	0.8	0.8
ALCOHOL	82.5	82.2	66.4	66.2	51.8	51.8
Binge Alcohol Use ³	--	--	--	--	23.1	22.6
Heavy Alcohol Use ³	--	--	--	--	6.7 ^a	6.2

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	685	602	403	366	188	154
Cigarettes	524	444	311	257	142	108
Smokeless Tobacco	178	155	109	89	48	36
Cigars	185	222	109	142	42	49
Pipe Tobacco ²	47	45	--	--	17	16
ALCOHOL	986	922	685	608	250	196
Binge Alcohol Use ³	--	--	--	--	80	83
Heavy Alcohol Use ³	--	--	--	--	10	4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	8.7	7.6	5.1	4.6	2.4	1.9
Cigarettes	6.6	5.6	3.9	3.3	1.8	1.4
Smokeless Tobacco	2.2	2.0	1.4	1.1	0.6	0.5
Cigars	2.3	2.8	1.4	1.8	0.5	0.6
Pipe Tobacco ²	0.6	0.6	--	--	0.2	0.2
ALCOHOL	12.5	11.7	8.7	7.7	3.2	2.5
Binge Alcohol Use ³	--	--	--	--	1.0	1.1
Heavy Alcohol Use ³	--	--	--	--	0.1	0.1

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	1,964	1,816	1,413	1,279	771	671
Cigarettes	1,588	1,468	1,091 ^a	966	595	506
Smokeless Tobacco	542	501	401	337	166	144
Cigars	820	772	573	538	219	214
Pipe Tobacco ²	149	176	--	--	47	53
ALCOHOL	2,903	2,842	2,313	2,227	998	959
Binge Alcohol Use ³	--	--	--	--	542	481
Heavy Alcohol Use ³	--	--	--	--	97	81

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	24.4 ^b	21.4	17.6 ^b	15.1	9.6 ^b	7.9
Cigarettes	19.8 ^b	17.3	13.6 ^b	11.4	7.4 ^a	6.0
Smokeless Tobacco	6.7	5.9	5.0 ^a	4.0	2.1	1.7
Cigars	10.2	9.1	7.1	6.3	2.7	2.5
Pipe Tobacco ²	1.9	2.1	--	--	0.6	0.6
ALCOHOL	36.1 ^a	33.5	28.8 ^b	26.3	12.4	11.3
Binge Alcohol Use ³	--	--	--	--	6.7 ^a	5.7
Heavy Alcohol Use ³	--	--	--	--	1.2	0.9

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	3,463	3,364	2,603	2,558	1,654	1,665
Cigarettes	2,875	2,861	2,058	2,064	1,304	1,325
Smokeless Tobacco	1,063	1,031	705	699	355	337
Cigars	1,688	1,681	1,199	1,259	527	595
Pipe Tobacco ²	344	353	--	--	91	112
ALCOHOL	4,733	4,847	3,985	4,105	2,071	2,171
Binge Alcohol Use ³	--	--	--	--	1,290	1,285
Heavy Alcohol Use ³	--	--	--	--	303	294

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	41.2	39.2	31.0	29.8	19.7	19.4
Cigarettes	34.2	33.3	24.5	24.0	15.5	15.4
Smokeless Tobacco	12.6	12.0	8.4	8.1	4.2	3.9
Cigars	20.1	19.6	14.3	14.7	6.3	6.9
Pipe Tobacco ²	4.1	4.1	--	--	1.1	1.3
ALCOHOL	56.3	56.4	47.4	47.8	24.6	25.3
Binge Alcohol Use ³	--	--	--	--	15.3	15.0
Heavy Alcohol Use ³	--	--	--	--	3.6	3.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	8,708	8,305	6,945	6,626	5,376	5,054
Cigarettes	7,646	7,268	5,766	5,552	4,381	4,272
Smokeless Tobacco	2,709	2,550	1,649 ^a	1,430	998 ^b	748
Cigars	5,073 ^a	4,702	3,473	3,313	1,771	1,683
Pipe Tobacco ²	1,209	1,189	--	--	339	291
ALCOHOL	10,574	10,059	9,452	8,998	6,615	6,328
Binge Alcohol Use ³	--	--	--	--	4,511	4,226
Heavy Alcohol Use ³	--	--	--	--	1,532	1,319

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	63.9 ^a	61.4	50.9	49.0	39.4 ^a	37.4
Cigarettes	56.1 ^a	53.7	42.3	41.0	32.1	31.6
Smokeless Tobacco	19.9	18.9	12.1 ^a	10.6	7.3 ^b	5.5
Cigars	37.2 ^a	34.8	25.5	24.5	13.0	12.4
Pipe Tobacco ²	8.9	8.8	--	--	2.5	2.2
ALCOHOL	77.6 ^b	74.4	69.3 ^b	66.5	48.5	46.8
Binge Alcohol Use ³	--	--	--	--	33.1	31.2
Heavy Alcohol Use ³	--	--	--	--	11.2 ^a	9.8

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	14,729	14,998	10,539	10,533	8,558	8,479
Cigarettes	13,578	13,661	8,967	8,964	7,321	7,216
Smokeless Tobacco	4,451	4,452	1,920	1,789	1,184	1,119
Cigars	8,914	8,838	4,262	4,256	2,087	2,041
Pipe Tobacco ²	2,068	2,141	--	--	280	344
ALCOHOL	18,615	18,842	17,331	17,432	14,302	14,486
Binge Alcohol Use ³	--	--	--	--	9,285	9,427
Heavy Alcohol Use ³	--	--	--	--	3,085	2,846

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	72.1	72.2	51.6	50.7	41.9	40.8
Cigarettes	66.4	65.7	43.9	43.1	35.8	34.7
Smokeless Tobacco	21.8	21.4	9.4	8.6	5.8	5.4
Cigars	43.6	42.5	20.9	20.5	10.2	9.8
Pipe Tobacco ²	10.1	10.3	--	--	1.4	1.7
ALCOHOL	91.1	90.7	84.8	83.9	70.0	69.7
Binge Alcohol Use ³	--	--	--	--	45.4	45.4
Heavy Alcohol Use ³	--	--	--	--	15.1 ^a	13.7

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	27,885 ^b	26,784	16,637	16,159	14,079	13,509
Cigarettes	26,289 ^b	25,161	14,082	13,669	12,283	11,495
Smokeless Tobacco	8,390	8,316	2,413	2,264	1,760	1,570
Cigars	16,432	15,773	5,245	5,207	2,489	2,753
Pipe Tobacco ²	3,240	2,816	--	--	290	284
ALCOHOL	33,098	32,779	28,960	28,831	23,563	23,230
Binge Alcohol Use ³	--	--	--	--	13,297	13,004
Heavy Alcohol Use ³	--	--	--	--	3,739	3,823

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	76.3 ^b	73.6	45.6	44.4	38.5	37.1
Cigarettes	72.0 ^b	69.1	38.6	37.5	33.6	31.6
Smokeless Tobacco	23.0	22.8	6.6	6.2	4.8	4.3
Cigars	45.0	43.3	14.4	14.3	6.8	7.6
Pipe Tobacco ²	8.9	7.7	--	--	0.8	0.8
ALCOHOL	90.6	90.0	79.3	79.2	64.5	63.8
Binge Alcohol Use ³	--	--	--	--	36.4	35.7
Heavy Alcohol Use ³	--	--	--	--	10.2	10.5

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	116,975	118,020	44,844	44,337	39,038	38,694
Cigarettes	110,439	110,935	36,324	35,655	32,311	31,897
Smokeless Tobacco	27,702	29,263	5,187	5,138	4,367	4,289
Cigars	55,647	56,517	11,198	10,265	6,138	5,530
Pipe Tobacco ²	27,323	27,811	--	--	1,091	1,036
ALCOHOL	138,355 ^b	141,457	105,656 ^a	108,222	83,575 ^a	86,014
Binge Alcohol Use ³	--	--	--	--	29,530	29,834
Heavy Alcohol Use ³	--	--	--	--	8,135	7,497

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	73.7	72.9	28.3	27.4	24.6	23.9
Cigarettes	69.6	68.5	22.9	22.0	20.4	19.7
Smokeless Tobacco	17.5	18.1	3.3	3.2	2.8	2.6
Cigars	35.1	34.9	7.1	6.3	3.9	3.4
Pipe Tobacco ²	17.2	17.2	--	--	0.7	0.6
ALCOHOL	87.2	87.4	66.6	66.8	52.7	53.1
Binge Alcohol Use ³	--	--	--	--	18.6	18.4
Heavy Alcohol Use ³	--	--	--	--	5.1	4.6

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	94,090	93,794	49,864	48,509	41,622	40,232
Cigarettes	85,396	84,626	37,012	35,937	31,378	30,331
Smokeless Tobacco	38,582	39,296	11,372	10,711	8,367	7,676
Cigars	65,233	64,738	20,181	19,392	10,583	10,179
Pipe Tobacco ²	30,236	29,886	--	--	1,739	1,728
ALCOHOL	105,494	106,455	85,674	86,725	70,765	70,736
Binge Alcohol Use ³	--	--	--	--	38,116	37,375
Heavy Alcohol Use ³	--	--	--	--	12,460	11,663

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	76.2	75.3	40.4 ^a	38.9	33.7 ^a	32.3
Cigarettes	69.2 ^a	67.9	30.0	28.8	25.4	24.3
Smokeless Tobacco	31.3	31.5	9.2	8.6	6.8	6.2
Cigars	52.8	51.9	16.4	15.6	8.6	8.2
Pipe Tobacco ²	24.5	24.0	--	--	1.4	1.4
ALCOHOL	85.5	85.4	69.4	69.6	57.3	56.8
Binge Alcohol Use ³	--	--	--	--	30.9	30.0
Heavy Alcohol Use ³	--	--	--	--	10.1	9.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	80,319	80,096	33,520	33,349	28,041	27,993
Cigarettes	77,544	77,173	31,587	31,189	26,957	26,488
Smokeless Tobacco	6,454	6,973	1,012	1,035	511	568
Cigars	23,527	23,767	5,880	5,588	2,691	2,686
Pipe Tobacco ²	4,144	4,645	--	--	416	410
ALCOHOL	103,770 ^a	105,292	82,708	83,697	60,609 ^a	62,649
Binge Alcohol Use ³	--	--	--	--	20,419	20,966
Heavy Alcohol Use ³	--	--	--	--	4,439	4,202

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	61.7 ^a	60.2	25.7	25.1	21.5	21.1
Cigarettes	59.6 ^a	58.0	24.3	23.5	20.7	19.9
Smokeless Tobacco	5.0	5.2	0.8	0.8	0.4	0.4
Cigars	18.1	17.9	4.5	4.2	2.1	2.0
Pipe Tobacco ²	3.2	3.5	--	--	0.3	0.3
ALCOHOL	79.7	79.2	63.5	62.9	46.6	47.1
Binge Alcohol Use ³	--	--	--	--	15.7	15.8
Heavy Alcohol Use ³	--	--	--	--	3.4	3.2

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.