

Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	874	926	920	900	852	846
Cigarettes	882	927	838	859	780	816
Smokeless Tobacco	700	752	351	326	314	278
Cigars	929	920	510	504	363	373
Pipe Tobacco ²	694	698	--	--	147	144
ALCOHOL	715	684	967	984	990	1,006
Binge Alcohol Use ³	--	--	--	--	752	734
Heavy Alcohol Use ³	--	--	--	--	422	423

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.34	0.36	0.36	0.35	0.34	0.33
Cigarettes	0.35	0.36	0.33	0.33	0.31	0.32
Smokeless Tobacco	0.28	0.29	0.14	0.13	0.12	0.11
Cigars	0.37	0.36	0.20	0.20	0.14	0.14
Pipe Tobacco ²	0.27	0.27	--	--	0.06	0.06
ALCOHOL	0.28	0.27	0.38	0.38	0.39	0.39
Binge Alcohol Use ³	--	--	--	--	0.30	0.29
Heavy Alcohol Use ³	--	--	--	--	0.17	0.16

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.0080		0.0213		0.0299	
Cigarettes	0.0028		0.0321		0.0308	
Smokeless Tobacco	0.5970		0.0750		0.0585	
Cigars	0.1910		0.0386		0.2401	
Pipe Tobacco ²	0.6847		--		0.8042	
ALCOHOL	0.4099		0.6550		0.9707	
Binge Alcohol Use ³	--		--		0.2901	
Heavy Alcohol Use ³	--		--		0.0256	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	39	35	29	27	21	19
Cigarettes	34	30	26	23	18	16
Smokeless Tobacco	20	17	15	13	11	10
Cigars	19	22	14	17	10	11
Pipe Tobacco ²	9	10	--	--	6	6
ALCOHOL	44	44	37	36	22	20
Binge Alcohol Use ³	--	--	--	--	12	13
Heavy Alcohol Use ³	--	--	--	--	3	2

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.47	0.43	0.35	0.33	0.26	0.23
Cigarettes	0.41	0.36	0.32	0.28	0.23	0.20
Smokeless Tobacco	0.24	0.22	0.19	0.17	0.14	0.12
Cigars	0.23	0.28	0.17	0.21	0.13	0.14
Pipe Tobacco ²	0.12	0.12	--	--	0.07	0.08
ALCOHOL	0.52	0.51	0.45	0.43	0.28	0.25
Binge Alcohol Use ³	--	--	--	--	0.15	0.16
Heavy Alcohol Use ³	--	--	--	--	0.04	0.02

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.1009		0.3379		0.2263	
Cigarettes	0.0616		0.1071		0.1672	
Smokeless Tobacco	0.4170		0.3341		0.4197	
Cigars	0.1860		0.1132		0.6400	
Pipe Tobacco ²	0.9026		--		0.9682	
ALCOHOL	0.2833		0.1295		0.0684	
Binge Alcohol Use ³	--		--		0.8559	
Heavy Alcohol Use ³	--		--		0.1171	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	58	62	49	50	38	39
Cigarettes	51	57	42	45	33	34
Smokeless Tobacco	32	32	26	26	16	15
Cigars	38	40	32	34	19	22
Pipe Tobacco ²	15	18	--	--	8	10
ALCOHOL	75	80	67	70	43	45
Binge Alcohol Use ³	--	--	--	--	32	31
Heavy Alcohol Use ³	--	--	--	--	14	11

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.63	0.66	0.55	0.56	0.45	0.45
Cigarettes	0.58	0.63	0.48	0.51	0.41	0.39
Smokeless Tobacco	0.38	0.36	0.32	0.30	0.20	0.18
Cigars	0.44	0.45	0.38	0.39	0.23	0.25
Pipe Tobacco ²	0.18	0.21	--	--	0.10	0.11
ALCOHOL	0.70	0.74	0.67	0.69	0.49	0.50
Binge Alcohol Use ³	--	--	--	--	0.38	0.35
Heavy Alcohol Use ³	--	--	--	--	0.17	0.13

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.0008		0.0012		0.0088	
Cigarettes	0.0044		0.0016		0.0113	
Smokeless Tobacco	0.0986		0.0169		0.1777	
Cigars	0.0739		0.1335		0.5387	
Pipe Tobacco ²	0.4271		--		0.7991	
ALCOHOL	0.0115		0.0093		0.1117	
Binge Alcohol Use ³	--		--		0.0333	
Heavy Alcohol Use ³	--		--		0.2299	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	89	80	77	70	58	57
Cigarettes	82	73	70	62	52	51
Smokeless Tobacco	43	43	34	36	24	23
Cigars	55	58	46	48	30	33
Pipe Tobacco ²	24	25	--	--	14	14
ALCOHOL	105	102	96	93	68	64
Binge Alcohol Use ³	--	--	--	--	54	46
Heavy Alcohol Use ³	--	--	--	--	24	21

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.78	0.75	0.74	0.69	0.60	0.58
Cigarettes	0.77	0.71	0.70	0.63	0.54	0.54
Smokeless Tobacco	0.48	0.47	0.39	0.40	0.27	0.27
Cigars	0.56	0.59	0.49	0.51	0.34	0.37
Pipe Tobacco ²	0.28	0.28	--	--	0.16	0.16
ALCOHOL	0.76	0.77	0.76	0.79	0.69	0.63
Binge Alcohol Use ³	--	--	--	--	0.57	0.50
Heavy Alcohol Use ³	--	--	--	--	0.28	0.24

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.0619		0.2443		0.7350	
Cigarettes	0.4031		0.6419		0.9205	
Smokeless Tobacco	0.3291		0.6735		0.4243	
Cigars	0.5275		0.5757		0.1756	
Pipe Tobacco ²	0.9723		--		0.3104	
ALCOHOL	0.9081		0.7223		0.4781	
Binge Alcohol Use ³	--		--		0.6029	
Heavy Alcohol Use ³	--		--		0.6249	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	200	219	168	183	140	145
Cigarettes	182	193	147	157	123	128
Smokeless Tobacco	96	86	76	61	61	44
Cigars	142	153	113	117	72	73
Pipe Tobacco ²	62	60	--	--	31	31
ALCOHOL	232	274	218	262	178	218
Binge Alcohol Use ³	--	--	--	--	142	168
Heavy Alcohol Use ³	--	--	--	--	79	92

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.74	0.76	0.74	0.78	0.72	0.76
Cigarettes	0.77	0.79	0.73	0.78	0.70	0.72
Smokeless Tobacco	0.60	0.58	0.50	0.44	0.42	0.32
Cigars	0.72	0.74	0.63	0.64	0.46	0.47
Pipe Tobacco ²	0.42	0.42	--	--	0.22	0.23
ALCOHOL	0.61	0.71	0.65	0.83	0.74	0.91
Binge Alcohol Use ³	--	--	--	--	0.71	0.85
Heavy Alcohol Use ³	--	--	--	--	0.51	0.58

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.0182		0.0675		0.0374	
Cigarettes	0.0291		0.2340		0.5690	
Smokeless Tobacco	0.2117		0.0219		0.0006	
Cigars	0.0158		0.2648		0.3777	
Pipe Tobacco ²	0.9006		--		0.3032	
ALCOHOL	0.0005		0.0067		0.1211	
Binge Alcohol Use ³	--		--		0.0846	
Heavy Alcohol Use ³	--		--		0.0406	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	262	285	210	219	178	184
Cigarettes	243	258	180	193	156	168
Smokeless Tobacco	122	121	75	76	57	59
Cigars	193	202	121	130	78	86
Pipe Tobacco ²	83	93	--	--	28	36
ALCOHOL	331	350	322	343	279	304
Binge Alcohol Use ³	--	--	--	--	197	225
Heavy Alcohol Use ³	--	--	--	--	110	100

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.59	0.55	0.63	0.62	0.61	0.59
Cigarettes	0.63	0.60	0.63	0.61	0.58	0.57
Smokeless Tobacco	0.49	0.46	0.33	0.33	0.26	0.26
Cigars	0.59	0.61	0.49	0.49	0.37	0.36
Pipe Tobacco ²	0.38	0.39	--	--	0.14	0.17
ALCOHOL	0.36	0.38	0.48	0.50	0.61	0.59
Binge Alcohol Use ³	--	--	--	--	0.63	0.66
Heavy Alcohol Use ³	--	--	--	--	0.47	0.41

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.8704		0.3148		0.1878	
Cigarettes	0.4170		0.3867		0.1613	
Smokeless Tobacco	0.6052		0.0959		0.2662	
Cigars	0.2002		0.5909		0.4478	
Pipe Tobacco ²	0.7289		--		0.1894	
ALCOHOL	0.4610		0.1924		0.7660	
Binge Alcohol Use ³	--		--		0.9522	
Heavy Alcohol Use ³	--		--		0.0220	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	269	283	317	316	306	298
Cigarettes	273	298	305	305	291	286
Smokeless Tobacco	237	259	152	131	132	108
Cigars	298	314	222	219	149	172
Pipe Tobacco ²	172	162	--	--	52	49
ALCOHOL	180	195	264	263	306	304
Binge Alcohol Use ³	--	--	--	--	302	298
Heavy Alcohol Use ³	--	--	--	--	185	200

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.74	0.78	0.87	0.87	0.84	0.82
Cigarettes	0.75	0.82	0.84	0.84	0.80	0.78
Smokeless Tobacco	0.65	0.71	0.42	0.36	0.36	0.30
Cigars	0.82	0.86	0.61	0.60	0.41	0.47
Pipe Tobacco ²	0.47	0.44	--	--	0.14	0.13
ALCOHOL	0.49	0.53	0.72	0.72	0.84	0.84
Binge Alcohol Use ³	--	--	--	--	0.83	0.82
Heavy Alcohol Use ³	--	--	--	--	0.51	0.55

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.0095		0.3287		0.2163	
Cigarettes	0.0088		0.3888		0.0653	
Smokeless Tobacco	0.8896		0.4888		0.2881	
Cigars	0.1666		0.9413		0.2117	
Pipe Tobacco ²	0.0756		--		0.9516	
ALCOHOL	0.4158		0.9183		0.5572	
Binge Alcohol Use ³	--		--		0.5562	
Heavy Alcohol Use ³	--		--		0.7259	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	771	819	792	756	747	716
Cigarettes	784	825	728	709	686	685
Smokeless Tobacco	620	656	277	266	254	241
Cigars	823	811	428	396	309	302
Pipe Tobacco ²	660	661	--	--	131	126
ALCOHOL	649	609	866	901	875	921
Binge Alcohol Use ³	--	--	--	--	624	604
Heavy Alcohol Use ³	--	--	--	--	347	324

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.49	0.51	0.50	0.47	0.47	0.44
Cigarettes	0.49	0.51	0.46	0.44	0.43	0.42
Smokeless Tobacco	0.39	0.41	0.17	0.16	0.16	0.15
Cigars	0.52	0.50	0.27	0.24	0.19	0.19
Pipe Tobacco ²	0.42	0.41	--	--	0.08	0.08
ALCOHOL	0.41	0.38	0.55	0.56	0.55	0.57
Binge Alcohol Use ³	--	--	--	--	0.39	0.37
Heavy Alcohol Use ³	--	--	--	--	0.22	0.20

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.2231		0.1858		0.2676	
Cigarettes	0.1168		0.1733		0.2774	
Smokeless Tobacco	0.2709		0.6787		0.6299	
Cigars	0.8132		0.0532		0.0986	
Pipe Tobacco ²	0.9409		--		0.6695	
ALCOHOL	0.7472		0.7316		0.5502	
Binge Alcohol Use ³	--		--		0.7336	
Heavy Alcohol Use ³	--		--		0.0932	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	506	558	667	605	626	581
Cigarettes	541	588	578	560	536	537
Smokeless Tobacco	613	652	337	311	302	264
Cigars	672	684	448	436	321	336
Pipe Tobacco ²	607	623	--	--	137	135
ALCOHOL	395	408	628	630	675	674
Binge Alcohol Use ³	--	--	--	--	577	567
Heavy Alcohol Use ³	--	--	--	--	353	355

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.41	0.45	0.54	0.49	0.51	0.47
Cigarettes	0.44	0.47	0.47	0.45	0.43	0.43
Smokeless Tobacco	0.50	0.52	0.27	0.25	0.25	0.21
Cigars	0.54	0.55	0.36	0.35	0.26	0.27
Pipe Tobacco ²	0.49	0.50	--	--	0.11	0.11
ALCOHOL	0.32	0.33	0.51	0.51	0.55	0.54
Binge Alcohol Use ³	--	--	--	--	0.47	0.45
Heavy Alcohol Use ³	--	--	--	--	0.29	0.28

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.1030		0.0339		0.0315	
Cigarettes	0.0425		0.0699		0.0713	
Smokeless Tobacco	0.7034		0.0883		0.0519	
Cigars	0.2223		0.1126		0.2828	
Pipe Tobacco ²	0.4412		--		0.8822	
ALCOHOL	0.9164		0.8009		0.4468	
Binge Alcohol Use ³	--		--		0.1773	
Heavy Alcohol Use ³	--		--		0.0569	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	648	660	572	564	535	542
Cigarettes	656	664	560	556	526	532
Smokeless Tobacco	245	281	89	97	78	82
Cigars	488	479	207	197	153	142
Pipe Tobacco ²	226	234	--	--	54	46
ALCOHOL	540	530	656	690	635	684
Binge Alcohol Use ³	--	--	--	--	441	441
Heavy Alcohol Use ³	--	--	--	--	201	207

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2010 and 2011

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.50	0.50	0.44	0.42	0.41	0.41
Cigarettes	0.50	0.50	0.43	0.42	0.40	0.40
Smokeless Tobacco	0.19	0.21	0.07	0.07	0.06	0.06
Cigars	0.37	0.36	0.16	0.15	0.12	0.11
Pipe Tobacco ²	0.17	0.18	--	--	0.04	0.03
ALCOHOL	0.41	0.40	0.50	0.52	0.49	0.51
Binge Alcohol Use ³	--	--	--	--	0.34	0.33
Heavy Alcohol Use ³	--	--	--	--	0.15	0.16

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOBACCO PRODUCTS¹	0.0312		0.2760		0.3996	
Cigarettes	0.0264		0.1860		0.1714	
Smokeless Tobacco	0.3040		0.9871		0.6831	
Cigars	0.6943		0.1364		0.7643	
Pipe Tobacco ²	0.2216		--		0.8332	
ALCOHOL	0.3460		0.4018		0.4196	
Binge Alcohol Use ³	--		--		0.8566	
Heavy Alcohol Use ³	--		--		0.2564	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.