

Table 2.47C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	874	926	920	900	852	846
GEOGRAPHIC DIVISION						
Northeast	337	398	332	394	309	364
New England	169	191	153	190	139	171
Middle Atlantic	288	347	292	346	275	321
Midwest	317	308	381	355	369	345
East North Central	266	260	303	293	287	288
West North Central	169	165	217	199	218	192
South	563	552	623	558	589	522
South Atlantic	441	429	461	430	439	394
East South Central	215	198	227	218	217	221
West South Central	282	280	291	269	287	259
West	471	524	453	456	409	425
Mountain	196	220	222	212	208	207
Pacific	427	471	396	397	355	362
COUNTY TYPE						
Large Metro	1,618	1,579	892	859	787	765
Small Metro	1,235	1,238	697	623	618	556
250K – 1 Mil. Pop.	995	1,015	532	488	471	443
< 250K Pop.	747	776	426	420	378	367
Nonmetro	898	814	550	486	496	445
Urbanized	716	527	356	298	324	275
Less Urbanized	650	674	411	381	368	349
Completely Rural	329	287	170	156	153	148

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.47D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.34	0.36	0.36	0.35	0.34	0.33
GEOGRAPHIC DIVISION						
Northeast	0.73	0.85	0.71	0.84	0.66	0.78
New England	1.38	1.55	1.25	1.55	1.13	1.39
Middle Atlantic	0.84	1.00	0.85	1.00	0.80	0.93
Midwest	0.57	0.55	0.69	0.64	0.67	0.62
East North Central	0.69	0.67	0.79	0.76	0.75	0.74
West North Central	1.00	0.97	1.29	1.17	1.30	1.13
South	0.61	0.58	0.67	0.59	0.63	0.55
South Atlantic	0.90	0.85	0.94	0.86	0.90	0.79
East South Central	1.42	1.29	1.51	1.42	1.44	1.44
West South Central	0.98	0.94	1.01	0.91	0.99	0.87
West	0.80	0.88	0.77	0.76	0.70	0.71
Mountain	1.09	1.21	1.23	1.17	1.16	1.14
Pacific	1.05	1.13	0.97	0.95	0.87	0.87
COUNTY TYPE						
Large Metro	0.50	0.54	0.50	0.49	0.47	0.46
Small Metro	0.61	0.58	0.65	0.58	0.61	0.56
250K – 1 Mil. Pop.	0.74	0.74	0.72	0.74	0.70	0.71
< 250K Pop.	1.07	0.92	1.21	1.01	1.11	0.94
Nonmetro	0.76	0.77	0.92	0.82	0.87	0.78
Urbanized	1.20	1.25	1.42	1.28	1.36	1.20
Less Urbanized	1.11	1.04	1.29	1.24	1.23	1.18
Completely Rural	2.39	1.99	2.50	2.14	2.33	2.12

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.47P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0080		0.0213		0.0299	
GEOGRAPHIC DIVISION						
Northeast	0.0060		0.9478		0.7777	
New England	0.2237		0.3485		0.4857	
Middle Atlantic	0.0120		0.6453		0.4908	
Midwest	0.1224		0.2457		0.5817	
East North Central	0.0468		0.4512		0.9740	
West North Central	0.9007		0.3299		0.3178	
South	0.7198		0.1166		0.1063	
South Atlantic	0.4710		0.3601		0.1617	
East South Central	0.8284		0.2957		0.3889	
West South Central	0.1221		0.3548		0.6473	
West	0.1690		0.0946		0.1032	
Mountain	0.0638		0.4787		0.6714	
Pacific	0.4999		0.1371		0.1124	
COUNTY TYPE						
Large Metro	0.0023		0.0509		0.0241	
Small Metro	0.5437		0.3847		0.6120	
250K – 1 Mil. Pop.	0.2804		0.6635		0.7206	
< 250K Pop.	0.6158		0.3979		0.7391	
Nonmetro	0.1305		0.3593		0.5921	
Urbanized	0.0165		0.8437		0.8743	
Less Urbanized	0.9333		0.1672		0.2954	
Completely Rural	0.8672		0.4194		0.2308	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.48C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	101	99	89	85	69	68
GEOGRAPHIC DIVISION						
Northeast	36	40	32	37	25	28
New England	15	17	14	17	12	14
Middle Atlantic	32	36	29	32	22	24
Midwest	36	37	33	33	27	28
East North Central	30	31	27	27	21	22
West North Central	21	21	19	19	17	17
South	68	60	62	51	48	40
South Atlantic	52	42	48	37	39	30
East South Central	21	21	19	18	14	15
West South Central	35	36	32	30	23	23
West	53	53	48	45	34	33
Mountain	26	25	23	22	18	17
Pacific	46	46	42	39	29	29
COUNTY TYPE						
Large Metro	94	84	80	70	57	53
Small Metro	70	67	53	57	38	43
250K – 1 Mil. Pop.	58	56	45	48	32	35
< 250K Pop.	40	39	31	32	22	25
Nonmetro	55	51	46	37	36	29
Urbanized	35	33	27	25	21	18
Less Urbanized	45	39	37	28	29	23
Completely Rural	12	16	11	13	9	10

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.48D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.42	0.40	0.37	0.34	0.28	0.27
GEOGRAPHIC DIVISION						
Northeast	0.84	0.93	0.77	0.86	0.60	0.67
New England	1.39	1.57	1.29	1.54	1.06	1.29
Middle Atlantic	1.02	1.12	0.92	1.01	0.71	0.77
Midwest	0.68	0.68	0.61	0.60	0.50	0.51
East North Central	0.79	0.81	0.72	0.70	0.56	0.58
West North Central	1.34	1.26	1.21	1.18	1.08	1.02
South	0.76	0.65	0.69	0.55	0.53	0.43
South Atlantic	1.17	0.91	1.08	0.80	0.88	0.65
East South Central	1.43	1.42	1.31	1.22	0.97	1.01
West South Central	1.16	1.15	1.05	0.94	0.75	0.71
West	0.90	0.88	0.82	0.75	0.59	0.55
Mountain	1.47	1.38	1.27	1.20	1.00	0.90
Pacific	1.13	1.11	1.05	0.95	0.72	0.69
COUNTY TYPE						
Large Metro	0.59	0.53	0.53	0.46	0.40	0.37
Small Metro	0.67	0.68	0.57	0.63	0.45	0.49
250K – 1 Mil. Pop.	0.82	0.83	0.72	0.76	0.57	0.58
< 250K Pop.	1.17	1.23	1.01	1.12	0.81	0.87
Nonmetro	1.02	0.97	0.96	0.79	0.81	0.70
Urbanized	1.48	1.34	1.32	1.09	1.03	0.93
Less Urbanized	1.54	1.55	1.52	1.30	1.32	1.10
Completely Rural	3.07	3.05	2.86	2.46	2.57	2.25

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.48P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0005		0.0064		0.0468	
GEOGRAPHIC DIVISION						
Northeast	0.1087		0.7527		0.3864	
New England	0.4724		0.8760		0.2869	
Middle Atlantic	0.1490		0.6596		0.6795	
Midwest	0.3669		0.1502		0.1615	
East North Central	0.4688		0.1867		0.4381	
West North Central	0.5996		0.5594		0.2032	
South	0.0019		0.0150		0.1512	
South Atlantic	0.0080		0.1251		0.1844	
East South Central	0.1700		0.6794		0.7962	
West South Central	0.2444		0.0512		0.3994	
West	0.3083		0.3546		0.0803	
Mountain	0.1140		0.1991		0.1078	
Pacific	0.7378		0.7028		0.2707	
COUNTY TYPE						
Large Metro	0.0122		0.0209		0.0581	
Small Metro	0.0081		0.3696		0.9084	
250K – 1 Mil. Pop.	0.0038		0.1215		0.4038	
< 250K Pop.	0.7635		0.4415		0.2740	
Nonmetro	0.5147		0.1981		0.2742	
Urbanized	0.8259		0.6853		0.6213	
Less Urbanized	0.3329		0.0835		0.1839	
Completely Rural	0.7358		0.8385		0.5135	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.49C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	159	160	171	171	166	167
GEOGRAPHIC DIVISION						
Northeast	65	66	70	67	66	67
New England	30	31	35	28	32	31
Middle Atlantic	57	59	60	61	58	59
Midwest	58	57	63	71	65	71
East North Central	48	46	51	60	53	58
West North Central	31	32	35	36	38	40
South	105	102	108	109	106	103
South Atlantic	84	74	85	79	83	79
East South Central	37	37	36	45	37	46
West South Central	50	57	57	61	55	49
West	86	86	96	92	90	92
Mountain	38	38	41	40	40	40
Pacific	76	78	87	83	81	82
COUNTY TYPE						
Large Metro	282	292	227	227	192	192
Small Metro	242	266	194	205	154	156
250K – 1 Mil. Pop.	173	163	137	123	108	106
< 250K Pop.	170	214	138	165	111	116
Nonmetro	137	131	118	109	104	95
Urbanized	98	95	82	79	73	70
Less Urbanized	100	90	88	72	76	63
Completely Rural	37	39	29	35	27	32

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.49D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.47	0.47	0.50	0.50	0.49	0.49
GEOGRAPHIC DIVISION						
Northeast	1.07	1.09	1.16	1.09	1.10	1.09
New England	1.86	1.90	2.15	1.73	1.97	1.92
Middle Atlantic	1.29	1.30	1.37	1.34	1.31	1.31
Midwest	0.78	0.78	0.84	0.96	0.87	0.96
East North Central	0.94	0.91	1.00	1.18	1.03	1.15
West North Central	1.33	1.42	1.49	1.59	1.60	1.75
South	0.85	0.81	0.87	0.87	0.85	0.82
South Atlantic	1.33	1.15	1.35	1.22	1.31	1.22
East South Central	1.87	1.86	1.81	2.25	1.88	2.28
West South Central	1.22	1.37	1.38	1.46	1.33	1.18
West	1.06	1.05	1.18	1.12	1.11	1.12
Mountain	1.51	1.53	1.62	1.63	1.61	1.61
Pacific	1.36	1.35	1.55	1.43	1.45	1.43
COUNTY TYPE						
Large Metro	0.69	0.65	0.72	0.69	0.69	0.68
Small Metro	0.82	0.80	0.90	0.86	0.88	0.85
250K – 1 Mil. Pop.	0.93	0.96	1.11	1.06	1.11	1.05
< 250K Pop.	1.46	1.34	1.50	1.37	1.46	1.37
Nonmetro	0.89	0.99	1.05	1.13	1.10	1.14
Urbanized	1.36	1.41	1.51	1.83	1.66	1.78
Less Urbanized	1.29	1.49	1.61	1.47	1.59	1.54
Completely Rural	3.90	3.59	3.29	3.79	3.48	4.44

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.49P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.1734		0.0588		0.0253	
GEOGRAPHIC DIVISION						
Northeast	0.2239		0.4226		0.1586	
New England	0.6316		0.6774		0.1303	
Middle Atlantic	0.2630		0.4998		0.4387	
Midwest	0.2059		0.1764		0.1166	
East North Central	0.3459		0.2435		0.0978	
West North Central	0.3781		0.4695		0.7219	
South	0.8126		0.1986		0.0710	
South Atlantic	0.8160		0.4024		0.1085	
East South Central	0.3626		0.5992		0.4506	
West South Central	0.1411		0.1137		0.0871	
West	0.6971		0.7462		0.7857	
Mountain	0.5170		0.9866		0.4397	
Pacific	0.9251		0.7254		0.9513	
COUNTY TYPE						
Large Metro	0.8583		0.6234		0.4555	
Small Metro	0.1267		0.0833		0.0943	
250K – 1 Mil. Pop.	0.3605		0.2380		0.2774	
< 250K Pop.	0.2025		0.2035		0.2158	
Nonmetro	0.2099		0.1066		0.0413	
Urbanized	0.1286		0.0177		0.0089	
Less Urbanized	0.9562		0.8579		0.6297	
Completely Rural	0.4050		0.5903		0.9030	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.50C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	818	877	870	857	813	805
GEOGRAPHIC DIVISION						
Northeast	313	386	316	376	297	346
New England	148	180	138	175	126	159
Middle Atlantic	272	340	281	334	267	308
Midwest	298	291	362	339	350	332
East North Central	247	246	288	281	274	276
West North Central	162	154	204	188	205	185
South	539	512	594	533	566	505
South Atlantic	424	396	438	403	419	376
East South Central	205	184	218	208	209	211
West South Central	271	261	275	258	272	252
West	431	493	425	425	386	393
Mountain	182	207	204	196	196	191
Pacific	392	443	376	370	336	334
COUNTY TYPE						
Large Metro	1,522	1,488	813	781	731	698
Small Metro	1,132	1,134	607	545	550	507
250K – 1 Mil. Pop.	922	956	472	441	425	410
< 250K Pop.	675	687	367	350	332	323
Nonmetro	827	742	492	429	448	400
Urbanized	652	475	313	256	289	243
Less Urbanized	591	612	357	337	327	313
Completely Rural	304	250	151	133	134	130

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.50D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.42	0.44	0.45	0.43	0.42	0.41
GEOGRAPHIC DIVISION						
Northeast	0.86	1.06	0.87	1.03	0.82	0.95
New England	1.56	1.87	1.45	1.83	1.32	1.66
Middle Atlantic	1.02	1.26	1.05	1.24	1.00	1.15
Midwest	0.70	0.68	0.85	0.79	0.82	0.77
East North Central	0.83	0.83	0.97	0.94	0.92	0.93
West North Central	1.26	1.18	1.59	1.44	1.59	1.41
South	0.75	0.70	0.83	0.73	0.79	0.69
South Atlantic	1.11	1.01	1.15	1.03	1.10	0.96
East South Central	1.76	1.56	1.87	1.76	1.79	1.79
West South Central	1.25	1.17	1.26	1.16	1.25	1.13
West	0.96	1.08	0.95	0.93	0.86	0.86
Mountain	1.32	1.50	1.48	1.41	1.43	1.38
Pacific	1.26	1.39	1.21	1.16	1.08	1.05
COUNTY TYPE						
Large Metro	0.60	0.68	0.62	0.61	0.58	0.57
Small Metro	0.73	0.68	0.78	0.71	0.74	0.69
250K – 1 Mil. Pop.	0.90	0.86	0.88	0.90	0.85	0.87
< 250K Pop.	1.32	1.14	1.49	1.24	1.37	1.19
Nonmetro	0.96	0.93	1.12	0.98	1.07	0.94
Urbanized	1.47	1.56	1.72	1.49	1.66	1.46
Less Urbanized	1.41	1.25	1.58	1.50	1.54	1.43
Completely Rural	2.97	2.32	2.96	2.63	2.72	2.60

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.50P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0416		0.0937		0.1136	
GEOGRAPHIC DIVISION						
Northeast	0.0151		0.7981		0.9267	
New England	0.2463		0.2468		0.3188	
Middle Atlantic	0.0311		0.7473		0.5375	
Midwest	0.2361		0.4802		0.9495	
East North Central	0.0806		0.7257		0.6197	
West North Central	0.6709		0.4594		0.4395	
South	0.9213		0.2712		0.2376	
South Atlantic	0.3064		0.5274		0.3123	
East South Central	0.7999		0.2488		0.2912	
West South Central	0.2444		0.7300		0.9747	
West	0.2222		0.1169		0.1064	
Mountain	0.1283		0.5689		0.6671	
Pacific	0.5235		0.1506		0.1165	
COUNTY TYPE						
Large Metro	0.0054		0.0903		0.0444	
Small Metro	0.2130		0.7335		0.9212	
250K – 1 Mil. Pop.	0.0954		0.9357		0.9923	
< 250K Pop.	0.7983		0.5062		0.8790	
Nonmetro	0.2280		0.5998		0.9479	
Urbanized	0.0552		0.8707		0.7542	
Less Urbanized	0.9916		0.2532		0.3949	
Completely Rural	0.7888		0.5043		0.2388	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.51C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	844	897	906	895	840	845
GEOGRAPHIC DIVISION						
Northeast	324	395	329	393	309	363
New England	157	186	153	187	140	169
Middle Atlantic	279	346	289	347	275	321
Midwest	306	296	379	352	367	343
East North Central	254	250	301	291	283	287
West North Central	168	157	216	196	220	190
South	551	525	618	555	583	521
South Atlantic	432	407	454	427	432	395
East South Central	210	190	225	216	216	218
West South Central	276	268	288	267	284	257
West	451	511	446	451	402	421
Mountain	185	211	219	209	207	205
Pacific	411	461	390	392	348	358
COUNTY TYPE						
Large Metro	1,595	1,560	874	843	775	753
Small Metro	1,212	1,220	682	611	609	551
250K – 1 Mil. Pop.	975	1,002	520	477	464	437
< 250K Pop.	731	758	416	412	372	361
Nonmetro	873	792	533	473	486	437
Urbanized	696	511	345	288	318	270
Less Urbanized	630	654	395	369	360	341
Completely Rural	323	279	164	151	148	144

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.51D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.37	0.39	0.40	0.38	0.37	0.36
GEOGRAPHIC DIVISION						
Northeast	0.77	0.93	0.78	0.92	0.73	0.85
New England	1.41	1.66	1.38	1.67	1.25	1.51
Middle Atlantic	0.90	1.10	0.93	1.10	0.88	1.02
Midwest	0.61	0.59	0.76	0.70	0.73	0.68
East North Central	0.73	0.72	0.87	0.83	0.81	0.82
West North Central	1.11	1.02	1.42	1.28	1.44	1.23
South	0.66	0.61	0.74	0.65	0.69	0.61
South Atlantic	0.97	0.89	1.02	0.94	0.97	0.87
East South Central	1.54	1.37	1.65	1.56	1.58	1.57
West South Central	1.07	1.01	1.12	1.01	1.10	0.97
West	0.85	0.95	0.84	0.84	0.76	0.78
Mountain	1.14	1.29	1.35	1.28	1.27	1.26
Pacific	1.12	1.23	1.06	1.04	0.95	0.96
COUNTY TYPE						
Large Metro	0.53	0.59	0.54	0.54	0.51	0.51
Small Metro	0.65	0.60	0.71	0.63	0.67	0.62
250K – 1 Mil. Pop.	0.79	0.77	0.80	0.81	0.77	0.78
< 250K Pop.	1.14	0.96	1.32	1.10	1.21	1.02
Nonmetro	0.84	0.83	1.00	0.90	0.96	0.86
Urbanized	1.30	1.35	1.54	1.41	1.49	1.33
Less Urbanized	1.22	1.11	1.42	1.36	1.37	1.29
Completely Rural	2.67	2.15	2.69	2.35	2.50	2.33

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.51P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0270		0.0437		0.0482	
GEOGRAPHIC DIVISION						
Northeast	0.0095		0.9191		0.7264	
New England	0.2435		0.3459		0.5346	
Middle Atlantic	0.0185		0.6768		0.4691	
Midwest	0.1661		0.3128		0.6791	
East North Central	0.0592		0.5422		0.9070	
West North Central	0.7884		0.3584		0.3792	
South	0.9541		0.1892		0.1380	
South Atlantic	0.2930		0.4492		0.2031	
East South Central	0.9603		0.3093		0.3801	
West South Central	0.1588		0.4912		0.7151	
West	0.2136		0.1160		0.1408	
Mountain	0.1013		0.5809		0.7956	
Pacific	0.5298		0.1458		0.1333	
COUNTY TYPE						
Large Metro	0.0060		0.0878		0.0380	
Small Metro	0.3407		0.4332		0.6199	
250K – 1 Mil. Pop.	0.1343		0.7872		0.7813	
< 250K Pop.	0.6052		0.3520		0.6649	
Nonmetro	0.1669		0.4434		0.6845	
Urbanized	0.0345		0.8997		0.9530	
Less Urbanized	0.9741		0.2061		0.3212	
Completely Rural	0.9066		0.4057		0.2018	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.