

**Table 2.67C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	715	684	967	984	990	1,006
<b>GEOGRAPHIC DIVISION</b>						
Northeast	274	276	362	383	403	420
New England	133	110	169	152	204	204
Middle Atlantic	237	253	317	350	340	365
Midwest	254	232	343	355	380	381
East North Central	221	188	285	307	311	322
West North Central	127	137	211	192	239	208
South	470	451	653	590	658	603
South Atlantic	352	332	493	473	495	466
East South Central	213	169	281	195	266	226
West South Central	225	235	322	304	353	322
West	415	390	551	566	549	588
Mountain	183	166	262	243	246	263
Pacific	373	352	480	507	492	521
<b>COUNTY TYPE</b>						
Large Metro	1,846	1,835	1,627	1,596	1,389	1,393
Small Metro	1,458	1,414	1,233	1,218	1,023	1,040
250K – 1 Mil. Pop.	1,185	1,147	1,006	968	831	820
< 250K Pop.	859	897	718	770	604	654
Nonmetro	980	918	819	726	685	595
Urbanized	760	643	654	504	554	412
Less Urbanized	723	742	576	576	461	481
Completely Rural	337	329	250	231	203	184

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.67D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.28	0.27	0.38	0.38	0.39	0.39
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.59	0.59	0.78	0.82	0.87	0.89
New England	1.09	0.89	1.38	1.23	1.67	1.65
Middle Atlantic	0.69	0.73	0.93	1.01	0.99	1.06
Midwest	0.46	0.42	0.62	0.64	0.69	0.68
East North Central	0.57	0.49	0.74	0.79	0.81	0.83
West North Central	0.76	0.81	1.25	1.13	1.42	1.23
South	0.51	0.47	0.70	0.62	0.71	0.63
South Atlantic	0.72	0.66	1.01	0.94	1.01	0.93
East South Central	1.42	1.10	1.86	1.27	1.77	1.48
West South Central	0.78	0.79	1.12	1.02	1.22	1.08
West	0.71	0.65	0.94	0.95	0.93	0.98
Mountain	1.02	0.91	1.45	1.34	1.37	1.45
Pacific	0.92	0.84	1.18	1.22	1.21	1.25
<b>COUNTY TYPE</b>						
Large Metro	0.41	0.38	0.53	0.52	0.56	0.56
Small Metro	0.50	0.48	0.65	0.66	0.70	0.67
250K – 1 Mil. Pop.	0.54	0.61	0.75	0.85	0.83	0.86
< 250K Pop.	1.00	0.74	1.27	1.03	1.30	1.09
Nonmetro	0.69	0.62	0.96	0.85	0.95	0.89
Urbanized	0.97	0.92	1.59	1.19	1.54	1.37
Less Urbanized	1.04	0.88	1.35	1.28	1.28	1.34
Completely Rural	2.20	2.00	2.96	2.32	2.89	2.56

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.67P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.4099		0.6550		0.9707	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.3327		0.6001		0.4730	
New England	0.9810		0.5951		0.5126	
Middle Atlantic	0.2838		0.7417		0.6570	
Midwest	0.7684		0.2150		0.3587	
East North Central	0.9730		0.3598		0.3701	
West North Central	0.5563		0.4131		0.7764	
South	0.9679		0.6801		0.2190	
South Atlantic	0.5658		0.9635		0.7278	
East South Central	0.7680		0.2671		0.6113	
West South Central	0.5201		0.8981		0.1583	
West	0.5821		0.4124		0.8024	
Mountain	0.4709		0.4000		0.7985	
Pacific	0.3361		0.1819		0.6707	
<b>COUNTY TYPE</b>						
Large Metro	0.1391		0.4679		0.8373	
Small Metro	0.4303		0.3850		0.3730	
250K – 1 Mil. Pop.	0.8975		0.5761		0.7678	
< 250K Pop.	0.1087		0.0191		0.0493	
Nonmetro	0.7661		0.3068		0.0530	
Urbanized	0.0917		0.6302		0.0265	
Less Urbanized	0.2942		0.4273		0.7051	
Completely Rural	0.9675		0.7320		0.5383	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.68C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	108	110	104	107	80	77
<b>GEOGRAPHIC DIVISION</b>						
Northeast	40	42	37	43	31	39
New England	19	18	17	18	14	14
Middle Atlantic	35	38	33	39	28	36
Midwest	40	41	38	39	30	32
East North Central	33	34	30	31	22	26
West North Central	24	23	24	24	21	19
South	72	67	68	64	53	44
South Atlantic	52	45	51	44	41	30
East South Central	23	24	22	23	16	14
West South Central	40	41	39	37	28	28
West	62	63	60	58	45	41
Mountain	27	28	25	28	19	18
Pacific	55	56	54	51	40	36
<b>COUNTY TYPE</b>						
Large Metro	118	130	105	113	73	69
Small Metro	87	85	74	74	46	47
250K – 1 Mil. Pop.	71	74	61	64	36	39
< 250K Pop.	52	44	43	40	28	26
Nonmetro	61	52	55	46	36	29
Urbanized	43	36	39	31	23	21
Less Urbanized	46	38	41	34	27	21
Completely Rural	15	19	13	16	9	10

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.68D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.44	0.44	0.43	0.43	0.33	0.31
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.93	0.98	0.88	1.01	0.73	0.90
New England	1.73	1.64	1.56	1.60	1.26	1.25
Middle Atlantic	1.10	1.19	1.05	1.25	0.88	1.14
Midwest	0.75	0.75	0.70	0.72	0.56	0.58
East North Central	0.87	0.89	0.80	0.82	0.58	0.67
West North Central	1.52	1.42	1.52	1.44	1.32	1.15
South	0.80	0.73	0.76	0.69	0.59	0.47
South Atlantic	1.16	0.97	1.14	0.96	0.91	0.65
East South Central	1.62	1.66	1.51	1.54	1.08	0.97
West South Central	1.34	1.29	1.28	1.18	0.94	0.88
West	1.06	1.05	1.03	0.97	0.77	0.68
Mountain	1.52	1.54	1.37	1.50	1.09	0.96
Pacific	1.37	1.35	1.35	1.23	1.00	0.88
<b>COUNTY TYPE</b>						
Large Metro	0.64	0.66	0.62	0.63	0.50	0.46
Small Metro	0.76	0.72	0.73	0.68	0.53	0.52
250K – 1 Mil. Pop.	0.92	0.92	0.90	0.86	0.62	0.63
< 250K Pop.	1.29	1.17	1.25	1.12	0.98	0.92
Nonmetro	1.01	0.92	0.96	0.90	0.79	0.68
Urbanized	1.50	1.29	1.41	1.31	0.96	1.00
Less Urbanized	1.51	1.46	1.50	1.38	1.28	1.03
Completely Rural	3.92	3.22	3.41	2.81	2.60	2.30

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.68P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.1372		0.1481		0.4804	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.7729		0.9423		0.2107	
New England	0.1687		0.2817		0.6836	
Middle Atlantic	0.3111		0.5431		0.1202	
Midwest	0.2044		0.9944		0.7072	
East North Central	0.9598		0.4578		0.1657	
West North Central	0.0271		0.3500		0.1921	
South	0.0209		0.0149		0.1832	
South Atlantic	0.0011		0.0122		0.1914	
East South Central	0.3217		0.4778		0.9658	
West South Central	0.6408		0.5417		0.5437	
West	0.4979		0.9073		0.3129	
Mountain	0.4624		0.9901		0.2085	
Pacific	0.2775		0.9030		0.6017	
<b>COUNTY TYPE</b>						
Large Metro	0.8372		0.8394		0.6498	
Small Metro	0.0370		0.1189		0.7136	
250K – 1 Mil. Pop.	0.0430		0.0478		0.4865	
< 250K Pop.	0.4940		0.8239		0.7556	
Nonmetro	0.1239		0.1720		0.6854	
Urbanized	0.9834		0.5830		0.7063	
Less Urbanized	0.0417		0.2103		0.7090	
Completely Rural	0.9156		0.9014		0.3384	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.69C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	114	131	138	157	169	184
<b>GEOGRAPHIC DIVISION</b>						
Northeast	44	49	54	62	68	74
New England	22	20	30	29	36	34
Middle Atlantic	38	45	46	55	58	66
Midwest	45	47	50	57	64	71
East North Central	36	40	39	47	50	59
West North Central	28	24	32	30	41	40
South	76	83	93	103	111	121
South Atlantic	61	63	74	78	86	93
East South Central	26	27	32	31	40	36
West South Central	38	42	44	52	55	60
West	63	75	79	83	93	95
Mountain	33	40	39	44	45	52
Pacific	53	64	68	72	80	81
<b>COUNTY TYPE</b>						
Large Metro	344	369	330	351	275	300
Small Metro	300	332	284	320	243	279
250K – 1 Mil. Pop.	219	212	203	202	179	170
< 250K Pop.	201	258	193	250	164	222
Nonmetro	150	154	139	147	121	118
Urbanized	114	115	106	110	92	84
Less Urbanized	109	102	101	98	85	82
Completely Rural	37	48	33	45	28	39

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.69D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.34	0.38	0.41	0.46	0.50	0.54
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.72	0.81	0.90	1.01	1.13	1.20
New England	1.36	1.25	1.82	1.79	2.22	2.09
Middle Atlantic	0.86	0.99	1.03	1.22	1.31	1.45
Midwest	0.60	0.64	0.67	0.77	0.85	0.97
East North Central	0.69	0.79	0.77	0.94	0.98	1.16
West North Central	1.17	1.04	1.38	1.32	1.72	1.78
South	0.62	0.66	0.75	0.81	0.89	0.96
South Atlantic	0.97	0.98	1.18	1.20	1.36	1.45
East South Central	1.33	1.35	1.61	1.57	2.04	1.81
West South Central	0.93	1.01	1.08	1.25	1.33	1.45
West	0.78	0.91	0.98	1.01	1.14	1.15
Mountain	1.31	1.62	1.57	1.78	1.80	2.12
Pacific	0.95	1.11	1.21	1.24	1.42	1.40
<b>COUNTY TYPE</b>						
Large Metro	0.52	0.54	0.61	0.64	0.72	0.71
Small Metro	0.55	0.62	0.69	0.76	0.87	0.99
250K – 1 Mil. Pop.	0.67	0.74	0.86	0.85	1.03	1.07
< 250K Pop.	0.97	1.09	1.18	1.47	1.60	1.99
Nonmetro	0.78	0.89	0.94	1.02	1.18	1.19
Urbanized	1.17	1.39	1.38	1.54	1.68	1.71
Less Urbanized	1.18	1.18	1.54	1.39	1.76	1.79
Completely Rural	2.37	2.99	3.16	3.52	4.36	3.63

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.69P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.0042		0.0089		0.2848	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0425		0.0706		0.3382	
New England	0.3198		0.4849		0.4186	
Middle Atlantic	0.0791		0.0928		0.5437	
Midwest	0.1096		0.0613		0.2256	
East North Central	0.0194		0.0441		0.3862	
West North Central	0.5007		0.7782		0.3832	
South	0.6411		0.9285		0.3038	
South Atlantic	0.9750		0.5229		0.2420	
East South Central	0.8040		0.5734		0.2905	
West South Central	0.2658		0.1274		0.7075	
West	0.0421		0.0409		0.1305	
Mountain	0.8668		0.6912		0.6909	
Pacific	0.0220		0.0298		0.1157	
<b>COUNTY TYPE</b>						
Large Metro	0.1045		0.0061		0.5469	
Small Metro	0.0307		0.2610		0.3394	
250K – 1 Mil. Pop.	0.1144		0.7266		0.5603	
< 250K Pop.	0.1626		0.1754		0.4260	
Nonmetro	0.1162		0.7797		0.8478	
Urbanized	0.0436		0.5434		0.0397	
Less Urbanized	0.9587		0.3624		0.1277	
Completely Rural	0.3572		0.6730		0.8045	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.70C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	680	641	925	954	946	978
<b>GEOGRAPHIC DIVISION</b>						
Northeast	259	260	354	371	394	404
New England	124	97	165	143	196	203
Middle Atlantic	225	239	310	339	334	346
Midwest	233	213	333	340	357	367
East North Central	204	175	275	292	293	310
West North Central	122	122	207	184	231	199
South	455	415	620	566	621	574
South Atlantic	332	306	464	453	466	444
East South Central	210	162	273	191	256	219
West South Central	219	215	316	293	344	313
West	385	364	522	545	516	570
Mountain	169	146	245	221	228	243
Pacific	348	334	457	494	466	511
<b>COUNTY TYPE</b>						
Large Metro	1,725	1,692	1,518	1,459	1,317	1,300
Small Metro	1,319	1,270	1,112	1,088	932	938
250K – 1 Mil. Pop.	1,083	1,065	924	898	771	772
< 250K Pop.	769	783	634	661	540	566
Nonmetro	887	830	736	654	631	546
Urbanized	681	579	581	451	501	379
Less Urbanized	649	669	511	512	422	433
Completely Rural	309	286	230	195	188	155

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.70D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.35	0.32	0.47	0.48	0.48	0.49
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.71	0.71	0.98	1.02	1.09	1.11
New England	1.30	1.01	1.73	1.49	2.06	2.11
Middle Atlantic	0.84	0.89	1.16	1.26	1.25	1.29
Midwest	0.55	0.50	0.78	0.79	0.84	0.86
East North Central	0.69	0.59	0.93	0.98	0.99	1.04
West North Central	0.95	0.93	1.61	1.41	1.80	1.52
South	0.63	0.57	0.87	0.77	0.87	0.78
South Atlantic	0.87	0.78	1.21	1.16	1.22	1.13
East South Central	1.80	1.37	2.34	1.62	2.20	1.85
West South Central	1.01	0.96	1.45	1.31	1.58	1.40
West	0.86	0.80	1.16	1.20	1.15	1.25
Mountain	1.23	1.05	1.78	1.60	1.66	1.76
Pacific	1.12	1.05	1.47	1.56	1.50	1.61
<b>COUNTY TYPE</b>						
Large Metro	0.50	0.47	0.67	0.66	0.70	0.72
Small Metro	0.61	0.57	0.82	0.82	0.87	0.82
250K – 1 Mil. Pop.	0.67	0.72	0.95	1.05	1.03	1.06
< 250K Pop.	1.26	0.87	1.62	1.27	1.66	1.31
Nonmetro	0.82	0.74	1.16	1.06	1.16	1.08
Urbanized	1.17	1.09	1.92	1.49	1.87	1.71
Less Urbanized	1.25	1.09	1.66	1.59	1.58	1.60
Completely Rural	2.75	2.20	3.73	2.74	3.54	2.97

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.70P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.9655		0.8274		0.7679	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4806		0.4085		0.4983	
New England	0.6627		0.3934		0.6411	
Middle Atlantic	0.3277		0.6082		0.6047	
Midwest	0.7222		0.3790		0.4960	
East North Central	0.5554		0.5309		0.4033	
West North Central	0.8045		0.5458		0.9674	
South	0.5679		0.4385		0.2062	
South Atlantic	0.2739		0.8194		0.7792	
East South Central	0.8539		0.2459		0.7061	
West South Central	0.6333		0.7965		0.1008	
West	0.8103		0.6350		0.9020	
Mountain	0.3128		0.3061		0.6121	
Pacific	0.4584		0.3154		0.9056	
<b>COUNTY TYPE</b>						
Large Metro	0.2475		0.8348		0.6708	
Small Metro	0.1260		0.1845		0.2378	
250K – 1 Mil. Pop.	0.5935		0.8480		0.9344	
< 250K Pop.	0.0557		0.0089		0.0279	
Nonmetro	0.7928		0.3645		0.0525	
Urbanized	0.3658		0.8395		0.0624	
Less Urbanized	0.2510		0.4037		0.4925	
Completely Rural	0.8909		0.7347		0.5798	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.71C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	694	656	953	977	974	1,006
<b>GEOGRAPHIC DIVISION</b>						
Northeast	263	266	359	380	400	415
New England	126	101	168	147	200	204
Middle Atlantic	228	245	315	348	338	359
Midwest	242	222	339	351	371	380
East North Central	210	181	281	303	304	321
West North Central	126	128	209	187	237	205
South	461	425	647	583	648	597
South Atlantic	338	313	487	467	489	464
East South Central	212	166	280	195	263	224
West South Central	224	221	324	300	352	320
West	393	375	536	559	538	588
Mountain	168	153	252	233	239	260
Pacific	356	341	470	504	484	522
<b>COUNTY TYPE</b>						
Large Metro	1,815	1,795	1,606	1,562	1,381	1,380
Small Metro	1,424	1,385	1,209	1,197	1,011	1,030
250K – 1 Mil. Pop.	1,156	1,122	985	951	822	813
< 250K Pop.	836	875	701	751	595	646
Nonmetro	951	894	791	706	671	588
Urbanized	736	625	631	488	542	405
Less Urbanized	702	723	558	562	453	474
Completely Rural	329	318	244	223	200	180

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.71D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.30	0.28	0.42	0.42	0.42	0.43
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.62	0.62	0.85	0.89	0.95	0.97
New England	1.13	0.90	1.50	1.31	1.79	1.82
Middle Atlantic	0.73	0.78	1.01	1.11	1.09	1.14
Midwest	0.48	0.44	0.68	0.70	0.74	0.76
East North Central	0.60	0.52	0.81	0.87	0.87	0.92
West North Central	0.83	0.83	1.37	1.22	1.56	1.33
South	0.55	0.49	0.77	0.68	0.77	0.69
South Atlantic	0.76	0.69	1.09	1.03	1.10	1.02
East South Central	1.56	1.20	2.05	1.41	1.93	1.62
West South Central	0.87	0.83	1.25	1.13	1.36	1.20
West	0.74	0.70	1.01	1.04	1.02	1.09
Mountain	1.03	0.94	1.55	1.43	1.47	1.59
Pacific	0.97	0.91	1.28	1.34	1.32	1.39
<b>COUNTY TYPE</b>						
Large Metro	0.44	0.40	0.58	0.58	0.61	0.63
Small Metro	0.52	0.50	0.71	0.71	0.76	0.73
250K – 1 Mil. Pop.	0.57	0.64	0.82	0.92	0.90	0.93
< 250K Pop.	1.06	0.76	1.37	1.10	1.41	1.17
Nonmetro	0.73	0.66	1.04	0.94	1.03	0.97
Urbanized	1.04	0.96	1.71	1.30	1.68	1.51
Less Urbanized	1.10	0.97	1.47	1.43	1.39	1.46
Completely Rural	2.51	2.00	3.39	2.50	3.25	2.73

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.71P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2010)</b>	<b>Lifetime (2011)</b>	<b>Past Year (2010)</b>	<b>Past Year (2011)</b>	<b>Past Month (2010)</b>	<b>Past Month (2011)</b>
<b>TOTAL</b>	0.6405		0.8354		0.9236	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.2990		0.5944		0.4074	
New England	0.8073		0.4822		0.5435	
Middle Atlantic	0.2072		0.7934		0.5452	
Midwest	0.9498		0.2327		0.3688	
East North Central	0.8923		0.3396		0.3320	
West North Central	0.9281		0.4975		0.8882	
South	0.6309		0.4688		0.1613	
South Atlantic	0.2790		0.7504		0.6377	
East South Central	0.8880		0.2286		0.5962	
West South Central	0.4935		0.9961		0.1285	
West	0.5426		0.4311		0.8946	
Mountain	0.3392		0.3620		0.6878	
Pacific	0.2589		0.1915		0.7185	
<b>COUNTY TYPE</b>						
Large Metro	0.1499		0.5279		0.7502	
Small Metro	0.2411		0.2881		0.3529	
250K – 1 Mil. Pop.	0.7817		0.7651		0.8269	
< 250K Pop.	0.0960		0.0211		0.0531	
Nonmetro	0.9879		0.4001		0.0593	
Urbanized	0.2163		0.8402		0.0427	
Less Urbanized	0.2478		0.4423		0.6380	
Completely Rural	0.9863		0.7442		0.5967	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.72C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	990	1,006	752	734	422	423
<b>GEOGRAPHIC DIVISION</b>						
Northeast	403	420	301	332	172	167
New England	204	204	148	174	82	91
Middle Atlantic	340	365	264	282	152	141
Midwest	380	381	334	294	176	174
East North Central	311	322	281	242	143	137
West North Central	239	208	182	159	104	106
South	658	603	452	473	264	268
South Atlantic	495	466	356	364	213	215
East South Central	266	226	152	160	88	91
West South Central	353	322	238	252	127	130
West	549	588	425	379	243	199
Mountain	246	263	206	183	122	88
Pacific	492	521	373	330	209	178
<b>COUNTY TYPE</b>						
Large Metro	1,389	1,393	738	718	349	334
Small Metro	1,023	1,040	533	553	294	259
250K – 1 Mil. Pop.	831	820	412	425	233	193
< 250K Pop.	604	654	335	363	178	166
Nonmetro	685	595	405	329	165	151
Urbanized	554	412	296	174	107	103
Less Urbanized	461	481	258	275	119	100
Completely Rural	203	184	104	83	60	53

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.72D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.39	0.39	0.30	0.29	0.17	0.16
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.87	0.89	0.65	0.71	0.37	0.36
New England	1.67	1.65	1.21	1.42	0.67	0.74
Middle Atlantic	0.99	1.06	0.77	0.81	0.44	0.41
Midwest	0.69	0.68	0.60	0.53	0.32	0.31
East North Central	0.81	0.83	0.73	0.63	0.37	0.35
West North Central	1.42	1.23	1.08	0.94	0.62	0.62
South	0.71	0.63	0.49	0.50	0.28	0.28
South Atlantic	1.01	0.93	0.73	0.73	0.43	0.43
East South Central	1.77	1.48	1.01	1.05	0.59	0.60
West South Central	1.22	1.08	0.83	0.85	0.44	0.44
West	0.93	0.98	0.72	0.63	0.41	0.33
Mountain	1.37	1.45	1.14	1.01	0.68	0.48
Pacific	1.21	1.25	0.91	0.79	0.51	0.43
<b>COUNTY TYPE</b>						
Large Metro	0.56	0.56	0.43	0.40	0.24	0.22
Small Metro	0.70	0.67	0.53	0.55	0.34	0.31
250K – 1 Mil. Pop.	0.83	0.86	0.63	0.67	0.41	0.35
< 250K Pop.	1.30	1.09	0.94	1.00	0.62	0.57
Nonmetro	0.95	0.89	0.75	0.63	0.36	0.34
Urbanized	1.54	1.37	1.15	0.79	0.54	0.53
Less Urbanized	1.28	1.34	0.97	1.04	0.52	0.47
Completely Rural	2.89	2.56	1.77	1.52	1.26	1.08

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.72P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.9707		0.2901		0.0256	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4730		0.9125		0.4903	
New England	0.5126		0.2263		0.5794	
Middle Atlantic	0.6570		0.5471		0.6437	
Midwest	0.3587		0.4472		0.5733	
East North Central	0.3701		0.3934		0.4836	
West North Central	0.7764		0.9235		0.9840	
South	0.2190		0.4600		0.4464	
South Atlantic	0.7278		0.8610		0.5398	
East South Central	0.6113		0.9867		0.7374	
West South Central	0.1583		0.1003		0.7502	
West	0.8024		0.0319		0.0215	
Mountain	0.7985		0.0688		0.1913	
Pacific	0.6707		0.1333		0.0533	
<b>COUNTY TYPE</b>						
Large Metro	0.8373		0.4958		0.1500	
Small Metro	0.3730		0.6185		0.3341	
250K – 1 Mil. Pop.	0.7678		0.5435		0.4472	
< 250K Pop.	0.0493		0.9524		0.5630	
Nonmetro	0.0530		0.0289		0.0977	
Urbanized	0.0265		0.0066		0.3466	
Less Urbanized	0.7051		0.6595		0.2092	
Completely Rural	0.5383		0.4820		0.5283	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.73C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	80	77	62	55	28	25
<b>GEOGRAPHIC DIVISION</b>						
Northeast	31	39	24	27	13	12
New England	14	14	12	10	5	4
Middle Atlantic	28	36	21	25	12	11
Midwest	30	32	23	22	9	10
East North Central	22	26	17	17	7	8
West North Central	21	19	16	13	6	6
South	53	44	38	31	14	15
South Atlantic	41	30	27	21	10	12
East South Central	16	14	13	11	4	5
West South Central	28	28	20	20	9	7
West	45	41	35	32	18	13
Mountain	19	18	18	12	9	6
Pacific	40	36	30	29	16	12
<b>COUNTY TYPE</b>						
Large Metro	73	69	54	46	23	20
Small Metro	46	47	32	33	13	13
250K – 1 Mil. Pop.	36	39	25	28	11	11
< 250K Pop.	28	26	19	19	6	7
Nonmetro	36	29	24	20	11	9
Urbanized	23	21	15	13	7	7
Less Urbanized	27	21	18	16	7	5
Completely Rural	9	10	8	4	2	2

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.73D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.33	0.31	0.25	0.22	0.11	0.10
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.73	0.90	0.58	0.62	0.30	0.28
New England	1.26	1.25	1.11	0.91	0.45	0.38
Middle Atlantic	0.88	1.14	0.68	0.78	0.38	0.36
Midwest	0.56	0.58	0.42	0.40	0.18	0.18
East North Central	0.58	0.67	0.44	0.46	0.19	0.21
West North Central	1.32	1.15	1.00	0.80	0.40	0.39
South	0.59	0.47	0.42	0.34	0.16	0.16
South Atlantic	0.91	0.65	0.61	0.45	0.23	0.26
East South Central	1.08	0.97	0.87	0.76	0.30	0.33
West South Central	0.94	0.88	0.67	0.65	0.30	0.23
West	0.77	0.68	0.60	0.53	0.31	0.22
Mountain	1.09	0.96	0.99	0.66	0.50	0.35
Pacific	1.00	0.88	0.74	0.70	0.39	0.28
<b>COUNTY TYPE</b>						
Large Metro	0.50	0.46	0.39	0.33	0.18	0.15
Small Metro	0.53	0.52	0.39	0.38	0.17	0.16
250K – 1 Mil. Pop.	0.62	0.63	0.45	0.46	0.22	0.20
< 250K Pop.	0.98	0.92	0.66	0.66	0.24	0.27
Nonmetro	0.79	0.68	0.56	0.51	0.26	0.24
Urbanized	0.96	1.00	0.74	0.72	0.43	0.42
Less Urbanized	1.28	1.03	0.84	0.81	0.36	0.31
Completely Rural	2.60	2.30	2.29	1.22	0.75	0.59

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.73P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.4804		0.1690		0.2621	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.2107		0.2993		0.8318	
New England	0.6836		0.4065		0.9864	
Middle Atlantic	0.1202		0.1363		0.8097	
Midwest	0.7072		0.4808		0.8441	
East North Central	0.1657		0.7185		0.6945	
West North Central	0.1921		0.1005		0.4666	
South	0.1832		0.1635		0.8203	
South Atlantic	0.1914		0.0564		0.8875	
East South Central	0.9658		0.8373		0.9049	
West South Central	0.5437		0.8314		0.5407	
West	0.3129		0.2077		0.0845	
Mountain	0.2085		0.0226		0.8043	
Pacific	0.6017		0.8391		0.0596	
<b>COUNTY TYPE</b>						
Large Metro	0.6498		0.2864		0.2937	
Small Metro	0.7136		0.8442		0.8891	
250K – 1 Mil. Pop.	0.4865		0.3625		0.2531	
< 250K Pop.	0.7556		0.3118		0.0955	
Nonmetro	0.6854		0.2573		0.5485	
Urbanized	0.7063		0.7037		0.6217	
Less Urbanized	0.7090		0.8507		0.2733	
Completely Rural	0.3384		0.0080		0.4580	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.74C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	169	184	168	189	127	125
<b>GEOGRAPHIC DIVISION</b>						
Northeast	68	74	67	71	47	46
New England	36	34	33	35	26	21
Middle Atlantic	58	66	57	62	39	41
Midwest	64	71	74	71	49	51
East North Central	50	59	61	60	40	41
West North Central	41	40	40	38	28	29
South	111	121	114	131	90	91
South Atlantic	86	93	92	110	79	79
East South Central	40	36	40	33	25	20
West South Central	55	60	57	58	37	37
West	93	95	87	91	64	52
Mountain	45	52	41	41	24	21
Pacific	80	81	76	82	59	47
<b>COUNTY TYPE</b>						
Large Metro	275	300	195	212	104	90
Small Metro	243	279	185	213	107	114
250K – 1 Mil. Pop.	179	170	128	113	66	53
< 250K Pop.	164	222	132	177	86	99
Nonmetro	121	118	94	95	48	40
Urbanized	92	84	72	70	34	27
Less Urbanized	85	82	67	66	35	27
Completely Rural	28	39	20	24	11	13

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.74D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.50	0.54	0.49	0.55	0.37	0.36
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1.13	1.20	1.10	1.15	0.78	0.75
New England	2.22	2.09	2.06	2.18	1.61	1.33
Middle Atlantic	1.31	1.45	1.30	1.37	0.88	0.90
Midwest	0.85	0.97	0.98	0.96	0.65	0.69
East North Central	0.98	1.16	1.19	1.17	0.78	0.81
West North Central	1.72	1.78	1.68	1.69	1.20	1.29
South	0.89	0.96	0.91	1.04	0.73	0.72
South Atlantic	1.36	1.45	1.46	1.71	1.26	1.22
East South Central	2.04	1.81	2.02	1.62	1.28	0.99
West South Central	1.33	1.45	1.38	1.39	0.90	0.89
West	1.14	1.15	1.08	1.11	0.79	0.63
Mountain	1.80	2.12	1.66	1.66	0.98	0.87
Pacific	1.42	1.40	1.36	1.43	1.05	0.82
<b>COUNTY TYPE</b>						
Large Metro	0.72	0.71	0.67	0.71	0.49	0.43
Small Metro	0.87	0.99	0.96	1.06	0.77	0.76
250K – 1 Mil. Pop.	1.03	1.07	1.05	1.05	0.77	0.64
< 250K Pop.	1.60	1.99	1.90	2.16	1.68	1.68
Nonmetro	1.18	1.19	1.15	1.19	0.76	0.72
Urbanized	1.68	1.71	1.77	1.84	1.14	1.01
Less Urbanized	1.76	1.79	1.68	1.79	1.12	1.09
Completely Rural	4.36	3.63	3.60	2.78	3.07	2.34

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.74P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.2848		0.3236		0.0037	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.3382		0.2168		0.0009	
New England	0.4186		0.2257		0.0280	
Middle Atlantic	0.5437		0.4673		0.0099	
Midwest	0.2256		0.7576		0.0611	
East North Central	0.3862		0.8612		0.0859	
West North Central	0.3832		0.3946		0.4118	
South	0.3038		0.5305		0.2756	
South Atlantic	0.2420		0.6032		0.3936	
East South Central	0.2905		0.3345		0.9068	
West South Central	0.7075		0.8250		0.3621	
West	0.1305		0.1411		0.9050	
Mountain	0.6909		0.3460		0.3524	
Pacific	0.1157		0.2371		0.6021	
<b>COUNTY TYPE</b>						
Large Metro	0.5469		0.3458		0.0024	
Small Metro	0.3394		0.8061		0.6273	
250K – 1 Mil. Pop.	0.5603		0.8743		0.4802	
< 250K Pop.	0.4260		0.8516		0.9405	
Nonmetro	0.8478		0.6326		0.1239	
Urbanized	0.0397		0.0503		0.0454	
Less Urbanized	0.1277		0.3663		0.5311	
Completely Rural	0.8045		0.6657		0.6822	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.75C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	946	978	715	699	398	396
<b>GEOGRAPHIC DIVISION</b>						
Northeast	394	404	294	313	161	160
New England	196	203	144	163	73	87
Middle Atlantic	334	346	258	265	144	135
Midwest	357	367	317	277	162	162
East North Central	293	310	268	227	133	130
West North Central	231	199	171	151	92	98
South	621	574	420	440	242	246
South Atlantic	466	444	325	324	191	191
East South Central	256	219	143	157	84	91
West South Central	344	313	228	247	123	123
West	516	570	399	370	229	194
Mountain	228	243	186	169	116	83
Pacific	466	511	354	327	198	174
<b>COUNTY TYPE</b>						
Large Metro	1,317	1,300	686	656	321	316
Small Metro	932	938	459	479	265	224
250K – 1 Mil. Pop.	771	772	367	397	218	190
< 250K Pop.	540	566	284	287	152	117
Nonmetro	631	546	372	296	156	141
Urbanized	501	379	258	154	98	95
Less Urbanized	422	433	235	244	110	93
Completely Rural	188	155	94	73	57	47

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.75D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.48	0.49	0.37	0.35	0.20	0.20
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1.09	1.11	0.81	0.86	0.44	0.44
New England	2.06	2.11	1.51	1.70	0.76	0.90
Middle Atlantic	1.25	1.29	0.96	0.99	0.54	0.50
Midwest	0.84	0.86	0.75	0.65	0.38	0.38
East North Central	0.99	1.04	0.90	0.76	0.45	0.44
West North Central	1.80	1.52	1.33	1.15	0.72	0.75
South	0.87	0.78	0.59	0.60	0.34	0.34
South Atlantic	1.22	1.13	0.85	0.83	0.50	0.49
East South Central	2.20	1.85	1.22	1.33	0.72	0.77
West South Central	1.58	1.40	1.05	1.10	0.57	0.55
West	1.15	1.25	0.89	0.81	0.51	0.42
Mountain	1.66	1.76	1.35	1.22	0.84	0.60
Pacific	1.50	1.61	1.14	1.03	0.64	0.55
<b>COUNTY TYPE</b>						
Large Metro	0.70	0.72	0.54	0.50	0.29	0.28
Small Metro	0.87	0.82	0.63	0.66	0.41	0.36
250K – 1 Mil. Pop.	1.03	1.06	0.76	0.82	0.50	0.44
< 250K Pop.	1.66	1.31	1.14	1.13	0.71	0.56
Nonmetro	1.16	1.08	0.91	0.76	0.45	0.41
Urbanized	1.87	1.71	1.39	0.97	0.67	0.65
Less Urbanized	1.58	1.60	1.20	1.23	0.65	0.56
Completely Rural	3.54	2.97	2.04	1.88	1.50	1.24

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.75P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.7679		0.5015		0.1783	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4983		0.7737		0.8359	
New England	0.6411		0.1056		0.9791	
Middle Atlantic	0.6047		0.5323		0.8225	
Midwest	0.4960		0.5884		0.9703	
East North Central	0.4033		0.3711		0.8369	
West North Central	0.9674		0.7281		0.7348	
South	0.2062		0.4333		0.7125	
South Atlantic	0.7792		0.8618		0.7511	
East South Central	0.7061		0.8135		0.7069	
West South Central	0.1008		0.0662		0.9848	
West	0.9020		0.0697		0.0246	
Mountain	0.6121		0.1282		0.1278	
Pacific	0.9056		0.1954		0.0771	
<b>COUNTY TYPE</b>						
Large Metro	0.6708		0.6879		0.5536	
Small Metro	0.2378		0.4378		0.4403	
250K – 1 Mil. Pop.	0.9344		0.3660		0.6544	
< 250K Pop.	0.0279		0.9408		0.4581	
Nonmetro	0.0525		0.0345		0.1979	
Urbanized	0.0624		0.0131		0.5737	
Less Urbanized	0.4925		0.5588		0.3051	
Completely Rural	0.5798		0.4931		0.4460	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.76C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	974	1,006	748	736	421	423
<b>GEOGRAPHIC DIVISION</b>						
Northeast	400	415	302	333	171	168
New England	200	204	147	173	82	91
Middle Atlantic	338	359	265	282	151	142
Midwest	371	380	332	291	175	173
East North Central	304	321	278	241	142	136
West North Central	237	205	181	157	103	106
South	648	597	453	473	265	268
South Atlantic	489	464	358	365	214	214
East South Central	263	224	150	161	88	92
West South Central	352	320	237	251	127	130
West	538	588	418	381	241	199
Mountain	239	260	204	185	122	88
Pacific	484	522	366	332	208	178
<b>COUNTY TYPE</b>						
Large Metro	1,381	1,380	733	713	347	334
Small Metro	1,011	1,030	527	548	294	258
250K – 1 Mil. Pop.	822	813	407	420	233	192
< 250K Pop.	595	646	330	359	178	165
Nonmetro	671	588	399	326	164	151
Urbanized	542	405	290	172	106	102
Less Urbanized	453	474	254	271	118	99
Completely Rural	200	180	102	83	59	53

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.76D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.42	0.43	0.33	0.32	0.18	0.18
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.95	0.97	0.71	0.78	0.40	0.39
New England	1.79	1.82	1.32	1.54	0.74	0.81
Middle Atlantic	1.09	1.14	0.85	0.90	0.49	0.45
Midwest	0.74	0.76	0.66	0.58	0.35	0.34
East North Central	0.87	0.92	0.80	0.69	0.41	0.39
West North Central	1.56	1.33	1.19	1.02	0.68	0.69
South	0.77	0.69	0.54	0.55	0.31	0.31
South Atlantic	1.10	1.02	0.80	0.80	0.48	0.47
East South Central	1.93	1.62	1.10	1.17	0.64	0.67
West South Central	1.36	1.20	0.92	0.95	0.49	0.49
West	1.02	1.09	0.79	0.71	0.46	0.37
Mountain	1.47	1.59	1.25	1.13	0.75	0.54
Pacific	1.32	1.39	1.00	0.89	0.57	0.47
<b>COUNTY TYPE</b>						
Large Metro	0.61	0.63	0.48	0.45	0.26	0.25
Small Metro	0.76	0.73	0.58	0.61	0.38	0.34
250K – 1 Mil. Pop.	0.90	0.93	0.70	0.73	0.45	0.39
< 250K Pop.	1.41	1.17	1.04	1.10	0.68	0.63
Nonmetro	1.03	0.97	0.82	0.69	0.40	0.37
Urbanized	1.68	1.51	1.27	0.89	0.59	0.59
Less Urbanized	1.39	1.46	1.07	1.14	0.58	0.51
Completely Rural	3.25	2.73	1.90	1.68	1.36	1.19

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 2.76P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Geographic Characteristic</b>	<b>Alcohol Use (2010)</b>	<b>Alcohol Use (2011)</b>	<b>Binge Alcohol Use (2010)</b>	<b>Binge Alcohol Use (2011)</b>	<b>Heavy Alcohol Use (2010)</b>	<b>Heavy Alcohol Use (2011)</b>
<b>TOTAL</b>	0.9236		0.3614		0.0322	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4074		0.9777		0.4817	
New England	0.5435		0.2025		0.5824	
Middle Atlantic	0.5452		0.4644		0.6302	
Midwest	0.3688		0.4950		0.5915	
East North Central	0.3320		0.3880		0.4758	
West North Central	0.8882		0.9513		0.9707	
South	0.1613		0.3829		0.4617	
South Atlantic	0.6377		0.9722		0.5378	
East South Central	0.5962		0.9949		0.7350	
West South Central	0.1285		0.0870		0.7916	
West	0.8946		0.0418		0.0289	
Mountain	0.6878		0.1049		0.2059	
Pacific	0.7185		0.1397		0.0685	
<b>COUNTY TYPE</b>						
Large Metro	0.7502		0.5772		0.1746	
Small Metro	0.3529		0.6073		0.3386	
250K – 1 Mil. Pop.	0.8269		0.4901		0.4917	
< 250K Pop.	0.0531		0.9819		0.5088	
Nonmetro	0.0593		0.0353		0.1082	
Urbanized	0.0427		0.0100		0.3660	
Less Urbanized	0.6380		0.6197		0.2210	
Completely Rural	0.5967		0.6395		0.5460	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.