

Table 2.77C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	291	332	266	307	204	239
GENDER						
Male	197	188	180	173	141	131
Female	170	213	156	199	118	159
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	251	278	230	262	182	207
White	219	239	203	224	160	180
Black or African American	97	88	82	77	60	56
American Indian or Alaska Native	14	12	11	12	9	7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	49	56	44	50	31	38
Two or More Races	30	42	26	40	19	30
Hispanic or Latino	138	148	125	129	85	90
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	145	128	135	119	108	96
Female, White, Not Hispanic	131	161	122	154	97	125
Male, Black, Not Hispanic	65	58	55	51	43	37
Female, Black, Not Hispanic	59	53	52	48	35	36
Male, Hispanic	93	99	85	87	63	65
Female, Hispanic	76	85	68	78	47	54

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.77D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.44	0.49	0.44	0.50	0.41	0.47
GENDER						
Male	0.60	0.61	0.60	0.60	0.55	0.54
Female	0.59	0.67	0.58	0.68	0.53	0.65
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.48	0.51	0.48	0.53	0.47	0.51
White	0.54	0.59	0.55	0.61	0.54	0.61
Black or African American	1.01	0.98	0.96	0.97	0.89	0.83
American Indian or Alaska Native	3.58	4.22	3.52	4.24	3.65	2.74
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.35	2.33	2.25	2.16	1.82	1.79
Two or More Races	2.34	2.31	2.35	2.39	2.05	2.25
Hispanic or Latino	1.01	1.01	1.00	1.01	0.84	0.84
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.75	0.71	0.75	0.71	0.71	0.67
Female, White, Not Hispanic	0.74	0.84	0.74	0.87	0.72	0.87
Male, Black, Not Hispanic	1.47	1.36	1.42	1.32	1.30	1.13
Female, Black, Not Hispanic	1.35	1.43	1.32	1.42	1.08	1.14
Male, Hispanic	1.33	1.37	1.37	1.37	1.22	1.25
Female, Hispanic	1.34	1.42	1.33	1.38	1.04	1.11

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.77P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0010		0.0033		0.0628	
GENDER						
Male	0.0011		0.0002		0.0005	
Female	0.1319		0.5442		0.4911	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0180		0.0529		0.1980	
White	0.0888		0.1262		0.2483	
Black or African American	0.0357		0.0867		0.0615	
American Indian or Alaska Native	0.1324		0.9410		0.5369	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7911		0.8312		0.2034	
Two or More Races	0.7250		0.6568		0.2977	
Hispanic or Latino	0.0090		0.0093		0.1248	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.0413		0.0293		0.0273	
Female, White, Not Hispanic	0.6411		0.8730		0.6945	
Male, Black, Not Hispanic	0.1420		0.2415		0.1903	
Female, Black, Not Hispanic	0.1211		0.2061		0.1585	
Male, Hispanic	0.0195		0.0020		0.0158	
Female, Hispanic	0.1635		0.5841		0.5622	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.78C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	2,217	2,134	1,960	1,885	1,681	1,656
GENDER						
Male	1,420	1,332	1,244	1,169	1,136	1,074
Female	1,276	1,334	1,115	1,197	909	1,018
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,111	2,016	1,868	1,778	1,607	1,557
White	1,912	1,841	1,703	1,638	1,468	1,446
Black or African American	708	670	582	550	492	446
American Indian or Alaska Native	*	105	57	94	50	83
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	421	497	378	421	328	364
Two or More Races	174	184	146	146	117	130
Hispanic or Latino	707	724	602	620	485	537
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,182	1,124	1,041	997	957	910
Female, White, Not Hispanic	1,126	1,126	1,000	1,004	818	889
Male, Black, Not Hispanic	423	420	358	348	325	309
Female, Black, Not Hispanic	475	430	385	360	318	279
Male, Hispanic	510	518	454	459	381	413
Female, Hispanic	407	425	319	367	256	298

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.78D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.32	0.29	0.44	0.44	0.45	0.45
GENDER						
Male	0.35	0.36	0.59	0.59	0.63	0.64
Female	0.48	0.45	0.58	0.59	0.56	0.59
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.34	0.29	0.46	0.47	0.47	0.48
White	0.32	0.28	0.50	0.50	0.52	0.53
Black or African American	1.08	1.01	1.37	1.22	1.34	1.17
American Indian or Alaska Native	*	4.44	5.05	4.64	4.51	4.48
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.47	2.11	2.68	2.44	2.50	2.39
Two or More Races	2.78	2.51	3.65	3.44	3.74	3.42
Hispanic or Latino	0.96	1.00	1.28	1.12	1.22	1.19
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.35	0.33	0.68	0.67	0.73	0.74
Female, White, Not Hispanic	0.49	0.44	0.66	0.64	0.67	0.70
Male, Black, Not Hispanic	1.39	1.20	1.87	1.72	1.94	1.73
Female, Black, Not Hispanic	1.39	1.44	1.81	1.70	1.74	1.54
Male, Hispanic	1.10	1.24	1.75	1.56	1.76	1.77
Female, Hispanic	1.47	1.54	1.65	1.67	1.52	1.55

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.78P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.9668		0.9398		0.8003	
GENDER						
Male	0.4461		0.3788		0.7576	
Female	0.5818		0.4832		0.4484	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2675		0.9250		0.9694	
White	0.1867		0.9767		0.9453	
Black or African American	0.6997		0.9239		0.7301	
American Indian or Alaska Native	*		0.2581		0.1469	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1925		0.4349		0.6111	
Two or More Races	0.3881		0.7868		0.7258	
Hispanic or Latino	0.0518		0.8262		0.3589	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.2891		0.4042		0.2891	
Female, White, Not Hispanic	0.3577		0.4322		0.2344	
Male, Black, Not Hispanic	0.2527		0.7280		0.9193	
Female, Black, Not Hispanic	0.1744		0.6783		0.5667	
Male, Hispanic	0.4083		0.5592		0.2419	
Female, Hispanic	0.1012		0.9422		0.7895	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.79C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	204	239	161	181	84	95
GENDER						
Male	141	131	118	107	66	59
Female	118	159	91	117	45	60
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	182	207	146	157	79	83
White	160	180	131	141	75	80
Black or African American	60	56	42	36	16	15
American Indian or Alaska Native	9	7	8	6	6	3
Native Hawaiian or Other Pacific Islander	*	*	*	*	1	*
Asian	31	38	20	28	7	11
Two or More Races	19	30	15	22	10	13
Hispanic or Latino	85	90	68	70	30	29
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	108	96	93	80	57	50
Female, White, Not Hispanic	97	125	76	94	42	48
Male, Black, Not Hispanic	43	37	31	28	14	11
Female, Black, Not Hispanic	35	36	23	23	6	10
Male, Hispanic	63	65	54	52	27	24
Female, Hispanic	47	54	37	40	13	18

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.79D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.41	0.47	0.35	0.39	0.21	0.23
GENDER						
Male	0.55	0.54	0.50	0.47	0.32	0.29
Female	0.53	0.65	0.44	0.53	0.23	0.30
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.51	0.41	0.43	0.25	0.26
White	0.54	0.61	0.49	0.53	0.32	0.34
Black or African American	0.89	0.83	0.67	0.59	0.27	0.27
American Indian or Alaska Native	3.65	2.74	3.46	2.36	2.74	1.30
Native Hawaiian or Other Pacific Islander	*	*	*	*	1.39	*
Asian	1.82	1.79	1.25	1.44	0.42	0.61
Two or More Races	2.05	2.25	1.72	1.85	1.30	1.18
Hispanic or Latino	0.84	0.84	0.72	0.72	0.38	0.34
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.71	0.67	0.67	0.62	0.46	0.43
Female, White, Not Hispanic	0.72	0.87	0.61	0.74	0.37	0.42
Male, Black, Not Hispanic	1.30	1.13	0.99	0.90	0.49	0.40
Female, Black, Not Hispanic	1.08	1.14	0.79	0.78	0.23	0.36
Male, Hispanic	1.22	1.25	1.12	1.08	0.66	0.55
Female, Hispanic	1.04	1.11	0.88	0.89	0.34	0.44

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.79P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.0628		0.0229		0.0153	
GENDER						
Male	0.0005		0.0008		0.0068	
Female	0.4911		0.9002		0.5358	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1980		0.1183		0.0486	
White	0.2483		0.0943		0.0524	
Black or African American	0.0615		0.7159		0.7222	
American Indian or Alaska Native	0.5369		0.2845		0.4765	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2034		0.5116		0.5293	
Two or More Races	0.2977		0.8013		0.9803	
Hispanic or Latino	0.1248		0.0453		0.1129	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.0273		0.0535		0.0697	
Female, White, Not Hispanic	0.6945		0.6027		0.3531	
Male, Black, Not Hispanic	0.1903		0.4926		0.6224	
Female, Black, Not Hispanic	0.1585		0.8025		0.9196	
Male, Hispanic	0.0158		0.0032		0.0837	
Female, Hispanic	0.5622		0.4610		0.9358	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.80C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Demographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	1,681	1,656	910	858	446	428
GENDER						
Male	1,136	1,074	709	665	375	362
Female	909	1,018	471	471	203	197
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,607	1,557	845	794	422	398
White	1,468	1,446	755	732	376	371
Black or African American	492	446	285	272	138	112
American Indian or Alaska Native	50	83	40	57	17	43
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	328	364	147	136	72	38
Two or More Races	117	130	62	57	31	37
Hispanic or Latino	485	537	370	364	137	158
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	957	910	589	573	319	311
Female, White, Not Hispanic	818	889	384	386	163	174
Male, Black, Not Hispanic	325	309	210	201	113	88
Female, Black, Not Hispanic	318	279	172	167	77	69
Male, Hispanic	381	413	311	311	117	148
Female, Hispanic	256	298	174	175	70	50

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.80D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Demographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.45	0.45	0.34	0.33	0.19	0.19
GENDER						
Male	0.63	0.64	0.54	0.53	0.33	0.33
Female	0.56	0.59	0.39	0.37	0.18	0.17
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.48	0.36	0.35	0.21	0.20
White	0.52	0.53	0.40	0.41	0.24	0.23
Black or African American	1.34	1.17	1.00	0.91	0.54	0.43
American Indian or Alaska Native	4.51	4.48	3.72	4.11	1.74	3.18
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.50	2.39	1.35	1.25	0.69	0.36
Two or More Races	3.74	3.42	2.61	2.06	1.29	1.24
Hispanic or Latino	1.22	1.19	1.05	0.96	0.46	0.48
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.73	0.74	0.66	0.65	0.42	0.41
Female, White, Not Hispanic	0.67	0.70	0.46	0.46	0.21	0.22
Male, Black, Not Hispanic	1.94	1.73	1.70	1.51	1.00	0.75
Female, Black, Not Hispanic	1.74	1.54	1.14	1.07	0.56	0.49
Male, Hispanic	1.76	1.77	1.67	1.65	0.77	0.89
Female, Hispanic	1.52	1.55	1.13	1.04	0.49	0.32

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.80P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Demographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.8003		0.5068		0.0684	
GENDER						
Male	0.7576		0.3818		0.1357	
Female	0.4484		0.8698		0.3007	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9694		0.8036		0.0461	
White	0.9453		0.8747		0.0874	
Black or African American	0.7301		0.7314		0.3818	
American Indian or Alaska Native	0.1469		0.9925		0.1106	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6111		0.6229		0.2039	
Two or More Races	0.7258		0.2267		0.6233	
Hispanic or Latino	0.3589		0.2571		0.8545	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.2891		0.6907		0.0593	
Female, White, Not Hispanic	0.2344		0.4221		0.8365	
Male, Black, Not Hispanic	0.9193		0.6485		0.6452	
Female, Black, Not Hispanic	0.5667		0.9793		0.3908	
Male, Hispanic	0.2419		0.6078		0.2615	
Female, Hispanic	0.7895		0.3162		0.1004	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.81C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	291	332	266	307	204	239
GEOGRAPHIC DIVISION						
Northeast	112	137	106	130	87	113
New England	49	69	46	59	35	52
Middle Atlantic	100	118	94	116	79	101
Midwest	116	114	107	106	81	82
East North Central	98	95	90	89	65	67
West North Central	62	61	57	55	47	44
South	201	226	181	209	141	168
South Atlantic	156	193	142	185	104	153
East South Central	66	56	57	48	44	33
West South Central	107	108	97	93	84	66
West	146	169	134	150	97	102
Mountain	77	81	64	71	48	49
Pacific	124	149	117	133	85	90
COUNTY TYPE						
Large Metro	225	251	206	227	154	167
Small Metro	186	214	166	202	128	163
250K – 1 Mil. Pop.	137	139	121	126	91	90
< 250K Pop.	126	165	115	159	90	136
Nonmetro	122	97	111	90	85	66
Urbanized	79	69	73	63	53	44
Less Urbanized	99	71	90	65	69	51
Completely Rural	31	35	26	32	19	24

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.81D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.44	0.49	0.44	0.50	0.41	0.47
GEOGRAPHIC DIVISION						
Northeast	0.96	1.07	0.97	1.13	0.97	1.20
New England	1.77	1.94	1.89	1.70	1.72	1.86
Middle Atlantic	1.13	1.27	1.12	1.40	1.15	1.49
Midwest	0.78	0.78	0.80	0.77	0.75	0.75
East North Central	0.91	0.91	0.92	0.91	0.82	0.89
West North Central	1.49	1.52	1.58	1.48	1.57	1.42
South	0.77	0.86	0.77	0.90	0.77	0.89
South Atlantic	1.15	1.45	1.17	1.56	1.11	1.58
East South Central	1.66	1.64	1.62	1.49	1.57	1.21
West South Central	1.34	1.15	1.35	1.14	1.38	1.09
West	1.01	1.08	1.01	1.07	0.83	0.86
Mountain	1.41	1.52	1.39	1.52	1.33	1.35
Pacific	1.31	1.40	1.31	1.37	1.04	1.08
COUNTY TYPE						
Large Metro	0.64	0.68	0.64	0.68	0.59	0.62
Small Metro	0.76	0.89	0.76	0.93	0.75	0.94
250K – 1 Mil. Pop.	0.91	0.96	0.91	0.95	0.85	0.81
< 250K Pop.	1.32	1.84	1.36	1.97	1.45	2.21
Nonmetro	1.05	0.93	1.08	0.97	1.03	0.92
Urbanized	1.55	1.33	1.68	1.42	1.44	1.27
Less Urbanized	1.51	1.45	1.59	1.42	1.65	1.42
Completely Rural	3.88	2.71	3.37	2.73	2.75	2.83

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.81P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0010		0.0033		0.0628	
GEOGRAPHIC DIVISION						
Northeast	0.7456		0.5649		0.5730	
New England	0.6812		0.9682		0.6232	
Middle Atlantic	0.8754		0.5357		0.7025	
Midwest	0.0516		0.2454		0.3343	
East North Central	0.2362		0.6256		0.9221	
West North Central	0.0853		0.1678		0.0497	
South	0.0004		0.0021		0.0761	
South Atlantic	0.0059		0.0538		0.3576	
East South Central	0.0547		0.0872		0.1546	
West South Central	0.1811		0.0525		0.2361	
West	0.7025		0.5645		0.2376	
Mountain	0.2248		0.6029		0.1533	
Pacific	0.8974		0.6883		0.5419	
COUNTY TYPE						
Large Metro	0.1898		0.0509		0.2044	
Small Metro	0.0191		0.1882		0.4300	
250K – 1 Mil. Pop.	0.0120		0.0838		0.1552	
< 250K Pop.	0.5236		0.9739		0.8624	
Nonmetro	0.0034		0.0357		0.1936	
Urbanized	0.0890		0.1661		0.0938	
Less Urbanized	0.0059		0.0509		0.6167	
Completely Rural	0.9191		0.6215		0.9752	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.82C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	2,217	2,134	1,960	1,885	1,681	1,656
GEOGRAPHIC DIVISION						
Northeast	844	885	744	815	681	720
New England	452	396	419	364	415	348
Middle Atlantic	710	793	613	729	539	631
Midwest	849	800	748	715	671	627
East North Central	683	659	602	582	532	514
West North Central	486	475	439	430	397	373
South	1,462	1,393	1,307	1,151	1,107	1,018
South Atlantic	1,118	1,059	1,014	871	856	766
East South Central	479	449	403	371	342	319
West South Central	836	780	741	651	630	592
West	1,178	1,165	1,041	1,068	883	929
Mountain	603	657	518	616	417	515
Pacific	999	968	894	873	772	773
COUNTY TYPE						
Large Metro	1,786	1,758	1,578	1,528	1,368	1,358
Small Metro	1,381	1,334	1,174	1,148	985	992
250K – 1 Mil. Pop.	1,132	1,105	970	938	811	802
< 250K Pop.	803	828	669	701	571	603
Nonmetro	931	873	779	688	663	571
Urbanized	723	606	624	473	537	392
Less Urbanized	677	703	537	541	441	456
Completely Rural	323	306	240	212	196	171

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.82D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.32	0.29	0.44	0.44	0.45	0.45
GEOGRAPHIC DIVISION						
Northeast	0.66	0.66	0.90	0.93	1.00	1.02
New England	1.21	0.94	1.60	1.35	1.88	1.91
Middle Atlantic	0.77	0.82	1.07	1.16	1.15	1.19
Midwest	0.50	0.45	0.72	0.73	0.78	0.79
East North Central	0.63	0.54	0.85	0.91	0.91	0.96
West North Central	0.86	0.85	1.46	1.28	1.66	1.39
South	0.58	0.52	0.81	0.71	0.81	0.72
South Atlantic	0.80	0.72	1.14	1.07	1.15	1.06
East South Central	1.66	1.25	2.19	1.48	2.04	1.71
West South Central	0.91	0.87	1.32	1.20	1.43	1.28
West	0.79	0.73	1.07	1.09	1.07	1.14
Mountain	1.12	0.98	1.64	1.51	1.54	1.66
Pacific	1.03	0.95	1.35	1.41	1.39	1.46
COUNTY TYPE						
Large Metro	0.46	0.42	0.61	0.61	0.65	0.66
Small Metro	0.56	0.52	0.76	0.76	0.81	0.77
250K – 1 Mil. Pop.	0.62	0.67	0.88	0.98	0.95	0.98
< 250K Pop.	1.12	0.78	1.47	1.15	1.51	1.21
Nonmetro	0.77	0.69	1.10	0.98	1.09	1.01
Urbanized	1.09	1.01	1.82	1.37	1.78	1.57
Less Urbanized	1.17	1.01	1.55	1.49	1.47	1.51
Completely Rural	2.59	2.09	3.55	2.62	3.37	2.86

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.82P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.9668		0.9398		0.8003	
GEOGRAPHIC DIVISION						
Northeast	0.4295		0.4689		0.4569	
New England	0.7602		0.4869		0.5314	
Middle Atlantic	0.3135		0.6262		0.6266	
Midwest	0.8487		0.2784		0.4131	
East North Central	0.6656		0.4112		0.3587	
West North Central	0.7628		0.4955		0.9369	
South	0.4932		0.3772		0.1452	
South Atlantic	0.2641		0.7403		0.6628	
East South Central	0.9974		0.1816		0.5099	
West South Central	0.6681		0.8589		0.1114	
West	0.7358		0.5209		0.9949	
Mountain	0.3120		0.3444		0.6475	
Pacific	0.3897		0.2448		0.8070	
COUNTY TYPE						
Large Metro	0.2589		0.7430		0.6486	
Small Metro	0.1531		0.2544		0.2951	
250K – 1 Mil. Pop.	0.6387		0.7942		0.9150	
< 250K Pop.	0.0645		0.0162		0.0450	
Nonmetro	0.8776		0.4484		0.0589	
Urbanized	0.2769		0.8780		0.0589	
Less Urbanized	0.2335		0.5093		0.5802	
Completely Rural	0.9468		0.6908		0.5508	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.83C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	204	239	161	181	84	95
GEOGRAPHIC DIVISION						
Northeast	87	113	71	78	36	35
New England	35	52	29	32	16	14
Middle Atlantic	79	101	64	72	32	32
Midwest	81	82	62	63	33	28
East North Central	65	67	49	54	25	23
West North Central	47	44	38	32	21	15
South	141	168	113	132	59	77
South Atlantic	104	153	83	121	47	73
East South Central	44	33	39	24	23	11
West South Central	84	66	68	51	31	23
West	97	102	77	77	37	34
Mountain	48	49	37	31	15	17
Pacific	85	90	67	70	34	30
COUNTY TYPE						
Large Metro	154	167	118	112	58	51
Small Metro	128	163	101	136	54	78
250K – 1 Mil. Pop.	91	90	69	65	35	31
< 250K Pop.	90	136	74	119	41	71
Nonmetro	85	66	68	51	35	22
Urbanized	53	44	42	33	20	13
Less Urbanized	69	51	55	40	28	17
Completely Rural	19	24	16	13	6	6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.83D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.41	0.47	0.35	0.39	0.21	0.23
GEOGRAPHIC DIVISION						
Northeast	0.97	1.20	0.87	0.90	0.49	0.48
New England	1.72	1.86	1.53	1.27	0.87	0.73
Middle Atlantic	1.15	1.49	1.04	1.14	0.59	0.59
Midwest	0.75	0.75	0.64	0.62	0.37	0.31
East North Central	0.82	0.89	0.70	0.75	0.40	0.36
West North Central	1.57	1.42	1.34	1.09	0.78	0.58
South	0.77	0.89	0.67	0.78	0.39	0.51
South Atlantic	1.11	1.58	0.95	1.40	0.62	0.94
East South Central	1.57	1.21	1.47	1.00	0.96	0.49
West South Central	1.38	1.09	1.23	0.93	0.61	0.49
West	0.83	0.86	0.70	0.75	0.39	0.36
Mountain	1.33	1.35	1.12	1.02	0.51	0.59
Pacific	1.04	1.08	0.88	0.96	0.52	0.44
COUNTY TYPE						
Large Metro	0.59	0.62	0.49	0.47	0.28	0.24
Small Metro	0.75	0.94	0.66	0.87	0.41	0.56
250K – 1 Mil. Pop.	0.85	0.81	0.72	0.67	0.42	0.35
< 250K Pop.	1.45	2.21	1.33	2.12	0.90	1.47
Nonmetro	1.03	0.92	0.91	0.76	0.52	0.39
Urbanized	1.44	1.27	1.37	1.06	0.70	0.52
Less Urbanized	1.65	1.42	1.40	1.21	0.82	0.60
Completely Rural	2.75	2.83	2.53	1.86	1.20	1.06

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.83P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.0628		0.0229		0.0153	
GEOGRAPHIC DIVISION						
Northeast	0.5730		0.7677		0.3534	
New England	0.6232		0.5681		0.8105	
Middle Atlantic	0.7025		0.9413		0.3585	
Midwest	0.3343		0.3281		0.0854	
East North Central	0.9221		0.7684		0.4114	
West North Central	0.0497		0.0201		0.0745	
South	0.0761		0.1693		0.2104	
South Atlantic	0.3576		0.6659		0.9472	
East South Central	0.1546		0.0961		0.1133	
West South Central	0.2361		0.3140		0.0497	
West	0.2376		0.0740		0.1964	
Mountain	0.1533		0.0124		0.9676	
Pacific	0.5419		0.4127		0.1471	
COUNTY TYPE						
Large Metro	0.2044		0.0454		0.0202	
Small Metro	0.4300		0.5689		0.9044	
250K – 1 Mil. Pop.	0.1552		0.1479		0.2135	
< 250K Pop.	0.8624		0.7071		0.5093	
Nonmetro	0.1936		0.1500		0.0067	
Urbanized	0.0938		0.1023		0.0049	
Less Urbanized	0.6167		0.7462		0.1757	
Completely Rural	0.9752		0.3365		0.9369	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.84C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2010 and 2011

Geographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	1,681	1,656	910	858	446	428
GEOGRAPHIC DIVISION						
Northeast	681	720	350	360	174	167
New England	415	348	175	198	80	90
Middle Atlantic	539	631	303	302	155	141
Midwest	671	627	394	365	180	184
East North Central	532	514	331	298	145	146
West North Central	397	373	216	209	104	114
South	1,107	1,018	521	553	265	265
South Atlantic	856	766	381	372	205	201
East South Central	342	319	171	180	92	93
West South Central	630	592	305	359	139	138
West	883	929	520	463	265	207
Mountain	417	515	250	252	130	95
Pacific	772	773	454	388	230	183
COUNTY TYPE						
Large Metro	1,368	1,358	721	702	339	329
Small Metro	985	992	500	510	281	232
250K – 1 Mil. Pop.	811	802	397	414	227	190
< 250K Pop.	571	603	305	310	164	128
Nonmetro	663	571	392	314	159	147
Urbanized	537	392	285	163	103	99
Less Urbanized	441	456	247	258	113	95
Completely Rural	196	171	99	80	58	52

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.84D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2010 and 2011

Geographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.45	0.45	0.34	0.33	0.19	0.19
GEOGRAPHIC DIVISION						
Northeast	1.00	1.02	0.76	0.82	0.42	0.41
New England	1.88	1.91	1.37	1.61	0.74	0.85
Middle Atlantic	1.15	1.19	0.91	0.94	0.51	0.47
Midwest	0.78	0.79	0.70	0.60	0.37	0.36
East North Central	0.91	0.96	0.84	0.72	0.43	0.41
West North Central	1.66	1.39	1.25	1.06	0.71	0.71
South	0.81	0.72	0.56	0.56	0.32	0.31
South Atlantic	1.15	1.06	0.83	0.79	0.48	0.45
East South Central	2.04	1.71	1.14	1.24	0.67	0.70
West South Central	1.43	1.28	0.96	1.01	0.53	0.51
West	1.07	1.14	0.83	0.75	0.47	0.39
Mountain	1.54	1.66	1.31	1.21	0.79	0.56
Pacific	1.39	1.46	1.05	0.94	0.59	0.50
COUNTY TYPE						
Large Metro	0.65	0.66	0.50	0.47	0.27	0.26
Small Metro	0.81	0.77	0.60	0.62	0.39	0.33
250K – 1 Mil. Pop.	0.95	0.98	0.73	0.76	0.47	0.40
< 250K Pop.	1.51	1.21	1.06	1.06	0.69	0.54
Nonmetro	1.09	1.01	0.86	0.71	0.42	0.38
Urbanized	1.78	1.57	1.33	0.91	0.62	0.60
Less Urbanized	1.47	1.51	1.13	1.17	0.61	0.53
Completely Rural	3.37	2.86	1.95	1.79	1.42	1.25

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.84P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Geographic Characteristic	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.8003		0.5068		0.0684	
GEOGRAPHIC DIVISION						
Northeast	0.4569		0.8526		0.5961	
New England	0.5314		0.1799		0.5967	
Middle Atlantic	0.6266		0.5628		0.7709	
Midwest	0.4131		0.5327		0.7563	
East North Central	0.3587		0.3553		0.5630	
West North Central	0.9369		0.8020		0.8132	
South	0.1452		0.3041		0.6127	
South Atlantic	0.6628		0.9123		0.4890	
East South Central	0.5099		0.7333		0.9768	
West South Central	0.1114		0.0604		0.9572	
West	0.9949		0.0585		0.0320	
Mountain	0.6475		0.1410		0.1867	
Pacific	0.8070		0.1674		0.0810	
COUNTY TYPE						
Large Metro	0.6486		0.7372		0.2778	
Small Metro	0.2951		0.5162		0.3214	
250K – 1 Mil. Pop.	0.9150		0.3980		0.5594	
< 250K Pop.	0.0450		0.9621		0.3649	
Nonmetro	0.0589		0.0432		0.2203	
Urbanized	0.0589		0.0156		0.6153	
Less Urbanized	0.5802		0.6375		0.2988	
Completely Rural	0.5508		0.5672		0.5397	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.