

**Table 4.13B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2010 and 2011**

Substance	Total (2010)	Total (2011)	Male (2010)	Male (2011)	Female (2010)	Female (2011)
<b>ILLCIT DRUGS<sup>1,2</sup></b>	20.6	19.6	19.1	17.4	21.9	21.4
Marijuana and Hashish	19.2	18.5	17.9	17.3	20.6	19.7
Cocaine	21.2	20.1	21.6	19.7	20.5	20.6
Crack	24.8	20.8	22.8	20.6	26.1	21.0
Heroin	21.4	22.1	22.4	22.9	19.9	21.2
Hallucinogens	18.3	18.7	18.3	19.0	18.3	18.4
LSD	19.0	18.6	18.7	18.7	19.4	18.5
PCP	17.6	17.8	18.1	18.9	16.9	16.2
Ecstasy	19.8	20.4	19.6	21.1	20.0	19.7
Inhalants	17.2	16.6	18.3	17.6	16.3	15.7
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	24.5	23.5	24.1 <sup>a</sup>	20.8	24.8	25.3
Pain Relievers	22.9	23.4	22.5	20.5	23.2	25.3
OxyContin <sup>®</sup>	24.8	23.9	23.3	24.2	26.5	23.5
Tranquilizers	26.7	25.6	24.6	22.7	28.1	27.3
Stimulants <sup>2</sup>	21.2	22.7	20.1	21.2	22.1	23.8
Sedatives	30.9	25.5	34.1	20.4	28.2	28.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1,2</sup></b>	21.6	21.0	20.8	19.1	22.3	22.5
<b>CIGARETTES</b>	17.4	17.4	17.1	17.6	17.7	17.2
Daily Cigarette Use <sup>4</sup>	19.1	19.1	18.6	19.2	19.8	19.0
<b>SMOKELESS TOBACCO<sup>5</sup></b>	20.4	20.1	20.6	20.4	19.9	18.9
<b>CIGARS</b>	22.4	23.0	20.6	23.8	24.5	21.9
<b>ALCOHOL</b>	18.0	17.3	18.7	16.9	17.4	17.6

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

<sup>5</sup> Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.