

Table 4.14B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2010 and 2011

Substance	Total (2010)	Total (2011)	Male (2010)	Male (2011)	Female (2010)	Female (2011)
ILLICIT DRUGS^{1,2}	14.9	14.9	15.0	15.0	14.8	14.8
Marijuana and Hashish	15.2	15.1	15.2	15.2	15.2	15.0
Cocaine	15.7	15.8	15.9	15.9	15.5	15.7
Crack	16.1	15.9	*	16.2	16.2	15.7
Heroin	15.3	15.1	*	15.9	15.3	14.1
Hallucinogens	15.5 ^a	15.7	15.6 ^a	15.9	15.4	15.5
LSD	15.5	15.7	15.4	15.9	15.6	15.3
PCP	15.2	14.9	15.7	15.1	14.9	14.7
Ecstasy	15.6	15.8	15.6 ^a	16.0	15.6	15.6
Inhalants	14.1	14.3	14.2	14.5	13.9	14.1
Nonmedical Use of Psychotherapeutics ^{2,3}	15.2	15.2	15.4	15.5	15.0	15.0
Pain Relievers	15.2	15.3	15.4	15.5	15.0	15.0
OxyContin [®]	15.6	15.5	15.8	15.5	15.4	15.5
Tranquilizers	15.6	15.4	15.8	15.5	15.4	15.2
Stimulants ²	15.5 ^a	15.1	15.7	15.5	15.4 ^a	14.8
Sedatives	14.8	15.0	14.8	15.5	14.8	14.7
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	14.8	15.0	15.1	15.2	14.6	14.8
CIGARETTES	15.1	15.1	15.2	15.1	15.1	15.1
Daily Cigarette Use ⁴	15.7	15.8	15.8	15.9	15.5	15.6
SMOKELESS TOBACCO⁵	15.2	15.3	15.2	15.2	15.3	15.6
CIGARS	15.4	15.4	15.4	15.4	15.3	15.5
ALCOHOL	15.0	15.0	15.1	15.1	14.9	14.9

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 4.15B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2010 and 2011

Substance	Total (2010)	Total (2011)	Male (2010)	Male (2011)	Female (2010)	Female (2011)
ILLCIT DRUGS^{1,2}	15.9	16.0	15.9	16.1	15.8	15.9
Marijuana and Hashish	16.2	16.2	16.2	16.2	16.2	16.2
Cocaine	17.3 ^a	17.8	17.6 ^a	18.1	16.9	17.4
Crack	17.3	17.3	17.6	17.8	17.0	16.8
Heroin	17.0	17.2	17.4	17.7	16.6	16.5
Hallucinogens	16.8	17.1	17.1	17.2	16.5 ^a	16.9
LSD	17.3	17.1	17.5	17.4	17.0	16.7
PCP	16.8 ^b	15.1	17.8 ^b	15.1	15.4	15.0
Ecstasy	17.0	17.1	17.3	17.2	16.6	17.0
Inhalants	15.1	15.3	15.4	15.7	14.8	14.9
Nonmedical Use of Psychotherapeutics ^{2,3}	16.5	16.5	16.8	16.7	16.3	16.4
Pain Relievers	16.5	16.4	16.7	16.6	16.2	16.3
OxyContin [®]	16.9	16.7	17.0	16.8	16.7	16.6
Tranquilizers	17.3 ^a	16.8	17.6	17.1	17.0	16.6
Stimulants ²	17.1	17.0	17.5	17.5	16.6	16.6
Sedatives	16.4	16.1	16.8	16.9	16.2	15.6
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	16.2	16.3	16.4	16.4	15.9	16.2
CIGARETTES	16.3	16.4	16.4	16.3	16.2	16.5
Daily Cigarette Use ⁴	17.2	17.2	17.3	17.3	17.1	17.1
SMOKELESS TOBACCO⁵	16.5	16.6	16.5	16.6	16.7	16.7
CIGARS	16.8	16.8	16.8	16.7	16.9	16.8
ALCOHOL	16.0	15.9	16.1	16.0	16.0	15.9

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.