

Table 6.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	5,368	4,962	7,383	7,285	4,949	5,022	4,947	5,185
Marijuana and Hashish	4,573	4,149	5,897	6,302	3,849	4,106	3,090	3,513
Cocaine	648	646	498	479	156	125	170	120
Crack	112	86	124	112	70 ^a	5	72	26
Heroin	51	53	47	65	70	51	71	112
Hallucinogens	543 ^b	314	402	370	130	160	132	128
LSD	77	56	48	55	16	15	13	25
PCP	4	6	8	18	*	*	24	*
Ecstasy	303 ^a	187	236	205	90	108	62	44
Inhalants	111	133	163	152	205	110	211	229
Nonmedical Use of Psychotherapeutics ^{3,4}	1,575	1,416	2,060 ^a	1,624	1,428	1,215	1,894	1,864
Pain Relievers	1,220 ^a	954	1,494	1,241	1,001	828	1,378	1,448
OxyContin [®]	136	126	193	161	90	49	147	98
Tranquilizers	458	499	677	529	512	395	512	418
Stimulants ⁴	373	359	338	232	133	150	238	228
Methamphetamine ⁴	72	155	115	60	66	72	100	152
Sedatives	72	48	127	32	30	49	146	101
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,323	2,025	2,714 ^a	2,270	1,715	1,484	2,265	2,241

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	31.8	31.3	17.7	17.2	6.8	6.7	4.0	4.2
Marijuana and Hashish	27.1	26.2	14.2	14.8	5.3	5.5	2.5	2.8
Cocaine	3.8	4.1	1.2	1.1	0.2	0.2	0.1	0.1
Crack	0.7	0.5	0.3	0.3	0.1 ^a	0.0	0.1	0.0
Heroin	0.3	0.3	0.1	0.2	0.1	0.1	0.1	0.1
Hallucinogens	3.2 ^b	2.0	1.0	0.9	0.2	0.2	0.1	0.1
LSD	0.5	0.4	0.1	0.1	0.0	0.0	0.0	0.0
PCP	0.0	0.0	0.0	0.0	*	*	0.0	*
Ecstasy	1.8	1.2	0.6	0.5	0.1	0.1	0.1	0.0
Inhalants	0.7	0.8	0.4	0.4	0.3	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	9.3	8.9	4.9 ^b	3.8	2.0	1.6	1.5	1.5
Pain Relievers	7.2	6.0	3.6 ^a	2.9	1.4	1.1	1.1	1.2
OxyContin [®]	0.8	0.8	0.5	0.4	0.1	0.1	0.1	0.1
Tranquilizers	2.7	3.1	1.6	1.2	0.7	0.5	0.4	0.3
Stimulants ⁴	2.2	2.3	0.8	0.5	0.2	0.2	0.2	0.2
Methamphetamine ⁴	0.4	1.0	0.3	0.1	0.1	0.1	0.1	0.1
Sedatives	0.4	0.3	0.3	0.1	0.0	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	13.7	12.8	6.5 ^a	5.3	2.4	2.0	1.9	1.8

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	287	267	737	658	364	441	1,079	1,157
Marijuana and Hashish	254	250	659	589	294 ^b	397	595 ^a	731
Cocaine	22	22	22	21	10	14	7	10
Crack	3	0	0	3	*	*	*	*
Heroin	2	1	3	2	1	10	2	1
Hallucinogens	51	46	83	76	30	35	50	58
LSD	17	8	15	21	7	1	6	4
PCP	2	2	2	2	*	*	3	*
Ecstasy	22	31	56	37	21	23	25	17
Inhalants	24	13	35	43	34	20	169	147
Nonmedical Use of Psychotherapeutics ^{3,4}	85	94	201	176	69	81	366	338
Pain Relievers	77	77	166	140	56	53	308	293
OxyContin [®]	13	17	18	23	2	2	14	9
Tranquilizers	28	42	44	62	10	23	51	35
Stimulants ⁴	23	29	43	21	10	22	32	25
Methamphetamine ⁴	5	12	11	9	2	8	6	7
Sedatives	5	2	10	7	*	5	20	9
ILLCIT DRUGS OTHER THAN MARIJUANA²	127	119	288	257	124	127	554	522

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	70.2	70.4	49.1	44.7	25.9	29.8	5.1	5.3
Marijuana and Hashish	62.1	66.1	43.8	40.1	20.9 ^b	26.9	2.8 ^a	3.4
Cocaine	5.3	5.8	1.4	1.4	0.7	1.0	0.0	0.0
Crack	0.8	0.0	0.0	0.2	*	*	*	*
Heroin	0.5	0.3	0.2	0.1	0.1	0.7	0.0	0.0
Hallucinogens	12.6	12.1	5.5	5.2	2.2	2.4	0.2	0.3
LSD	4.2	2.0	1.0	1.4	0.5	0.1	0.0	0.0
PCP	0.4	0.4	0.1	0.1	*	*	0.0	*
Ecstasy	5.3	8.3	3.8	2.5	1.5	1.5	0.1	0.1
Inhalants	5.8	3.4	2.3	2.9	2.4	1.4	0.8	0.7
Nonmedical Use of Psychotherapeutics ^{3,4}	20.7	24.7	13.4	12.0	4.9	5.5	1.7	1.6
Pain Relievers	18.8	20.3	11.1	9.5	4.0	3.6	1.5	1.4
OxyContin [®]	3.2	4.4	1.2	1.6	0.2	0.1	0.1	0.0
Tranquilizers	6.8	11.2	2.9	4.2	0.7	1.6	0.2	0.2
Stimulants ⁴	5.6	7.6	2.8	1.5	0.7	1.5	0.2	0.1
Methamphetamine ⁴	1.3	3.1	0.7	0.6	0.2	0.6	0.0	0.0
Sedatives	1.2	0.6	0.7	0.5	*	0.4	0.1	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	31.1	31.5	19.2	17.5	8.8	8.6	2.6	2.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	2,251	2,019	2,746	2,903	1,257	1,391	1,091	1,038
Marijuana and Hashish	1,997	1,809	2,354 ^a	2,659	1,116	1,226	845	831
Cocaine	287	247	159	170	33	27	23	20
Crack	29 ^a	7	31	12	9	1	3	1
Heroin	30	21	38	33	13	31	9	22
Hallucinogens	318 ^b	198	227	229	84	79	44	52
LSD	56	47	33	34	10	8	6	4
PCP	3	1	3	10	*	1	*	*
Ecstasy	175 ^a	108	133	135	67	46	28	27
Inhalants	39	49	58	49	23	36	16	11
Nonmedical Use of Psychotherapeutics ^{3,4}	634	574	733 ^b	569	286	290	344	279
Pain Relievers	456	394	562 ^a	436	230	201	262 ^a	189
OxyContin [®]	55	52	105	61	25	27	53 ^b	18
Tranquilizers	203	182	224	173	89	105	54	85
Stimulants ⁴	170	166	149 ^a	95	32	42	63	41
Methamphetamine ⁴	15	25	47 ^a	17	11	14	10	23
Sedatives	20	14	16	9	4	12	19	8
ILLCIT DRUGS OTHER THAN MARIJUANA²	998 ^a	824	954	840	368	378	389	342

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	48.8	48.5	29.9	30.6	17.7	19.4	8.3	7.7
Marijuana and Hashish	43.3	43.4	25.6 ^a	28.0	15.7	17.1	6.4	6.2
Cocaine	6.2	5.9	1.7	1.8	0.5	0.4	0.2	0.2
Crack	0.6 ^a	0.2	0.3	0.1	0.1	0.0	0.0	0.0
Heroin	0.6	0.5	0.4	0.3	0.2	0.4	0.1	0.2
Hallucinogens	6.9 ^a	4.7	2.5	2.4	1.2	1.1	0.3	0.4
LSD	1.2	1.1	0.4	0.4	0.1	0.1	0.0	0.0
PCP	0.1	0.0	0.0	0.1	*	0.0	*	*
Ecstasy	3.8	2.6	1.4	1.4	0.9	0.6	0.2	0.2
Inhalants	0.8	1.2	0.6	0.5	0.3	0.5	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	13.7	13.8	8.0 ^b	6.0	4.0	4.0	2.6	2.1
Pain Relievers	9.9	9.5	6.1 ^b	4.6	3.2	2.8	2.0 ^a	1.4
OxyContin [®]	1.2	1.2	1.1 ^a	0.6	0.4	0.4	0.4 ^b	0.1
Tranquilizers	4.4	4.4	2.4	1.8	1.3	1.5	0.4	0.6
Stimulants ⁴	3.7	4.0	1.6 ^a	1.0	0.4	0.6	0.5	0.3
Methamphetamine ⁴	0.3	0.6	0.5 ^a	0.2	0.2	0.2	0.1	0.2
Sedatives	0.4	0.3	0.2	0.1	0.1	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	21.6	19.8	10.4 ^a	8.8	5.2	5.3	3.0	2.5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	2,831	2,676	3,899	3,724	3,328	3,190	2,777	2,990
Marijuana and Hashish	2,322	2,090	2,884	3,053	2,439	2,483	1,650	1,952
Cocaine	340	377	317	288	113	84	140	89
Crack	80	79	93	97	61	4	69	24
Heroin	19	32	6	31	56	10	60	88
Hallucinogens	174	70	92	65	17	46	38	18
LSD	4	1	*	*	*	7	*	17
PCP	*	3	3	7	*	*	21	*
Ecstasy	106	48	47	33	*	39	10	*
Inhalants	49	71	70	60	148	54	26	71
Nonmedical Use of Psychotherapeutics ^{3,4}	857	749	1,126	879	1,072	845	1,184	1,247
Pain Relievers	687	483	765	665	715	574	808	966
OxyContin [®]	67	57	70	77	62	20	81	71
Tranquilizers	227	274	409	294	412	267	407	298
Stimulants ⁴	180	165	146	116	91	86	143	162
Methamphetamine ⁴	51	118	56	34	52	49	85	122
Sedatives	47	32	101	17	25	32	108	84
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,198	1,081	1,472	1,173	1,223	978	1,322	1,376

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
ILLCIT DRUGS²	23.8	23.6	12.6	11.8	5.2	4.8	3.2	3.4
Marijuana and Hashish	19.6	18.5	9.3	9.7	3.8	3.7	1.9	2.2
Cocaine	2.9	3.3	1.0	0.9	0.2	0.1	0.2	0.1
Crack	0.7	0.7	0.3	0.3	0.1	0.0	0.1	0.0
Heroin	0.2	0.3	0.0	0.1	0.1	0.0	0.1	0.1
Hallucinogens	1.5	0.6	0.3	0.2	0.0	0.1	0.0	0.0
LSD	0.0	0.0	*	*	*	0.0	*	0.0
PCP	*	0.0	0.0	0.0	*	*	0.0	*
Ecstasy	0.9	0.4	0.2	0.1	*	0.1	0.0	*
Inhalants	0.4	0.6	0.2	0.2	0.2	0.1	0.0	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	7.2	6.6	3.6	2.8	1.7	1.3	1.3	1.4
Pain Relievers	5.8	4.3	2.5	2.1	1.1	0.9	0.9	1.1
OxyContin [®]	0.6	0.5	0.2	0.2	0.1	0.0	0.1	0.1
Tranquilizers	1.9	2.4	1.3	0.9	0.6	0.4	0.5	0.3
Stimulants ⁴	1.5	1.5	0.5	0.4	0.1	0.1	0.2	0.2
Methamphetamine ⁴	0.4	1.0	0.2	0.1	0.1	0.1	0.1	0.1
Sedatives	0.4	0.3	0.3	0.1	0.0	0.0	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	10.1	9.6	4.8	3.7	1.9	1.5	1.5	1.5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2010 and 2011

Drug	Total (2010)	Total (2011)	Aged 12-20 (2010)	Aged 12-20 (2011)	Aged 21+ (2010)	Aged 21+ (2011)
ILLICIT DRUGS¹	17,701	17,269	3,896	3,985	13,805	13,284
Marijuana or Hashish	14,318	14,557	3,464	3,674	10,854	10,884
Cocaine (Including Crack)	1,301	1,250	204	253	1,097	997
Heroin	169	169	45	43	124	126
Hallucinogens	1,075 ^a	844	455	381	620	463
Inhalants	479	395	134	146	346	249
Pain Relievers ²	3,714 ^b	3,023	774	653	2,941 ^b	2,370
Tranquilizers ²	1,647	1,423	260	328	1,387	1,095
Stimulants ^{2,3}	844	742	224	194	620	548
Methamphetamine ³	253	287	56	47	197	240
Sedatives ²	228	130	27	30	201	100
ILLICIT DRUGS OTHER THAN MARIJUANA¹	6,751 ^b	5,779	1,425	1,326	5,326 ^b	4,453

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2010 and 2011

Drug	Total (2010)	Total (2011)	Aged 12-20 (2010)	Aged 12-20 (2011)	Aged 21+ (2010)	Aged 21+ (2011)
ILLCIT DRUGS¹	13.5	12.9	39.2	41.3	11.4	10.7
Marijuana or Hashish	10.9	10.9	34.9 ^b	38.1	8.9	8.8
Cocaine (Including Crack)	1.0	0.9	2.1	2.6	0.9	0.8
Heroin	0.1	0.1	0.5	0.4	0.1	0.1
Hallucinogens	0.8 ^a	0.6	4.6	4.0	0.5 ^a	0.4
Inhalants	0.4	0.3	1.3	1.5	0.3	0.2
Pain Relievers ²	2.8 ^b	2.3	7.8	6.8	2.4 ^b	1.9
Tranquilizers ²	1.3	1.1	2.6 ^a	3.4	1.1 ^a	0.9
Stimulants ^{2,3}	0.6	0.6	2.3	2.0	0.5	0.4
Methamphetamine ³	0.2	0.2	0.6	0.5	0.2	0.2
Sedatives ²	0.2	0.1	0.3	0.3	0.2	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.1 ^b	4.3	14.3	13.7	4.4 ^b	3.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Numbers in Thousands, 2010 and 2011

Drug¹	Total (2010)	Total (2011)	Aged 12-20 (2010)	Aged 12-20 (2011)	Aged 21+ (2010)	Aged 21+ (2011)
ILLICIT DRUGS²	8,066	7,929	1,857	1,904	6,209	6,025
Marijuana or Hashish	7,215	7,276	1,761	1,812	5,454	5,464
Cocaine (Including Crack)	304	279	48	52	256	227
Heroin	62	55	15	14	47	42
Hallucinogens	99	98	41	30	58	68
Inhalants	36	35	15	17	20	18
Pain Relievers ³	591	470	99	78	492	392
Tranquilizers ³	170 ^a	84	35	26	134 ^a	58
Stimulants ³	103	66	13 ^a	4	90	63
Methamphetamine ⁴	85	80	5	6	80	74
Sedatives ³	15	35	3	3	12	31
ILLICIT DRUGS OTHER THAN MARIJUANA²	1,230	1,081	213	210	1,017	871

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Percentages, 2010 and 2011

Drug¹	Total (2010)	Total (2011)	Aged 12-20 (2010)	Aged 12-20 (2011)	Aged 21+ (2010)	Aged 21+ (2011)
ILLCIT DRUGS²	6.2	6.0	19.0	20.1	5.1	4.9
Marijuana or Hashish	5.5	5.5	18.1	19.2	4.5	4.4
Cocaine (Including Crack)	0.2	0.2	0.5	0.5	0.2	0.2
Heroin	0.0	0.0	0.2	0.1	0.0	0.0
Hallucinogens	0.1	0.1	0.4	0.3	0.0	0.1
Inhalants	0.0	0.0	0.2	0.2	0.0	0.0
Pain Relievers ³	0.5	0.4	1.0	0.8	0.4	0.3
Tranquilizers ³	0.1 ^a	0.1	0.4	0.3	0.1 ^a	0.0
Stimulants ³	0.1	0.0	0.1 ^a	0.0	0.1	0.1
Methamphetamine ⁴	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives ³	0.0	0.0	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.9	0.8	2.2	2.2	0.8	0.7

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.17A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2010 and 2011

Drug	Total (2010)	Total (2011)	Aged 12-17 (2010)	Aged 12-17 (2011)	Aged 18-25 (2010)	Aged 18-25 (2011)	Aged 26+ (2010)	Aged 26+ (2011)
ILLCIT DRUGS¹	17,701	17,269	1,389	1,365	6,254	6,314	10,058	9,590
Marijuana or Hashish	14,318	14,557	1,207	1,237	5,467	5,695	7,644	7,626
Cocaine (Including Crack)	1,301	1,250	53	57	479	444	769	749
Heroin	169	169	6	13	81	85	81	72
Hallucinogens	1,075 ^a	844	165	157	628 ^a	505	282	182
Inhalants	479	395	93	76	119	133	267	186
Pain Relievers ²	3,714 ^b	3,023	299	270	1,248 ^b	1,031	2,167 ^a	1,722
Tranquilizers ²	1,647	1,423	82 ^a	127	516	461	1,049	834
Stimulants ^{2,3}	844	742	76	73	351	303	418	366
Methamphetamine ³	253	287	19	29	74	56	160	201
Sedatives ²	228	130	15	15	41	35	172	80
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6,751 ^b	5,779	538	503	2,320 ^a	2,042	3,892 ^a	3,233

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.17B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2010 and 2011

Drug	Total (2010)	Total (2011)	Aged 12-17 (2010)	Aged 12-17 (2011)	Aged 18-25 (2010)	Aged 18-25 (2011)	Aged 26+ (2010)	Aged 26+ (2011)
ILLCIT DRUGS¹	13.5	12.9	41.8	41.0	29.9	30.3	9.4	8.8
Marijuana or Hashish	10.9	10.9	36.4	37.2	26.1	27.4	7.1	7.0
Cocaine (Including Crack)	1.0	0.9	1.6	1.7	2.3	2.1	0.7	0.7
Heroin	0.1	0.1	0.2	0.4	0.4	0.4	0.1	0.1
Hallucinogens	0.8 ^a	0.6	5.0	4.7	3.0 ^a	2.4	0.3	0.2
Inhalants	0.4	0.3	2.8	2.3	0.6	0.6	0.2	0.2
Pain Relievers ²	2.8 ^b	2.3	9.0	8.1	6.0 ^b	5.0	2.0 ^a	1.6
Tranquilizers ²	1.3	1.1	2.5 ^a	3.8	2.5	2.2	1.0	0.8
Stimulants ^{2,3}	0.6	0.6	2.3	2.2	1.7	1.5	0.4	0.3
Methamphetamine ³	0.2	0.2	0.6	0.9	0.4	0.3	0.1	0.2
Sedatives ²	0.2	0.1	0.5	0.4	0.2	0.2	0.2	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.1 ^b	4.3	16.2	15.1	11.1 ^a	9.8	3.6 ^a	3.0

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2010 and 2011

Drug¹	Total (2010)	Total (2011)	Aged 12-17 (2010)	Aged 12-17 (2011)	Aged 18-25 (2010)	Aged 18-25 (2011)	Aged 26+ (2010)	Aged 26+ (2011)
ILLCIT DRUGS²	8,066	7,929	628	619	3,170	3,162	4,268	4,147
Marijuana or Hashish	7,215	7,276	607	582	2,957	2,993	3,651	3,701
Cocaine (Including Crack)	304	279	9	12	118	95	177	171
Heroin	62	55	*	5	32	29	31	22
Hallucinogens	99	98	14	11	56	62	29	25
Inhalants	36	35	12	1	15	16	9	18
Pain Relievers ³	591	470	32	31	182	161	377	277
Tranquilizers ³	170 ^a	84	9	10	56	40	105 ^a	34
Stimulants ³	103	66	3	2	50	23	50	41
Methamphetamine ⁴	85	80	2	6	13	5	70	70
Sedatives ³	15	35	3	3	2	*	10	30
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,230	1,081	66	71	401	393	763	617

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2010 and 2011

Drug¹	Total (2010)	Total (2011)	Aged 12-17 (2010)	Aged 12-17 (2011)	Aged 18-25 (2010)	Aged 18-25 (2011)	Aged 26+ (2010)	Aged 26+ (2011)
ILLCIT DRUGS²	6.2	6.0	19.4	19.2	15.3	15.3	4.0	3.8
Marijuana or Hashish	5.5	5.5	18.8	18.2	14.3	14.5	3.4	3.4
Cocaine (Including Crack)	0.2	0.2	0.3	0.4	0.6	0.5	0.2	0.2
Heroin	0.0	0.0	*	0.1	0.2	0.1	0.0	0.0
Hallucinogens	0.1	0.1	0.4	0.3	0.3	0.3	0.0	0.0
Inhalants	0.0	0.0	0.4	0.0	0.1	0.1	0.0	0.0
Pain Relievers ³	0.5	0.4	1.0	1.0	0.9	0.8	0.4	0.3
Tranquilizers ³	0.1 ^a	0.1	0.3	0.3	0.3	0.2	0.1 ^a	0.0
Stimulants ³	0.1	0.0	0.1	0.1	0.2	0.1	0.0	0.0
Methamphetamine ⁴	0.1	0.1	0.1	0.2	0.1	0.0	0.1	0.1
Sedatives ³	0.0	0.0	0.1	0.1	0.0	*	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.9	0.8	2.0	2.2	1.9	1.9	0.7	0.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.