

Table 6.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	11,327	11,406
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,600	3,256	5,279	4,987
Cigars	7,204	6,691	6,069	6,174
Pipe Tobacco	1,180	1,149	974	988
ALCOHOL	38,575	37,764	92,799	95,621
Binge Alcohol Use ²	25,484	24,154	33,051	34,187
Heavy Alcohol Use ²	9,231	8,716	7,668	7,148

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	5.8	5.7
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	6.2	5.7	2.7	2.5
Cigars	12.3	11.8	3.1	3.1
Pipe Tobacco	2.0	2.0	0.5	0.5
ALCOHOL	66.1	66.5	47.5	47.6
Binge Alcohol Use ²	43.7	42.5	16.9	17.0
Heavy Alcohol Use ²	15.8	15.3	3.9	3.6

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	572	551
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	286	278	284	239
Cigars	479	549	310	309
Pipe Tobacco	111	120	44	62
ALCOHOL	1,223	1,171	2,096	2,156
Binge Alcohol Use ²	945	864	967	985
Heavy Alcohol Use ²	285	249	124	129

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	2.6	2.4
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	14.0	14.3	1.3	1.0
Cigars	23.5 ^b	28.3	1.4	1.3
Pipe Tobacco	5.4	6.2	0.2	0.3
ALCOHOL	59.9	60.4	9.4	9.4
Binge Alcohol Use ²	46.3	44.6	4.3	4.3
Heavy Alcohol Use ²	14.0	12.9	0.6	0.6

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	2,232	2,044
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,310	1,159	873 ^a	708
Cigars	2,438	2,439	1,420	1,285
Pipe Tobacco	431	399	188	236
ALCOHOL	9,143	8,880	11,774	11,934
Binge Alcohol Use ²	7,246	6,948	6,550	6,705
Heavy Alcohol Use ²	2,950 ^a	2,646	1,667	1,519

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	10.0 ^a	9.0
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	11.2	10.1	3.9 ^b	3.1
Cigars	20.8	21.2	6.3	5.6
Pipe Tobacco	3.7	3.5	0.8	1.0
ALCOHOL	78.1	77.3	52.6	52.3
Binge Alcohol Use ²	61.9	60.5	29.3	29.4
Heavy Alcohol Use ²	25.2 ^a	23.0	7.4	6.7

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	8,524	8,810
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	2,004	1,819	4,122	4,040
Cigars	4,287	3,702	4,340	4,581
Pipe Tobacco	639	630	742	690
ALCOHOL	28,210	27,713	78,928	81,532
Binge Alcohol Use ²	17,292	16,342	25,535	26,496
Heavy Alcohol Use ²	5,996	5,821	5,877	5,500

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2010 and 2011

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
TOBACCO PRODUCTS¹	N/A	N/A	5.7	5.7
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.5	4.2	2.7	2.6
Cigars	9.6	8.5	2.9	3.0
Pipe Tobacco	1.4	1.5	0.5	0.4
ALCOHOL	63.3	63.9	52.4	52.6
Binge Alcohol Use ²	38.8	37.7	17.0	17.1
Heavy Alcohol Use ²	13.4	13.4	3.9	3.5

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.